

Original article

Cost-effectiveness of implantable cardioverter-defibrillators for primary prevention of sudden cardiac death



Aida Ribera,^{a,b,*} Emmanuel Giménez,^c Gerard Oristrell,^{d,e} Dimelza Osorio,^{b,f} Josep Ramón Marsal,^{a,b} Lidia García-Pérez,^{g,h,i} Mónica Ballesteros,^{b,f} Eduard Ródenas,^d Yassin Belahnech,^d Roxana Escalona,^d Núria Rivas,^d Ivo Roca-Luque,^j Ignacio Ferreira-González,^{a,b,d} and Mireia Espallargues^{c,i}

^a Unidad de Epidemiología Cardiovascular, Servicio de Cardiología, Hospital Universitario Vall d'Hebron, Institut de Recerca Vall d'Hebron (VHIR), Barcelona, Spain

^b Centro de Investigación en Red de Epidemiología y Salud Pública (CIBERESP), Spain

^c Agència de Qualitat i Avaluació Sanitàries de Catalunya (AQuAS), Departament de Salut, Barcelona, Spain

^d Servicio de Cardiología, Hospital Universitario Vall d'Hebron, Barcelona, Spain

^e Centro de Investigación en Red de Enfermedades Cardiovasculares (CIBERCV), Spain

^f Unidad de Calidad-Mejora de la Práctica Clínica, Dirección de Procesos y Calidad, Hospital Universitario Vall d'Hebron, Barcelona, Spain

^g Servicio de Evaluación del Servicio Canario de la Salud (SESCS), Tenerife, Spain

^h Red Española de Agencias de Evaluación de Tecnologías Sanitarias y Prestaciones del Sistema Nacional de Salud (RedETS), Spain

ⁱ Red de Investigación en Servicios de Salud en Enfermedades Crónicas (REDISSEC), Madrid, Spain

^j Unidad de Arritmias, Institut Cardiovascular, Hospital Clínic de Barcelona, Barcelona, Spain

Article history:

Received 27 November 2020

Accepted 23 February 2021

Available online 5 June 2021

Keywords:

Implantable cardioverter defibrillator

Cost-effectiveness

Primary prevention

ABSTRACT

Introduction and objectives: Implantable cardioverter-defibrillators (ICD) are a cost-effective alternative for secondary prevention of sudden cardiac death, but their efficiency in primary prevention, especially among patients with nonischemic heart disease, is still uncertain.

Methods: We performed a cost-effectiveness analysis of ICD plus conventional medical treatment (CMT) vs CMT for primary prevention of cardiac arrhythmias from the perspective of the national health service. We simulated the course of the disease by using Markov models in patients with ischemic and nonischemic heart disease. The parameters of the model were based on the results obtained from a meta-analysis of clinical trials published between 1996 and 2018 comparing ICD plus CMT vs CMT, the safety results of the DANISH trial, and analysis of real-world clinical practice in a tertiary hospital.

Results: We estimated that ICD reduced the likelihood of all-cause death in patients with ischemic heart disease (HR, 0.70; 95%CI, 0.58–0.85) and in those with nonischemic heart disease (HR, 0.79; 95%CI, 0.66–0.96). The incremental cost-effectiveness ratio (ICER) estimated with probabilistic analysis was €19 171/quality adjusted life year (QALY) in patients with ischemic heart disease and €31 084/QALY in those with nonischemic dilated cardiomyopathy overall and €23 230/QALY in patients younger than 68 years.

Conclusions: The efficiency of single-lead ICD systems has improved in the last decade, and these devices are cost-effective in patients with ischemic and nonischemic left ventricular dysfunction younger than 68 years, assuming willingness to pay as €25 000/QALY. For older nonischemic patients, the ICER was around €30 000/QALY.

© 2021 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Coste-efectividad del desfibrilador automático implantable para la prevención primaria de la muerte súbita cardíaca

RESUMEN

Introducción y objetivos: El desfibrilador automático implantable (DAI) es una alternativa coste-efectiva para la prevención secundaria de la muerte súbita cardíaca, pero sigue habiendo dudas sobre su eficiencia en prevención primaria, sobre todo en pacientes con cardiopatía no isquémica.

Métodos: Análisis de coste-utilidad del DAI más tratamiento médico convencional frente a tratamiento médico convencional para la prevención primaria de arritmias cardíacas desde la perspectiva del Sistema Nacional de Salud. Se simuló el curso de la enfermedad mediante modelos de Markov en pacientes con y sin cardiopatía isquémica. Los parámetros del modelo se basaron en los resultados obtenidos mediante metanálisis de los ensayos clínicos publicados entre 1996 y 2018 en los que se comparaba el DAI con el

Palabras clave:

Desfibrilador automático implantable

Coste-efectividad

Prevención primaria

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2021.07.022>

* Corresponding author: Unidad de Epidemiología Cardiovascular, Servicio de Cardiología, Hospital Universitario Vall d'Hebron, Pg. Vall d'Hebron 119-129, 08035 Barcelona, Spain.

E-mail address: aida.ribera@vhir.org (A. Ribera).

<https://doi.org/10.1016/j.rec.2021.05.004>

1885-5857/© 2021 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

tratamiento médico convencional, los resultados de seguridad del ensayo DANISH y el análisis de la práctica clínica habitual en un hospital terciario.

Resultados: Se estimó un beneficio del DAI sobre la muerte por cualquier causa con HR = 0,70 (IC95%, 0,58–0,85) en cardiopatía isquémica y HR = 0,79 (IC95%, 0,66–0,96) en no isquémica. La razón de coste-efectividad incremental estimada mediante análisis probabilístico fue de 19.171 euros/año de vida ajustado por calidad (AVAC) en pacientes con cardiopatía isquémica, 31.084 euros/AVAC en pacientes con miocardiopatía dilatada no isquémica y 23.230 euros/AVAC en los menores de 68 años.

Conclusiones: La eficiencia del DAI monocameral ha mejorado en la última década y este resulta coste-efectivo para los pacientes con disfunción ventricular izquierda de origen isquémico o no isquémico menores de 68 años considerando una disposición a pagar 25.000 euros/AVAC. En pacientes no isquémicos mayores, la razón de coste-efectividad incremental estimada se sitúa alrededor de los 30.000 euros/AVAC.

© 2021 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Abbreviations

CMT: conventional medical treatment
EF: ejection fraction
ICD: implantable cardioverter-defibrillator
ICER: incremental cost-effectiveness ratio
NHS: National Health Service
QALY: quality-adjusted life year

INTRODUCTION

The implantable cardioverter-defibrillator (ICD) is an effective device used to treat life-threatening ventricular arrhythmia.¹ More than a decade after the publication of the earliest pivotal trials,^{2,3} the efficacy of ICD for primary prevention of sudden cardiac death has been a subject of debate due to the publication of the last clinical trial performed in patients with nonischemic heart failure.⁴ This debate led to the publication of numerous systematic reviews and meta-analyses, concluding that ICD in primary prevention improves the survival of patients with ischemic^{5,6} and nonischemic^{6–17} heart failure and ejection fraction (EF) $\leq 35\%$ compared with conventional medical treatment (CMT).

According to the official registry published in *Revista Española de Cardiología*,¹⁸ the prophylactic use of ICD accounts for 62% of all indications, with high variability between hospitals, and is the fastest-growing indication for patients with dilated cardiomyopathy: 63.5% of indications vs 49.9% in patients with ischemic heart disease.

Health technology assessment is a basic tool used in decision-making and resource allocation to ensure high quality services and system sustainability.¹⁹ A technology review report prepared in our setting in 2011²⁰ concluded that ICDs were a cost-effective alternative for primary prevention in patients with ischemic dilated cardiomyopathy who met the criteria of the MADIT trial.²¹ Conversely, the indication of ICD in primary prevention was not cost-effective, assuming willingness to pay as €30 000/quality-adjusted life year (QALY), in patients who met the MADIT II² criteria and had nonischemic dilated cardiomyopathy.

There is evidence^{22,23} that the arrhythmia mortality rate and the rate of appropriate therapies have dropped in recent years and that they vary according to age and comorbidity. Consequently, many patients with ICD will never receive an appropriate discharge or will die due to a nonarrhythmic cause.²⁴ A secondary finding²⁵ of the DANISH study was that the analyses of predefined subgroups observed a positive effect of ICD in terms of reducing the risk of mortality among patients with nonischemic ventricular dysfunction who received an ICD at earlier ages. Therefore, it is

essential to select patients with a higher expected benefit of the ICD and a lower probability of complications to allow an efficient use of ICD, particularly in indications for primary prevention.

The aim of this study was to evaluate the cost-utility of ICD in primary prevention for patients receiving CMT, updating the 2011 review with any new evidence, and to determine in which patient subgroups it would be cost-effective according to age and type of heart disease.

METHODS

Cost-utility economic model based on a Markov decision analysis.

Decision model

A Markov model was used to estimate the costs and survival of patients on CMT who receive an ICD for primary prevention and those who do not. A cost-utility analysis was performed from the perspective of the national health service, using patient lifetime as the time frame and applying a discount rate of 3% per year for both costs and benefits. The analysis considered CMT vs ICD plus CMT for primary prevention in patients with 3 different profiles: a) ischemic ventricular dysfunction; b) nonischemic ventricular dysfunction, and c) nonischemic ventricular dysfunction in patients age 68 years or less (predefined subgroup of the DANISH study seen to benefit from the ICD).

The model used in a previous evaluation²⁶ was adapted with the TreeAge Pro software (version 2019 R2.0). A hypothetical cohort was simulated of 60-year-old patients with functional class II–III heart failure and EF $< 35\%$, with or without a history of ischemic heart disease, who received a single-chamber ICD (CMT + ICD group) for primary prevention or who did not receive it (CMT group). Figure 1 shows the various states through which patients could move in monthly cycles and the transitions between them. All patients started with a “stable” state and, from there, were at particular risk of all-cause death or hospitalization for heart failure in each month. Additionally, patients in the CMT + ICD group could experience a device-related complication at any time or a perioperative complication in the first month or when the battery was replaced (assumed to be 8 years after the first implant).

Mortality

The likelihoods of death are shown for each group and each model in table 1. All likelihoods were obtained from a systematic review and meta-analysis on ICD efficacy in primary prevention (methodology described in supplementary data I), updating the




Figure 1. Markov model structure. Health states are shown as solid-line boxes, and transition conditions as dashed-line boxes. The arrows represent the transitions between states. The dashed arrow indicates that it starts from all health states. CMT, conventional medical treatment; HF, heart failure; ICD, implantable cardioverter-defibrillator.

search performed in a previous study²⁰ up to May 30, 2018. Based on the meta-analysis, the efficacy data from the MADIT,²¹ MADIT II,² and SCD-HeFT³ trials were used to estimate the likelihood of death among ischemic patients. However, the CABG-Patch,²⁶ DINAMIT,²⁷ and IRIS²⁸ studies were excluded because they took place in the setting of coronary revascularization surgery or recent myocardial infarction. The likelihood of death among patients with nonischemic ventricular dysfunction was estimated using efficacy data from the CAT,²⁹ DEFINITE,³⁰ DANISH,⁴ and SCD-HeFT³ studies (placebo-controlled, except for the amiodarone treatment arm). The estimate calculation excluded the AMIOVIRT study,³¹ as it used amiodarone as a comparator, as well as the Pro-ICD study,³² which included patients wait-listed for a heart transplant. The COMPANION trial was also excluded because efficacy data could not be collected separately for patients with and without ischemic heart disease.

The likelihoods were estimated from the cumulative rates of all randomized clinical trials included in both treatment groups, using the following formula:

$$1 - e^{-rt}$$

The effect of ICD on the likelihood of all-cause death was assumed to be constant over the entire period, even though the randomized clinical trials had a mean follow-up period of 16 to 41 months (table 1 of the supplementary data). To extrapolate mortality rates over the patient's entire lifetime, an age adjustment was performed using the 2017 mortality tables of the Spanish population published by the National Institute of Statistics (figure 4 of the supplementary data). All model parameters for the base case and for the sensitivity analyses are summarized in table 2.

Complications

Perioperative complications (superficial or deep infection, major bleeds, pneumothorax) were differentiated from those potentially occurring throughout follow-up (malfunction due to rupture or dislodgment or inappropriate therapies).

Complication probabilities were obtained from the recent literature. Because registries and observational studies tend to underestimate complication rates,³³ the perioperative complication rates and monthly rate of inappropriate therapies described in the DANISH trial were used, as this is the most recent randomized

Table 1
Baseline likelihoods of transition to all-cause death

	HR (95%CI)*	Monthly likelihood		Sources
		CMT+ICD group	CMT group	
Ischemic heart disease	0.70 (0.58-0.85)	0.0074	0.0110	MADIT, ²¹ MADIT II, ² SCD-HeFT ³
Nonischemic heart disease	0.79 (0.66-0.96)	0.0038	0.0049	CAT, ²⁹ DEFINITE, ³⁰ DANISH, ⁴ SCD-HeFT ³
DANISH patients younger than 68 years	0.70 (0.51-0.96)	0.0016	0.0031	DANISH ⁴

95%CI, 95% confidence interval; CAT, Cardiomyopathy Trial; CMT, conventional medical treatment; DANISH, DANish trial to assess efficacy of Implantable cardioverter defibrillators in patients with non-ischemic Systolic Heart failure; DEFINITE, Defibrillators in non-ischemic Cardiomyopathy Treatment Evaluation trial; HR, hazard ratio; ICD, implantable cardioverter defibrillator; MADIT, Multicenter Automatic Defibrillator Implantation Trial; SCD-HeFT, Sudden Cardiac Death in Heart Failure Trial.

* Summation effect obtained in the meta-analyses.

Table 2

Summary of the model parameters for the base case and sensitivity analyses

	Base case		Deterministic analysis		Probabilistic analysis		
	Value	Source	Value	Source	Function	Value	Source
Likelihoods of transition							
CMT + ICD group							
Annual (monthly) likelihood of all-cause death in ischemic heart disease	0.0847 (0.0074)	MADIT, ²¹ MADIT II, ² SCD-HeFT ³	± 20%	—	Beta	Alpha = 107 Beta = 1.159	MADIT, ²¹ MADIT II, ² SCD-HeFT ³
Annual (monthly) likelihood of all-cause death in nonischemic heart disease	0.0450 (0.0038)	CAT, ²⁹ DEFINITE, ³⁰ DANISH, ⁴ SCD-HeFT	± 20%	-	Beta	Alpha = 55 Beta = 1.177	CAT, ²⁹ DEFINITE, ³⁰ DANISH, ⁴ SCD-HeFT
Annual likelihood of all-cause death in DANISH patients younger than 68 y	0.0016	DANISH ⁴	± 20%	-	Beta	Alpha = 3 Beta = 163	DANISH ⁴
Likelihood of infection (45% superficial; 55% deep)	0.0026	DANISH ⁴	± 20%	-	-	-	-
Likelihood of major bleed	0.00003	DANISH ⁴	± 20%	-	-	-	-
Likelihood of pneumothorax	0.00029	DANISH ⁴	± 20%	-	-	-	-
Monthly likelihood of rupture, dislodgment, or displacement	0.0017	Ezzat et al. OpenHeart 2015 ³³	± 20%	-	-	-	-
Monthly likelihood of inappropriate therapy	0.0009	DANISH ⁴	± 20%	-	-	-	-
Monthly likelihood of hospitalization due to heart failure (also in CMT group)	0.0165	García-Pérez et al. ²⁰	± 20%	-	-	-	-
CMT group							
Annual (monthly) likelihood of all-cause death in ischemic heart disease	0.1245 (0.0110)	MADIT, ²¹ MADIT II, ² SCD-HeFT ³	± 20%	-	Beta	Alpha = 129 Beta = 913	
Annual (monthly) likelihood of all-cause death in nonischemic heart disease	0.0571 (0.0049)	CAT, ²⁹ DEFINITE, ³⁰ DANISH, ⁴ SCD-HeFT	± 20%	-	Beta	Alpha = 71 Beta = 1165	
Annual likelihood of all-cause death in DANISH patients younger than 68 y	0.0031	DANISH ⁴	± 20%	-	Beta	Alpha = 6 Beta = 174	
Utilities							
Death state	0	Assumed	-	-	-	-	-
Well state	0.75	Martín et al. ³⁶	± 20%		Beta	μ = 0.6 σ = 0.2	VIDA-IC study ⁴⁷
Hospitalized state due to heart failure or ICD complications	0.4	García-Pérez ³⁷	± 20%	-	Beta	μ = 0.4 σ = 0.1	
Resource utilization							
Generator replacement rate	8	VH analytical accounting	6 or 10	-	-	-	-
Cost of ICD, €	10 764		± 20%	-	-	-	-
Cost of ICD implantation, €	12 385		± 20%	-	-	-	-
Cost of generator replacement, €	11 864		-	-	-	-	-
Cost of emergency visits, €	208.16		-	-	-	-	-
Cost of superficial infection, €	24 382		-	-	-	-	-
Cost of deep infection, €	32 992				-	-	-
Cost of major bleed, €	12 940		-	-	-	-	-
Cost of pneumothorax, €	4315		-	-	-	-	-
Cost of follow-up, €	49.55 + 28.45		-	-	-	-	-
Cost of inappropriate therapies or reprogramming, €	208.16		180.86	Canary Islands	.		-
Cost of lead dislodgment or displacement, €	4167		-	-	-	-	-
Monthly cost of medical treatment for ischemic heart disease, €	28.48	Pharmacy Nomenclature of the Ministry of Health, Consumer Affairs, and Social Welfare	± 20%	-	-	-	-
Monthly cost of medical treatment for nonischemic heart disease, €	12.45		± 20%	-	-	-	-
Cost of hospitalization due to heart failure, €	4315	VH analytical accounting	± 20%	-	-	-	-
Other							
Proportion between superficial and deep infection	0.45	DANISH ⁴	-	-			
Discount rate	3%	Guidelines	0%; 5%				

CAT, Cardiomyopathy Trial; CMT, conventional medical treatment; DANISH, DANish trial to assess efficacy of Implantable cardioverter defibrillators in patients with non-ischemic Systolic Heart failure; DEFINITE, Defibrillators in non-ischemic Cardiomyopathy Treatment Evaluation trial; ICD, implantable cardioverter defibrillator; MADIT, Multicenter Automatic Defibrillator Implantation Trial; SCD-HeFT, Sudden Cardiac Death in Heart Failure Trial; VH, Vall d'Hebron Hospital.

The cost breakdown is listed in [table 4 of the supplementary data](#).

Table 3

Base case results (probabilistic analysis), CMT + ICD vs CMT

		Cost (90%CrI)	QALYs	ICER
Ischemic	CMT + ICD	27 125 (25 238–28 533)	5.12 (4.16–6.04)	
	CMT	4887 (3040–6974)	3.96 (3.25–4.66)	19 171 (15 622–24 393)
Nonischemic	CMT + ICD	34 252 (32 530–36 171)	7.26 (5.94–8.64)	
	CMT	7856 (4787–11503)	6.41 (5.24–7.52)	31 084 (22 025–39 633)
DANISH patients younger than 68 y	CMT + ICD	39 263 (35 145–43 447)	8.93 (6.90–11.05)	
	CMT	9529 (5719–14 013)	7.65 (5.97–9.34)	23 230 (17 208–61 785)

90%CrI, 90% credibility interval; CMT, conventional medical treatment; DANISH, DANish trial to assess efficacy of Implantable cardioverter defibrillators in patients with nonischemic Systolic Heart failure; ICD, implantable cardioverter defibrillator; ICER, incremental cost-effectiveness ratio; QALYs, quality-adjusted life years.

clinical trial and, therefore, the most representative of current clinical practice. The monthly probabilities of mechanical complications during follow-up (rupture, dislodgment, or displacement) were obtained from the articles by Ezzat et al.³³ and Koneru et al.³⁴

Utilities

All utilities were calculated using the same assumptions defined in the 2011 study.²⁰ Utility is an index measuring preference-based quality of life from 0 to 1, where 0 represents death and 1 represents perfect health. It was assumed that 1 year of life with ventricular dysfunction was equal to 0.75 years with perfect health, based on previous studies.^{35,36} It was also assumed that, in the absence of complications, ICD usage did not affect quality of life. In health states involving patient worsening, quality of life was assumed to deteriorate by 0.35 points per year of life during the days the patient remained hospitalized.³⁷

Resource use and costs

Resource use and costs were analyzed using direct health costs. In particular, the use of resources not directly ICD-related was presumed to be similar and not excessively high in either group and, therefore, these costs were not included. The regular drugs used in patients with heart failure with or without ischemic heart disease were included, assuming a percentage of use obtained from experts (clinical cardiologists and arrhythmia specialists) in the Cardiology Department of Vall d'Hebron Hospital, as well as the target, maximum, or most common doses and the costs according to the Pharmaceutical Provision System of the Ministry of Health, Consumer Affairs, and Social Welfare (table 2 of the supplementary data).

Table 3 of the supplementary data contains a description of the events defining each state, the assumptions made for each, the unit costs, and the sources of cost information. Resource use in the CMT + ICD group was based on routine clinical practice at Vall d'Hebron Hospital. The hospitalization costs for a first implant or complication-related replacement were estimated from a 22-patient sample provided by the analytical accounting at the same hospital. The unit costs include 20% overhead costs (eg, maintenance, services).

All costs are expressed as euros in 2018. When costs were not available for 2018, they were inflation-adjusted using the variation described by the National Institute of Statistics.

Sensitivity analysis

To evaluate the degree of uncertainty of the estimates, probabilistic sensitivity analyses were performed by assigning

various distribution functions to the parameters: beta distribution for probabilities and utilities, and gamma-uniform distributions for resource use. The cost-effectiveness planes were obtained by Monte Carlo simulations of 10 000 iterations. The willingness-to-pay threshold for the base case was assumed to be the upper limit of the range of €20 000 to €25 000/QALY, empirically determined by Vallejo-Torres et al.³⁸ Acceptability curves were also plotted to represent the probabilities of accepting ICD as a cost-effective alternative for different willingness-to-pay levels.^{39–41} To evaluate the impact on the result from the model estimations for each parameter, sensitivity analyses were performed using the variability found in the literature or variations of 20%. For intervention costs, the values used in the sensitivity analysis considered the variability between Vall d'Hebron Hospital and data provided by the Canary Islands Health Service. The base case assumes the most affordable device of those used at Vall d'Hebron Hospital and, therefore, this parameter was modified only with higher values in the sensitivity analysis. The results of this analysis were plotted in tornado charts.

RESULTS

Table 3 lists the results of the base case for the probabilistic analysis of the 3 scenarios analyzed. In the 3 cases, ICD implantation represents a QALY gain and a higher cost. The mean incremental cost-effectiveness ratios (ICERs) for CMT + ICD vs CMT were estimated at €19 171, €31 084, and €23 230/QALY for the base case in patients with ischemic heart disease, patients with nonischemic heart disease, and DANISH patients (with nonischemic heart disease) younger than 68 years, respectively. The points on the cost-effectiveness planes (figure 2) show the ICER result for each Monte Carlo simulation. Most points are in the upper right quadrant, indicating a higher cost and effectiveness for the CMT + ICD group compared with CMT in the 3 scenarios. In the nonischemic patient cohort, the percentage of cases to the left of the willingness-to-pay threshold of €25 000/QALY is higher. The probability that the CMT + ICD alternative would be cost-effective above a willingness-to-pay threshold of €25 000/QALY was 80% in patients with ischemic heart disease, 25% in nonischemic patients, and 55% in DANISH patients younger than 68 years (figures 4 and 5 of the supplementary data).

The sensitivity analyses (figure 3) show that the 3 models were sensitive, apart from the annual discount rate, device cost, utility value of the stable state (higher utility value in patients with stable heart failure signifies larger QALY gain if death is avoided by ICD use), and battery replacement rate (more replacements over a patient's lifetime will yield a higher cost per patient). In patients with ischemic heart disease, considering only device costs at the upper end of the range analyzed would lead to a different interpretation of the results compared with a threshold of €25 000/QALY. Two other important factors, but with a less




Figure 2. Incremental cost-effectiveness planes for ICD plus CMT vs CMT in patients with ischemic heart disease (A), patients with nonischemic heart disease (B), and DANISH patients younger than 68 years (C). The dashed line shows the threshold of €25 000/QALY. CMT, conventional medical treatment; ICD, implantable cardioverter-defibrillator; QALYs, quality-adjusted life years.




Figure 3. Results of univariate sensitivity analysis (tornado chart) in patients with ischemic heart disease (A), patients with nonischemic heart disease (B), and DANISH patients younger than 65 years (C). The middle value of the chart is the result for the base case, and the impact of an upward or downward parameter variation are shown to the right and left. The willingness-to-pay reference threshold is assumed to be €25 000/QALY. HF, heart failure; ICD, implantable cardioverter defibrillator; ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life year; WTP, willingness to pay in €/QALY.

relevant individual impact, would be to vary the cost of the hospitalization state (by reducing the number and/or length of hospitalizations) or to reduce the possibility of serious complications requiring repeat surgery and long hospital stays to negligible levels. In the case of patients with nonischemic heart disease, no single parameter change showed an interpretation economically favorable to the use of the ICD below a threshold of €25 000/QALY.

DISCUSSION

The present economic assessment from the perspective of the national health care system yielded similar results to those obtained in the last assessment in our setting carried out in 2011,¹⁹ but with differences explained by new data related to efficacy, resource use, and costs. An ICD implant for primary prevention in patients with ischemic heart disease yielded a slight gain in QALY (1.15 for a time horizon of 40 years) at a higher cost (€22 170), with an ICER of €19 278/QALY. In patients with ventricular dysfunction not due to ischemic heart disease, ICD implant represents a lower QALY gain (0.82) at a higher cost (€26 245), with an ICER of €32 006 /QALY, higher than the threshold of €25 000 /QALY. However, according to the DANISH trial data, the result for patients with nonischemic ventricular dysfunction would be more favorable for ICD in younger patients (younger than 68 years), in whom the benefit was somewhat higher (QALY difference, 1.22) and the ICER was near the threshold of €25 000 (€24 142 /QALY).

A willingness-to-pay threshold of €25 000/QALY³⁸ was selected based on the only empirical study in our setting and accepted as a reference level by the Spanish Network of Agencies for Health Technology Assessment. However, the cost-effectiveness of ICD in patients with nonischemic ventricular dysfunction would be around the threshold of €30 000/QALY most commonly accepted until recently in Spain.

The differences compared with the 2011 evaluation were almost entirely due to differences in the incremental cost. The QALY gain was also small and very similar to that obtained in this study (0.95 in the ischemic subgroup; 0.79 in the nonischemic subgroup) whereas the estimated cost increase was higher, €36 376 in ischemic patients (ICER, €38 371) and €52 694 in nonischemic patients (ICER, €52 694). This cost difference is mainly attributable to the decrease in device cost and the longer mean life of the generators. This cost difference between ischemic and nonischemic patients is mainly attributable to differences in the likelihood of death due to higher survival of nonischemic patients and, therefore, a higher cumulative cost during follow-up.

In view of the ICER distribution in the patient subgroup with nonischemic ventricular dysfunction according to age, a high percentage of patients were unlikely to benefit from ICD, and decisions for these patients should be made understanding that the opportunity cost of implanting an ICD is high. Nevertheless, there is still some uncertainty regarding this patient profile, and efforts should focus on improving and encouraging individualized decisions. Several other prediction models based on other factors along with EF have been shown to have predictive value regarding the risk of sudden cardiac death, such as left ventricular end-diastolic diameter, ventricular tachycardia induction in the electrophysiologic study, the presence of spontaneous nonsustained ventricular tachycardia, left bundle-branch block, fragmented QRS,⁴² and particularly the presence, location, and extent of myocardial fibrosis.⁴³ The health benefit obtained for each euro invested would be enhanced by encouraging the use of these or other models in differential risk prediction for sudden and

nonsudden death to determine the patient profile most benefitting from prophylactic ICD implantation.

Likewise, it is important to consider patient preferences under various circumstances. For instance, this analysis did not include end-of-life replacements or ICD generator depletion with no discharges to the patient. In both cases, the need for replacement should be reassessed according to the patient's preferences and updated life expectancy.

Limitations

Cost-utility estimation is based on a decision-making model, and hence its validity depends on the validity of the model's assumptions and parameters. Ultimately, the validity of the model depends on the sources of scientific evidence used to estimate the model parameters. The main aim of the present study was to include new evidence on efficacy and complication costs and rates that has become available since 2011. The clinical efficacy evidence was updated by using a systematic review and meta-analysis, incorporating the DANISH trial in the nonischemic ventricular dysfunction subgroup. The effectiveness evidence in patients with ischemic heart disease was the same as that used in 2011 and, therefore, may reflect a situation not entirely consistent with current clinical practice.

Likewise, resource use and costs are based on expert opinion, the health care process, and the analytical accounting of a single site, although we believe that the data are a more accurate picture of reality than those obtained from rates or from the literature. The external validity of the device cost estimate is more limited because the final fixed cost often depends on negotiations between the supply center and industry. Our approach chose the most affordable option (single-chamber generator from a specific supplier), but sensitivity analyses were carried out for other assumptions.

Although some patients received a dual-chamber device (14% according to the Spanish Cardiology Society registry), it was assumed that this was indicated less often for primary prevention.

Another important limitation is that ICD was not compared with ICD plus cardiac resynchronization therapy in patients with an indication to switch to this therapy due to heart failure progression. There is some debate about whether the expected benefits of ICD are greater in patients with no indication for CRT than in patients with an indication⁴⁴ and, therefore, economic models should be developed with a structure that includes this possibility. However, this would require more sound data on efficacy and effectiveness, adverse events, and quality of life in these patients. Equally, we did not take into account the introduction of new alternative drug treatments, such as sacubitril-valsartan, despite proven efficacy and potentially higher cost-effectiveness than those of ICDs,^{45,46} possibly reducing the number of patients who may be eligible for an ICD. Follow-up also did not include long-term consequences that may have a different weight among patients with or without an ICD, such as the possibility of receiving a transplant or ventricular assist device.

CONCLUSIONS

The efficiency of the single-chamber ICD has improved in the past decade and is cost-effective in patients with ischemic and nonischemic left ventricular dysfunction younger than 68 years, assuming a willingness-to-pay threshold of €25 000/QALY. In older nonischemic patients, the ICER is around €30 000 /QALY.

WHAT IS KNOWN ABOUT THE TOPIC?

- Compared with conventional medical treatment, ICD in primary prevention improves the survival of patients with ischemic and nonischemic heart failure with EF \leq 35%.
- Prophylactic ICD use accounts for more than half of all indications, and is the fastest-growing indication in patients with dilated cardiomyopathy.
- A 2011 assessment found that ICD was cost-effective for primary prevention in patients with ischemic dilated cardiomyopathy who met the criteria for the MADIT trial, but not in patients who met the MADIT II criteria or had nonischemic dilated cardiomyopathy.

WHAT DOES THIS STUDY ADD?

- Assuming a willingness-to-pay threshold of €25 000/QALY, ICD was is cost-effective for patients with ischemic left ventricular dysfunction.
- An ICD implant in patients with ventricular dysfunction not due to ischemic heart disease yields a lower QALY gain at a higher cost and would only be cost-effective below the threshold of €25 000 /QALY in younger patients (younger than 68 years), who obtain greater benefit.

FUNDING

This study was funded by the Ministry of Health, Consumer Affairs, and Social Welfare in the development framework of activities in the Annual Work Plan of Spanish Network of Health Technology Assessment and National Health Care Services (2018).

AUTHORS' CONTRIBUTIONS

A. Ribera, E. Giménez, and G. Oristrell prepared the first draft of the article. E. Giménez and J.R. Marsal performed the economic modeling. A. Ribera, J.R. Marsal, D. Osorio, and M. Ballesteros performed the meta-analysis. A. Ribera, J.R. Marsal, D. Osorio, M. Ballesteros, E. Ródenas, Y. Belahnech, and R. Escalona reviewed the abstracts and full articles to collect information for the systematic review. G. Oristrell, N. Rivas, I. Roca-Luque, and I. Ferreira-González participated as a clinical expert committee in determining the parameters of the economic model. L. García-Pérez developed the initial economic model and participated in its current adaptation. A. Ribera, I. Ferreira-González, and M. Espallargues sponsored the study and assisted with obtaining funding for its conduct. All authors contributed substantially to the interpretation of the results obtained in the systematic review and meta-analysis phase and in the financial modeling. All authors performed a critical review of the various draft versions and provided considerable intellectual input.

CONFLICTS OF INTEREST

N. Rivas received fees from Abbott and Boston-Scientific unrelated to the article submitted. I. Roca-Luque received fees from Abbott, Medtronic, and Boston unrelated to the article

submitted. E. Ródenas received an unconditional grant from Biotronik unrelated to the article submitted.

APPENDIX. SUPPLEMENTARY DATA

Supplementary data associated with this article can be found in the online version available at <https://doi.org/10.1016/j.rec.2021.05.004>

REFERENCES

1. Priori SG, Blomström-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur Heart J*. 2015;36:2793–2867.
2. Moss AJ, Zareba W, Hall WJ, et al. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. 2002;346:877–883.
3. Bardy GH, Lee KL, Mark DB, et al. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med*. 2005;325:225–237.
4. Køber L, Thune JJ, Nielsen JC, et al. Defibrillator implantation in patients with nonischemic systolic heart failure. *N Engl J Med*. 2016;375:1221–1230.
5. Friedman DJ, Al-Khatib SM, Zeitler EP, et al. New York Heart Association class and the survival benefit from primary prevention implantable cardioverter defibrillators: A pooled analysis of 4 randomized controlled trials. *Am Heart J*. 2017;191:21–29.
6. Shun-Shin MJ, Zheng SL, Cole GD, Howard JP, Whinnett ZI, Francis DP. Implantable cardioverter defibrillators for primary prevention of death in left ventricular dysfunction with and without ischaemic heart disease: a meta-analysis of 8567 patients in the 11 trials. *Eur Heart J*. 2017;38:1738–1746.
7. Alba AC, Foroutan F, Duero Posada J, et al. Implantable cardiac defibrillator and mortality in non-ischaemic cardiomyopathy: an updated meta-analysis. *Heart*. 2018;104:230–236.
8. Beggs SAS, Jhund PS, Jackson CE, McMurray JJV, Gardner RS. Non-ischaemic cardiomyopathy, sudden death and implantable defibrillators: a review and meta-analysis. *Heart*. 2018;104:144–150.
9. Siddiqui WJ, Aggarwal S, Rafique M, Singh S, Kutalek S, Eisen HJ. Prophylactic use of the implantable cardioverter-defibrillator and its effect on the long-term survival, cardiovascular and sudden cardiac death in nonischemic cardiomyopathy patients—a systematic review and meta-analysis. *Heart Fail Rev*. 2018;23:181–190.
10. Barakat AF, Saad M, Elgendy AY, et al. Primary prevention implantable cardioverter defibrillator in patients with non-ischaemic cardiomyopathy: a meta-analysis of randomised controlled trials. *BMJ Open*. 2017;7:e016352.
11. Cavalcanti R, Aboul-Hosn N, Morales G, Abdel-Latif A. Implantable cardioverter defibrillator for the primary prevention of sudden cardiac death in patients with nonischemic cardiomyopathy. *Angiology*. 2018;69:297–302.
12. Al-Khatib SM, Fonarow GC, Joglar JA, et al. Primary prevention implantable cardioverter defibrillators in patients with nonischemic cardiomyopathy: A meta-analysis. *JAMA Cardiol*. 2017;2:685–688.
13. Xing Z, Tang L, Chen C, Huang J, Zhu Z, Hu X. Effectiveness of implantation of cardioverter-defibrillators therapy in patients with non-ischemic heart failure: an updated systematic review and meta-analysis. *Brazilian J Cardiovasc Surg*. 2017;32:417–422.
14. Kunz WG, Hunink MGM, Sommer WH, et al. Cost-effectiveness of endovascular stroke therapy. *Stroke*. 2016;47:2797–2804.
15. Wolff G, Lin Y, Karathanos A, et al. Implantable cardioverter/defibrillators for primary prevention in dilated cardiomyopathy post-DANISH: an updated meta-analysis and systematic review of randomized controlled trials. *Clin Res Cardiol*. 2017;106:501–513.
16. Stavrakis S, Asad Z, Reynolds D. Implantable cardioverter defibrillators for primary prevention of mortality in patients with nonischemic cardiomyopathy: a meta-analysis of randomized controlled trials. *J Cardiovasc Electrophysiol*. 2017;28:659–665.
17. Anantha Narayanan M, Vakil K, Reddy YN, et al. Efficacy of implantable cardioverter-defibrillator therapy in patients with nonischemic cardiomyopathy: a systematic review and meta-analysis of randomized controlled trials. *JACC Clin Electrophysiol*. 2017;3:962–970.
18. Fernández Lozano I, Osca Asensi J, Alzueta Rodríguez J. Spanish Implantable Cardioverter-defibrillator Registry 16th Official Report of the Heart Rhythm Association of the Spanish Society of Cardiology (2019). *Rev Esp Cardiol*. 2020;73:1026–1037.
19. Campillo-Artero C, Ortún V. El análisis de coste-efectividad: por qué y cómo. *Rev Esp Cardiol*. 2016;69:370–373.
20. García Pérez L, Pinilla Domínguez P, García García FJ, García Quintana A, Caballero Dorta E, Linertová R. Coste-efectividad del desfibrilador automático implantable en prevención primaria en España. Ministerio de Sanidad, Servicios Sociales e Igualdad. Servicio de Evaluación del Servicio Canario de la Salud; 2011. Informes de Evaluación de Tecnologías Sanitarias. Available at: <https://www3.gobiernodecanarias.org/sanidad/scs/contenidoGenerico.jsp?idDocumento=5cc0e975-ae0a-11e4-9053-9d1690bb437a&idCarpeta=993a9b1d-7aed-11e4-a62a-758e414b4260>. Accessed 6 Feb 2021.

21. Moss AJ, Hall WJ, Cannom DS, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. *N Engl J Med*. 1996;335:1933–1940.
22. Kramer DB, Friedman PA, Kallinen LM, et al. Development and validation of a risk score to predict early mortality in recipients of implantable cardioverter-defibrillators. *Hear Rhythm*. 2012;9:42–46.
23. Bilchick KC, Stukenborg GJ, Kamath S, Cheng A. Prediction of mortality in clinical practice for medicare patients undergoing defibrillator implantation for primary prevention of sudden cardiac death. *J Am Coll Cardiol*. 2012;60:1647–1655.
24. Koller MT, Schaefer B, Wolbers M, Sticherling C, Bucher HC, Osswald S. Death without prior appropriate implantable cardioverter-defibrillator therapy: a competing risk study. *Circulation*. 2008;117:1918–1926.
25. Elming MB, Nielsen JC, Haarbo J, et al. Age and outcomes of primary prevention implantable cardioverter-defibrillators in patients with nonischemic systolic heart failure. *Circulation*. 2017;136:1772–1780.
26. Bigger JT. Prophylactic use of implanted cardiac defibrillators in patients at high risk for ventricular arrhythmias after coronary-artery bypass graft surgery. *N Engl J Med*. 1997;337:1569–1575.
27. Hohnloser SH, Kuck KH, Dorian P, et al. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med*. 2004;351:2481–2488.
28. Steinbeck G, Andresen D, Seidl K, et al. Defibrillator implantation early after myocardial infarction. *N Engl J Med*. 2009;361:1427–1436.
29. Bönsch D, Antz M, Boczor S, et al. Primary prevention of sudden cardiac death in idiopathic dilated cardiomyopathy: The Cardiomyopathy Trial (CAT). *Circulation*. 2002;105:1453–1458.
30. Kadish A, Dyer A, Daubert JP, et al. Prophylactic defibrillator implantation in patients with nonischemic dilated cardiomyopathy. *N Engl J Med*. 2004;350:2151–2158.
31. Strickberger SA, Hummel JD, Bartlett TG, et al. Amiodarone versus implantable cardioverter-defibrillator: Randomized trial in patients with nonischemic dilated cardiomyopathy and asymptomatic nonsustained ventricular tachycardia - AMIO-VIRT. *J Am Coll Cardiol*. 2003;41:1707–1712.
32. Pezawas T, Grimm M, Ristl R, et al. Primary preventive cardioverter-defibrillator implantation (Pro-ICD) in patients awaiting heart transplantation. A prospective, randomized, controlled 12-year follow-up study. *Transpl Int*. 2015;28:34–41.
33. Ezzat VA, Lee V, Ahsan S, et al. A systematic review of ICD complications in randomised controlled trials versus registries: is our 'real-world' data an underestimation? *Open Heart*. 2015;2:e000198.
34. Koneru JN, Jones PW, Hammill EF, Wold N, Ellenbogen KA. Risk factors and temporal trends of complications associated with transvenous implantable cardiac defibrillator leads. *J Am Heart Assoc*. 2018;7:e007691.
35. Gandjour A, Dipl-Ges-Ök AH, Adarkwah CC, Adarkwah CC. Cost-effectiveness of implantable defibrillators after myocardial infarction based on 8-year follow-up data (MADIT II). *Value Heal*. 2011;14:812–817.
36. Martí B, Delgado J, Oliva J, et al. PCV120 quality of life in chronic symptomatic heart failure patients in Spain, insight from the Inoescaro Study. *Value Heal*. 2011;14:A386.
37. García-Pérez L, Pinilla Domínguez P, Davila Ramos MF, Copca Alvarez A, Ruiz Hernández JJ, Díaz Escofet M. Utilidades a partir del EQ-5D en pacientes hospitalizados con insuficiencia cardíaca en Canarias. *Gac Sanit*. 2012;26(Espec Congr 1):49.
38. Vallejo-Torres L, García-Lorenzo B, Castilla I, et al. Valor monetario de un año de vida ajustado por calidad: estimación empírica del coste de oportunidad en el Sistema Nacional de Salud. 2015. Available at: <http://www3.gobiernodecanarias.org/sanidad/scs/contenidoGenerico.jsp?idDocumento=e690e0c1-cbed-11e5-a9c5-a398589805dc&idCarpeta=993a9b1d-7aed-11e4-a62a-758e414b4260>. Accessed 16 Feb 2021.
39. Sacristán JA, Oliva J, Llano J, Del Prieto L, Pinto JL. ¿Qué es una tecnología sanitaria eficiente en España? *Gac Sanit*. 2002;16:334–343. Available at: http://scielo.isciii.es/scielo.php?pid=S0213-91112002000400008&script=sci_abstract&lng=pt. Accessed 5 Apr 2019.
40. De Cock E, Miravittles M, González-Juanatey JR, Azanza-Perea JR. Valor umbral del coste por año de vida ganado para recomendar la adopción de tecnologías sanitarias en España: evidencias procedentes de una revisión de la literatura. *Pharmacoeconomics Spanish Res Artic*. 2007;4:97–107.
41. Briggs A, Fenn P. Confidence intervals or surfaces? Uncertainty on the cost-effectiveness plane. *Health Econ*. 1998;7:723–740.
42. Goldberger JJ, Subačius H, Patel T, Cunnane R, Kadish AH. Sudden cardiac death risk stratification in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol*. 2014;63:1879.
43. Kuruvilla S, Adenaw N, Katwal AB, Lipinski MJ, Kramer CM, Salerno M. Late gadolinium enhancement on cardiac magnetic resonance predicts adverse cardiovascular outcomes in nonischemic cardiomyopathy: A systematic review and meta-analysis. *Circ Cardiovasc Imaging*. 2014;7:250–257.
44. Golwala H, Bajaj NS, Arora G, Arora P. Implantable Cardioverter-defibrillator for nonischemic cardiomyopathy: an updated meta-analysis. *Circulation*. 2017;135:201–203.
45. McMurray JJV, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993–1004.
46. Zacà V. Sacubitril/valsartan or an implantable cardioverter-defibrillator in heart failure with reduced ejection fraction patients: a cost-effectiveness analysis. *J Cardiovasc Med (Hagerstown)*. 2018;19:597–605.
47. Comín-Colet J, Anguita M, Formiga F, et al. Calidad de vida relacionada con la salud de los pacientes con insuficiencia cardíaca crónica sistólica en España: resultados del estudio VIDA-IC. *Rev Esp Cardiol*. 2016;69:256–271.