

Letters to the Editor

Looking for ECG signs of acute coronary syndrome



En búsqueda de signos ECG para el síndrome coronario agudo

To the Editor,

We read with interest the article published by Fernández-Sánchez et al.¹ in which they described the South African flag sign. This is one of many signs already described to help identify particular electrocardiogram patterns associated with acute coronary syndrome, such as the shark fin sign, the triangular wave, and the lambda-like wave. Nonetheless, in our opinion, the South African flag sign is an ill-suited teaching tool,² as one must first be familiar with the flag. The sign also demands a certain degree of imagination and only applies to 3 × 4 display formats. In addition, the ST depression mirror image is also observed in II and aVF, meaning it is positioned outside the flag. This is, however, not the main point we wish to make in reference to the article by Fernández-Sánchez et al.

The pattern described by the authors is a characteristic finding in first diagonal or intermediate branch occlusion. Although this pattern has traditionally been considered to depict high lateral ischemia due to circumflex artery occlusion, when not associated with ST depression in V₁-V₂,³ it is a manifestation of occlusion of the first diagonal or intermediate branch, which both supply blood to the mid-low anterolateral wall, not the high lateral wall.⁴ Strictly speaking, thus, the pattern represents mid-anterior myocardial infarction, which in its subacute or chronic phase can be confirmed by delayed enhancement cardiac magnetic resonance imaging.⁵

We agree with the authors that it is a grave error to expect ST elevation in 2 contiguous leads when assessing a potential case of acute coronary syndrome. In this particular case, however, the I and aVL leads displaying ST elevation in the frontal plane can be considered contiguous.

FUNDING

Article funded by *Fundación Instituto de Investigación Sanitaria de les Illes Balears* (IdISBa).

STATEMENT ON THE USE OF ARTIFICIAL INTELLIGENCE

Artificial intelligence has not been used.

Looking for ECG signs of acute coronary syndrome. Response



En búsqueda de signos ECG para el síndrome coronario agudo. Respuesta

To the Editor,

We thank Fiol et al. for their comments on our article. After reviewing the literature cited, we agree that the term *high lateral infarction* (from the classic Myers classification) is incorrect and

AUTHORS' CONTRIBUTIONS

All the authors contributed to revising this letter to the editor.

CONFLICTS OF INTEREST

No conflicts of interest.

Miquel Fiol Sala,^{a,*} Andrés Carrillo López,^b and Alberto Rodríguez Salgado^c

^a*Instituto de Investigación Sanitaria de les Illes Balears (IdISBa), Palma, Balearic Islands, Spain*

^b*Servicio de Medicina Intensiva, Hospital Universitario Miguel Servet, Zaragoza, Spain*

^c*Servicio de Medicina Intensiva, Hospital Universitario Son Espases, Instituto de Investigación Sanitaria de les Illes Balears (HUSE-IdISBa), Palma, Balearic Islands, Spain*

* Corresponding author:

E-mail address: miguel.fiol@ssib.es (M. Fiol Sala).

✉ @miquelfiol (M. Fiol Sala).

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2022.12.014>

<https://doi.org/10.1016/j.rec.2023.10.008>

REFERENCES

1. Fernández-Sánchez JA, Arriaga-Jiménez AE, Jiménez-Fernández M. IAMCEST lateral alto con signo de la bandera de Sudáfrica. *Rev Esp Cardiol.* 2023;76:665.
2. Littmann sl. South African flag sign: a teaching tool for easier ECG recognition of high lateral infarct. *Am J Emerg Med.* 2016;34:107–109.
3. Birnbaum Y, Hasdai D, Sclarovsky S, Herz I, Strasberg B, Rechavia E. Acute myocardial infarction entailing ST segment elevation in lead AVL: electrocardiographic differentiation among occlusion of the left anterior descending, first diagonal and first obtuse marginal coronary arteries. *Am Heart J.* 1996;131:38–42.
4. Bayés de Luna A, Wagner G, Birnbaum Y, et al. A new terminology for the left ventricular walls and for the location of myocardial infarcts that present Q wave based on the standard of cardiac magnetic resonance imaging: a statement for healthcare professionals from a committee appointed by the International Society for Holter and Non Invasive Electrocardiography. *Circulation.* 2006;114:1755–1760.
5. Bayes de Luna A, Cino JM, Pujadas S, et al. Concordance of electrocardiographic patterns and healed myocardial infarction location detected by cardiovascular magnetic resonance. *Am J Cardiol.* 2006;97:443–451.

<https://doi.org/10.1016/j.rec.2023.09.013>

1885-5857/© 2023 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Cardiología.

should no longer be used. Given the variable blood supply to the mid-lateral segment (and more specifically to echocardiographic segments 12 and 16, which can be supplied by the circumflex artery or the first diagonal artery), we consider that *limited anterolateral wall myocardial infarction*¹ is a more appropriate term.

While it is certainly true that the I and aVL leads are contiguous in the frontal plane, they appear separately in ECG tracings (whether 12-lead or 3 × 4 display formats), leaving them open to misinterpretation by an inexperienced reader.