

Use of Antithrombotic Therapy in Patients With Atrial Fibrillation in Primary Care. Importance of INR Control



Uso del tratamiento antitrombótico en pacientes con fibrilación auricular en atención primaria. Importancia del control del INR

To the Editor,

It was with great interest that we read the letter by Barrios et al¹ about the use of antithrombotic therapy by applying the CHA₂DS₂-VASc scale in patients with atrial fibrillation in primary care. Their objective was to determine whether there were any differences in the use of antithrombotic therapy, depending on whether the CHADS₂ or CHA₂DS₂-VASc scale was applied. To do this, they analyzed data from the VAL-FAAP study,² a registry constructed in primary care between September 2009 and May 2010. The authors affirm that antithrombotic therapy use is far from optimal, since they found that oral anticoagulants were used in 44.3% of patients with CHA₂DS₂-VASc 0 but not in 33.5% of patients with CHA₂DS₂-VASc ≥2 (ie, 66.5% received oral anti-coagulation). They conclude that in Spain treatment to prevent thromboembolic complications is administered inadequately and is apparently prescribed independently of the risk stratification scale being used.

On the basis of the findings in the aforementioned letter,¹ we would ask the following questions: Can we consider that antithrombotic therapy use is "far from optimal" or "inadequate"? Can we determine that its use is "far from optimal" without taking account of International Normalized Ratio (INR) control? Currently, what is the role of the new anticoagulants that require no INR monitoring? In our opinion, the CHA₂DS₂-VASc scale should not have been used, nor should the European Society of Cardiology 2010 recommendations for oral anticoagulant use, since they were subsequent to the construction of the VAL-FAAP registry. Perhaps the results should be interpreted from a more positive perspective in the light of recent publications^{3–5} showing clear improvements in the use of this treatment. Kirchhof et al³ analyzed the management of atrial fibrillation in seven European countries, including Spain, between January 2012 and January 2013 (post-VAL-FAAP) and, in fact, found that 85.6% of patients with CHA₂DS₂-VASc ≥2 received oral anticoagulants, a far higher percentage than that reported in the VAL-FAAP study. Furthermore, they also found that 62.5% of patients with CHA₂DS₂-VASc 0 received oral anticoagulants and stressed that only 35% of patients treated with acenocoumarol had INR values within the therapeutic range in their last 3 determinations, which constitutes a high added risk of bleeding or thrombotic complications. Recently, Lip et al⁴ have shown that 70.9% of patients with CHA₂DS₂-VASc ≥2 receive oral anticoagulants. They also reported that new anticoagulation agents were administered to 6.9% of all patients receiving anticoagulation therapy. Focussing on data from primary care in Spain, Lobos-Bejarano et al⁵ found that 84.1% of all patients with atrial fibrillation received anticoagulation therapy. Of these,

68.8% had CHADS₂ ≥2 vs 31.2% with CHADS₂ ≤1. Of patients receiving oral anticoagulants, 66% had their last INR value within the therapeutic range.

These findings show a positive tendency in antithrombotic therapy use, although there remains room for improvement. They also help us reflect on what is most important in antithrombotic therapy. Clearly, correct indication is fundamental but we should never forget the importance of adequate INR control in these patients, since severe bleeding complications can be fatal. We are convinced that primary care can play a fundamental role in identifying patients with inadequate control of anticoagulation with vitamin K antagonists, in which case, we should consider using the new anticoagulants. Currently, these drugs are clearly underused, despite being much safer in terms of bleeding complications because they require no dose adjustment as a function of the patient's INR value and represent an important therapeutic advance.⁴ We should remember that our final objective is to prevent complications in patients with atrial fibrillation, which is impossible with inadequate control.

Mariano Leal-Hernández^{a,*} and Francisco Marín^b

^aCentro de Salud de San Andrés, Murcia, Spain

^bServicio de Cardiología, Hospital Clínico Universitario Virgen de la Arrixaca, Instituto Murciano de Investigación en Biomedicina (IMIB), El Palmar, Murcia, Spain

* Corresponding author.

E-mail address: arboleja@yahoo.es (M. Leal-Hernández).

Available online 3 April 2014

REFERENCES

- Barrios V, Escobar C, Calderón A, Rodríguez G, Llisterri JL, Polo J. Uso del tratamiento antitrombótico según la escala CHAD₂DS₂-VASc en los pacientes con fibrilación auricular en atención primaria. Rev Esp Cardiol. 2014;67:150–1.
- Barrios V, Calderón A, Escobar C, De la Figuera M. Pacientes con fibrilación auricular asistidos en consultas de atención primaria Estudio Val-FAAP. Rev Esp Cardiol. 2012;65:47–53.
- Kirchhof P, Ammentorp B, Darius H, De Caterina R, Le Heuzey JY, Schilling RJ, et al. Management of atrial fibrillation in seven European countries after the publication of the 2010 ESC Guidelines on atrial fibrillation: primary results of the PREvention of thromboembolic events—European Registry in Atrial Fibrillation (PREFFER in AF). Europace. 2014;16:6–14.
- Lip GY, Laroche C, Dan GA, Santini M, Kalarus Z, Rasmussen LH, et al. 'Real-world' antithrombotic treatment in atrial fibrillation: the EURObservational Research Programme Atrial Fibrillation General Pilot survey. Am J Med. 2014. Jan 28. doi: 10.1016/j.amjmed.2013.12.022.
- Lobos-Bejarano JM, Del Castillo-Rodríguez JC, Mena-González A, Alemán-Sánchez JJ, Cabrera de León A, Barón-Esquivias G, et al. Características de los pacientes y abordaje terapéutico de la fibrilación auricular en atención primaria en España Estudio FIATE. Med Clin (Barc). 2013;141:279–86.

SEE RELATED ARTICLES:

<http://dx.doi.org/10.1016/j.rec.2013.07.009>

<http://dx.doi.org/10.1016/j.rec.2014.03.001>

<http://dx.doi.org/10.1016/j.rec.2014.02.003>