

## ECG Contest

## Response to ECG, October 2019

## Respuesta al ECG de octubre de 2019


Pasquale Crea,<sup>a,\*</sup> Angela Nicotera,<sup>b</sup> and Teresa Crea<sup>c</sup><sup>a</sup> Cardiology Unit, Department of Clinical and Experimental Medicine, University of Messina, Italy<sup>b</sup> Cardiology Unit, Papardo Hospital, Messina, Italy<sup>c</sup> Internal Medicine, Department of Clinical and Experimental Medicine, University of Messina, Italy

Figure 1.

The correct answer is number 3. As shown in Figure 1, the P-P cycle exactly doubles. This finding strongly suggests a 2:1 sinoatrial (S-A) block. Sympathovagal imbalance and respiratory sinus arrhythmia would have generated a gradually lengthening and shortening of the P-P cycle in a cyclical fashion, usually corresponding to the phases of the respiratory cycle or other autonomic reflexes. Thus, answers 1 and 2 are incorrect. A concealed atrioventricular (A-V) node reentry cannot explain sinus node depression. Moreover, the ECG would have shown a trigger, preceding the pause, such as a Wenckebach phenomenon or a premature atrial impulse.<sup>1</sup> For this reason, answer 4 is incorrect.

The sinus impulse is only recorded by the ECG once it has left the sinus node and activated the atrium, thereby resulting in the P wave.<sup>2</sup> Delay in conduction at the S-A junction, such as that occurring during a first-degree S-A block is, therefore, not visible electrocardiographically. Second-degree S-A blocks can be revealed by a careful analysis of the ECG. A Wenckebach conduction defect at the S-A level will present as progressive acceleration of the P-P intervals, analogously to the progressive acceleration of the R-R intervals in an A-V Wenckebach disturbance. Pause is less than twice the preceding P-P cycle.<sup>3</sup> In contrast, an abrupt doubling of the P-P cycle excludes a second-degree Mobitz type 2 or a 2:1 S-A block. It is worth noting that sinus bradycardia and 2:1 S-A block are indistinguishable on an ECG, unless an abrupt change in heart rate clarifies the underlying mechanism. Advanced S-A block will be revealed by a pause that is a multiple of > 2 of the P-P cycle. ECG of third-degree S-A block is analogous to sinus arrest.

## REFERENCES

1. Oreto G, Consolo A, Scimone IM. Manifest and concealed AV nodal reentry in Wenckebach type of AV conduction with AV junctional escape rhythm. *J Electrocardiol.* 1996;29:333–336.
2. de Almeida Henriques MA, Schamroth L. Sinus rhythm complicated by second-degree sino-atrial block. *Heart Lung.* 1976;5:137–138.
3. Schamroth L, Dove E. The Wenckebach phenomenon in sino-atrial block. *Br Heart J.* 1966;28:350–358.

## SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2018.11.006>

\* Corresponding author:

E-mail address: [pasqualecrea@hotmail.it](mailto:pasqualecrea@hotmail.it) (P. Crea).

<https://doi.org/10.1016/j.rec.2018.11.005>

1885-5857/© 2018 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.