

Cardiopulmonary exercise testing in patients with severe aortic stenosis: lights and shadows. Response



Prueba de esfuerzo con consumo de oxígeno en pacientes con estenosis aórtica grave: luces y sombras. Respuesta

To the Editor,

We appreciate the critical analysis regarding our study provided by Lacalzada-Almeida et al., in which they highlight their experience in the use of cardiopulmonary exercise testing (CPET) in patients with aortic stenosis (AS).

In their study, Lacalzada-Almeida et al.¹ found a higher percentage of abnormal CPET results—defined by symptoms and electrocardiographic and blood pressure changes—in patients with severe or paradoxical low-gradient AS than in patients with moderately severe AS. No differences were found in the peak oxygen consumption (pVO_2) or the VE/VCO_2 slope between these 3 groups. This may be explained by the limited population included and possible selection bias, as the authors point out, by excluding patients with inconclusive symptoms attributable to AS. In addition, a nonnegligible percentage of patients reached an $\text{RER} < 1.1$, which could lead to erroneous analyses if submaximal CPET parameters were not taken into account.¹

Currently, CPET is the only objective tool available to estimate myocardial oxygen consumption, ventilatory efficacy, and muscle performance on exertion, information of value to determine the cause of a patient's symptoms.² Hence, we propose CPET use in patients with asymptomatic severe AS, as many of them are falsely asymptomatic because of several concurrent factors, such as advanced age, frailty, obesity, and others.² pVO_2 values < 20 mL/kg/min and VE/VCO_2 slope values > 30 are pathological and imply a higher risk of adverse events on follow-up, as described by Guazzi et al.³ in the first published report proposing an algorithm for the study of valve disease with CPET.

Thus, our group proposes an objective algorithm to identify these falsely asymptomatic patients using CPET. In the absence of randomized studies, we believe it could be a useful instrument for this purpose.

FUNDING

No funding was received for the present publication.

AUTHORS' CONTRIBUTIONS

All authors have contributed to writing and critical review of the article.

CONFLICTS OF INTEREST

P. Avanzas is an associate editor of *Revista Española de Cardiología*; the editorial procedure established by the journal has been followed to guarantee impartial management of the manuscript. The other authors declare that they have no conflicts of interest.

Néstor Báez-Ferrer,^a Pablo Avanzas,^{b,c,d} and Alberto Domínguez-Rodríguez^{a,e,f,*}

^aDepartamento de Cardiología, Hospital Universitario de Canarias, Tenerife, Spain

^bÁrea del Corazón, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

^cInstituto de Investigación del Principado de Asturias, Oviedo, Asturias, Spain

^dDepartamento de Medicina, Facultad de Medicina, Universidad de Oviedo, Oviedo, Asturias, Spain

^eFacultad de Ciencias de la Salud, Universidad Europea de Canarias, Tenerife, Spain

^fCentro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Spain

*Corresponding author.

E-mail address: adrvd@hotmail.com (A. Domínguez-Rodríguez).

Available online 21 December 2022

REFERENCES

1. Lacalzada-Almeida J, Izquierdo-Gómez MM, Laynez-Cerdeña I, et al. Role of Exercise Testing and Speckle Tracking Echocardiography in Paradoxical Severe Aortic Stenosis. *Cureus*. 2021. <http://dx.doi.org/10.7759/cureus.18266>.
2. Báez-Ferrer N, Avanzas P, Domínguez-Rodríguez A. Papel de la prueba de esfuerzo con consumo de oxígeno en pacientes con estenosis aórtica grave. *Rev Esp Cardiol*. 2022. <http://dx.doi.org/10.1016/j.recresp.2022.07.005>.
3. Guazzi M, Arena R, Halle M, Piepoli MF, Myers J, Lavie CJ. 2016 focused update: Clinical recommendations for cardiopulmonary exercise testing data assessment in specific patient populations. *Eur Heart J*. 2018;39:1144–1161.

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2022.10.011>

<https://doi.org/10.1016/j.rec.2022.10.013>
1885-5857/

© 2022 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Variability and equity, a debatable relationship



Variabilidad y equidad, una relación discutible

To the Editor,

In a recent study published in *Revista Española de Cardiología*, de la Torre Hernández et al.¹ conclude that cardiovascular technology use varied considerably across Spain's autonomous communities and that the variability observed could not be explained by economic factors or number of hospital visits. The authors made several references to «equal opportunities» and even mentioned

«country-wide deficiencies in equitable access to cardiovascular interventions of proven clinical effectiveness». Based on the study's findings, we believe it may be somewhat rash to draw an inverse correlation between variability in technology use and equitable access to health care interventions. The authors acknowledge certain limitations of their study,¹ and these are neatly illustrated in an editorial on the subject.² Age is one of many factors not assessed in the study that can influence the use of cardiovascular procedures. An autonomous community with an older population, for example, may perform more procedures. Using a simple analysis of data from 2019, we observed a direct regional correlation between mean population age and the number of




Figure 1. Correlation adjusted for mean age of the autonomous communities. CRT, cardiac resynchronization therapy; GDP, gross domestic product per capita; HS, healthcare spending per capita; ICD, implantable cardioverter defibrillator; PCI, percutaneous coronary intervention; TAVI, transcatheter aortic valve implant.

cardiovascular procedures performed in Spain (figure 1). Variability in clinical practice is certainly an important issue and one that calls for in-depth analysis. Notwithstanding, clinical guidelines leave ample room for case-by-case decision-making. They do not provide recommendations on many clinical situations and should not be followed indiscriminately.

FUNDING

None

AUTHORS' CONTRIBUTIONS

Both authors contributed equally.

CONFLICTS OF INTEREST

None.

J. Alberto San Román^{a,b,*} and Itziar Gómez Salvador^{a,b}

^aServicio de Cardiología, Hospital Clínico Universitario, Valladolid, Spain

^bCentro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Spain

* Corresponding author:

E-mail address: asanroman@secardiologia.es (J.A. San Román).

Available online 21 December 2022

REFERENCES

- de la Torre Hernández JM, Lozano González M, García Camarero T, et al. Variabilidad interregional en el uso de tecnologías cardiovasculares (2011–2019). Correlación con índices económicos y frecuencia y mortalidad hospitalarias. *Rev Esp Cardiol.* 2022;75:806–816.
- Marrugat J, Subirana I, Dégano IR. ¿Hasta qué nivel regional se puede analizar la calidad asistencial? *Rev Esp Cardiol.* 2022;75:784–786.
- Ojeda S, Romaguera R, Cruz-González I, Moreno R. Registro Español de Hemodinámica y Cardiología Intervencionista. XXIX Informe Oficial de la Asociación de Cardiología Intervencionista de la Sociedad Española de Cardiología (1990–2019). *Rev Esp Cardiol.* 2020;73:927–936.

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2022.10.011>

<https://doi.org/10.1016/j.rec.2022.02.016>

<https://doi.org/10.1016/j.rec.2022.10.015>

1885–5857/

© 2022 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.