

## UPDATE

### Diabetes and Cardiovascular Diseases (VII)

# Coronary Angioplasty in Diabetic Patients. Current and Future Perspectives

Joaquín J. Alonso, Juan M. Durán, Federico Gimeno, Benigno Ramos, Ana Serrador and Francisco Fernández-Avilés

Instituto de Ciencias del Corazón (ICICOR). Hospital Universitario de Valladolid. Valladolid. España.

It has been estimated that 15-25% of patients who undergo percutaneous or surgical coronary angioplasty are diabetics. The indications for coronary revascularization and initial results of the procedure do not differ substantially between patients with diabetes mellitus and non-diabetics. However, the long-term results of both percutaneous and surgical coronary angioplasty are less favorable in diabetics in terms of mortality and the need for new revascularization procedures.

The development and widespread use of stents and glycoprotein IIb/IIIa receptor inhibiting drugs have improved the clinical evolution of diabetics treated with angioplasty. Currently available data show that the administration of glycoprotein IIb/IIIa inhibitors to patients undergoing coronary angioplasty is especially useful in diabetics and improves short-term and long-term results, decreasing one-year mortality by 45%. There seem to be indications for the routine use of glycoprotein IIb/IIIa inhibitors in diabetics treated with angioplasty. While the use of stents has improved long-term and short-term results in diabetics, the success rates of angioplasty in diabetics are still lower than in non-diabetics. Diabetes is still an independent predictor of restenosis and long-term events after stenting interventions.

Analysis of the studies comparing percutaneous and surgical revascularization in diabetic patients with multivessel disease shows that surgery is superior in terms of long-term mortality and need for new revascularization procedures. Stenting has improved, but not substantially, the results of multivessel angioplasty in diabetics. Therefore, the indications for angioplasty in multivessel diabetics should be evaluated individually.

Factors that contribute to the less favorable post-angioplasty evolution of diabetic patients are more rapid progression of atherosclerosis and, especially, a higher rate of restenosis. New angioplasty techniques, such as brachytherapy and drug-eluting stents, are likely to significantly improve the results of percutaneous interventions

in diabetics, thus allowing the indications for angioplasty in diabetics to be extended even further in the near future.

**Key words:** *Diabetes. Coronary angioplasty stent. Coronary surgery. Glycoprotein IIb/IIIa inhibitors.*

Full English text available at: [www.revespcardiol.org](http://www.revespcardiol.org)

### Angioplastia coronaria en el paciente diabético. Situación actual y perspectivas futuras

Se estima que un 15-25% de los pacientes en los que se indica angioplastia o cirugía coronaria son diabéticos. Las indicaciones de revascularización coronaria y los resultados iniciales de ésta en los pacientes con diabetes mellitus no difieren sustancialmente de los no diabéticos. Sin embargo, los resultados a largo plazo, tanto de la angioplastia coronaria como de la cirugía, son peores en los diabéticos en términos de mortalidad y de necesidad de nuevos procedimientos de revascularización.

El desarrollo y la expansión del uso del *stent* y de los fármacos que inhiben los receptores IIb/IIIa de las plaquetas han mejorado la evolución clínica de los diabéticos tratados con angioplastia. Los datos disponibles en el momento actual revelan que el uso de los inhibidores de la glucoproteína IIb/IIIa en el contexto de la angioplastia coronaria es especialmente beneficioso en los diabéticos, mejorando los resultados a corto y largo plazos y reduciendo la mortalidad al año en un 45%. Su utilización rutinaria en los diabéticos que son tratados con angioplastia parece indicada. El *stent*, aunque ha mejorado los resultados a corto y largo plazos en los diabéticos, no ha podido igualar los resultados de la angioplastia a los de los no diabéticos. La diabetes continúa siendo un factor predictor independiente de reestenosis y de acontecimientos a largo plazo tras angioplastia con *stent*.

El análisis de los estudios que han comparado la revascularización percutánea y quirúrgica en los pacientes diabéticos con enfermedad multivaso demuestran que la cirugía es superior en términos de mortalidad y necesidad de nuevos procedimientos de revascularización a largo plazo. El *stent* ha mejorado, pero no parece haber cambiado sustancialmente los resultados de la angioplastia multivaso en los diabéticos. Por tanto, las indicaciones

Section sponsored by Laboratorio Dr. Esteve

Correspondence: Dr. J.J. Alonso.  
Servicio de Cardiología. Instituto de Ciencias del Corazón.  
Hospital Universitario de Valladolid.  
Ramón y Cajal, 3. 47011 Valladolid. España.  
E-mail: [jalonso@secardiologia.es](mailto:jalonso@secardiologia.es)

de angioplastia en los diabéticos multivaso deben considerarse de forma individualizada.

Los factores determinantes de la peor evolución post-angioplastia de los pacientes diabéticos son la más rápida progresión de la arteriosclerosis y, sobre todo, la mayor tasa de reestenosis. Las nuevas técnicas de angioplastia, como la braquiterapia o los *stents* liberadores de fármacos que inhiben la hiperplasia intimal es probable que mejoren significativamente los resultados del intervencionismo percutáneo en los diabéticos y permitan, en un futuro próximo, la expansión de indicaciones de angioplastia en éstos.

**Palabras clave:** *Diabetes mellitus. Angioplastia coronaria. Stent. Cirugía coronaria. Inhibidores de la glucoproteína IIb/IIIa.*

## INTRODUCTION

The relation between diabetes and coronary artery disease is marked by two circumstances: the high incidence of coronary artery disease in diabetic patients and its poor prognosis in diabetics compared to non-diabetics. It is estimated that more than 50% of adult diabetics have significant coronary atherosclerosis, a prevalence 10 times greater than that of the general population, which is about 2%-4%.<sup>1-3</sup> In fact, at present diabetes is considered not only a risk factor, but also a marker of cardiovascular disease from the point of view of prevention. Consequently, the recommended prevention interventions are the same for diabetics as for patients with coronary artery disease.<sup>3,4</sup> Diabetes not only increases the incidence of coronary disease, but also contributes to a less favorable prognosis. Thus, cardiovascular mortality is twice as frequent in diabetic men and four times as frequent in diabetic women. In general, if we analyze any subgroup of diabetics with coronary artery disease, they have a worse long-term clinical evolution in terms of cardiovascular morbidity and mortality.<sup>2,3</sup> These factors, together with the high prevalence of diabetes in the industrialized world, have made the association of diabetes and ischemic heart disease a serious public health problem in these countries, a frequent cause of disability among the citizens of these countries, and a major expense for health resources (in the U.S., it has been calculated that diabetics use one-fourth of the Medicare budget).

A topic that has received much attention in recent years is coronary revascularization in diabetic patients. The high incidence of coronary artery disease in diabetics, and its greater extension and severity, mean that coronary revascularization is considered in many of these patients. The proportion of diabetics among patients who undergo revascularization is high and it is calculated that 15%-25% of the patients in which

percutaneous or surgical revascularization is indicated are diabetics.<sup>2,3</sup> The interest in revascularization in diabetics not only is due to its frequency, but also to two other factors. The first is related with the results of both angioplasty and coronary surgery, which are less favorable in diabetics, and the second with the controversy regarding the suitability of percutaneous revascularization for diabetics with multivessel disease.<sup>6-8</sup> This subgroup of patients is the only one in the general population of patients with ischemic heart disease in which percutaneous revascularization is associated with a greater long-term mortality than surgical revascularization.

Nevertheless, the continuous advances that are appearing in the field of coronary angioplasty are improving results, even in unfavorable subgroups like diabetics, and expanding the indications. Paradigms of this progress are the development and dissemination of the use of stents and glycoprotein IIb/IIIa inhibitors, the availability of techniques that inhibit intimal hyperplasia, the main factor responsible for intra-stent restenosis in diabetics, such as intracoronary brachytherapy and, more recently, coated stents that release drugs to inhibit the proliferation of smooth muscle cells. Due to these advances, percutaneous coronary revascularization in diabetic patients is a continuously evolving area of cardiological therapeutics that requires frequent updates. The fundamental aims of this update are to review the indications for revascularization in diabetic patients with coronary artery disease, analyze the results of percutaneous revascularization in diabetics, especially in comparison with surgical revascularization, and evaluate the future expectations for new devices.

## INDICATIONS FOR REVASCULARIZATION IN DIABETIC PATIENTS WITH CORONARY ARTERY DISEASE

The indication for any intervention should be based on an analysis of the evidence of its influence on mortality, the clinical evolution, and short and long-term quality of life. Coronary revascularization has been shown to be useful in reducing mortality and improving quality of life in many clinical situations and diverse angiographic contexts. The indications for coronary revascularization were recently reviewed by the Sociedad Española de Cardiología in various documents, but the specific subgroup of patients with diabetes has not been examined in depth.<sup>9,10</sup>

### Indications for revascularization in diabetic patients with stable chronic ischemic heart disease

Coronary revascularization is a useful tool for improving the prognosis and controlling symptoms in

some groups of patients with coronary artery disease.<sup>11-13</sup> In fact, there is a clear surgical indication in patients with acute angina in spite of adequate medical treatment, regardless of the number of diseased blood vessels. This indication is also valid in diabetic patients, but limited by the fact that, as is well known, the perception of ischemia is sometimes altered by diabetic neuropathy. Diabetics with asymptomatic acute ischemia or atypical clinical manifestations are not uncommon.<sup>2,3</sup> For that reason, the evaluation of revascularization indications in asymptomatic or mildly symptomatic patients is especially interesting. These indications have been analyzed in many studies. The overall evidence obtained from evaluating these studies has allowed scientific societies and expert groups to issue recommendations on the indications for revascularization based on the clinical manifestations and anatomy of each patient.<sup>9-15</sup> Generally speaking, benefits have been shown in terms of mortality in patients with significant disease of the coronary trunk, three vessels, or one or two vessels with involvement of the proximal portion of the anterior descending coronary artery, especially if left ventricular ejection fraction is abnormal or there is extensive ischemia. The validity of these studies has been questioned because they were carried out in the 1980s. The surgical or angioplasty technique used in most cases, and especially the medical treatment, were substantially different from those now used. Nevertheless, because of the consistency of the results they are still considered applicable and valid.<sup>14,15</sup>

The morbidity and mortality of surgery has always been found to be higher in diabetics compared with non-diabetics.<sup>16,17</sup> However, although most studies did not specifically analyze the benefit of revascularization in diabetic patients, they did demonstrate that the effectiveness of revascularization was always greater in patients with more extensive coronary artery disease and/or a more impaired left ventricular function. Indeed, among diabetics there is a greater proportion of patients with these characteristics and, therefore, a larger number of patients than could potentially benefit from surgery. Thus, in 317 diabetic patients over the age of 65 years in the CASS study,<sup>16</sup> after a mean follow-up of 13 years the mortality was 57% greater than in the 1843 matched non-diabetics. Nevertheless, the benefit of surgery in relation to medical treatment was similar to that of non-diabetics, so surgery was associated with a reduction in mortality of 44%. To summarize, there is a general consensus that surgical revascularization prolongs the life of diabetics with acute coronary disease so, consequently, it seems reasonable to order examination to detect extensive ischemic heart disease in diabetics, in which it is common.

## Indications for revascularization in diabetic patients with acute coronary syndrome without ST-segment elevation

The management of acute coronary syndrome without elevation of the ST segment has recently changed as a result of the publication of two large clinical trials that have compared an aggressive strategy, consisting of coronariography and early revascularization, with conservative management guided by ischemia. In both studies (FRISC II and TACTIS-TIMI-18)<sup>18-21</sup> an early invasive strategy in patients with acute coronary syndrome without ST-segment elevation reduced the incidence of major cardiac events at 1 year and at 6 months, respectively. The clinical impact of the two studies has been so important that it has forced both the ACC/AHA and Sociedad Española de Cardiología to update their guidelines for clinical practice in the management of acute coronary syndrome.

In the FRISC II study,<sup>18,20</sup> 2457 patients with acute coronary syndrome without ST-segment elevation, with ST-segment depression or an increase in CK-MB or TnT ( $>0.10$ ) were randomized to receive either an invasive strategy (1222 patients) consisting of catheterization and revascularization in the first 7 days, or noninvasive management in which revascularization was indicated for refractory angina or ischemia in the exercise stress test. Of the 1222 patients assigned to invasive therapy, angioplasty was performed in 522 patients (43%) (a stent was used in 61% of the procedures and abciximab in 10%) and surgery in 430 patients (35%). Within one year, the invasive strategy was associated with a reduction in mortality of 43% (RR, 0.57; 95% CI, 0.36-0.90) and in myocardial infarction of 26% (RR, 0.74; 95% CI, 0.59-0.94). In the FRISC II study, 298 patients (12%) were diabetics (154 in the invasive group and 144 in the conservative group). The invasive strategy reduced the 1-year mortality in diabetics by 38% (RR, 0.62; 95% CI, 0.31-1.24) and the incidence of death or infarction by 30% (RR, 0.70; 95% CI, 0.47-1.04). These figures were similar to those observed in non-diabetics. In addition, in diabetics that received invasive treatment a decrease in the need for medication and improvement in symptoms was observed at 6 months (RR, 0.59; 95% CI, 0.49-0.64). In summary, the findings of this study are consistent with the idea that invasive management that includes revascularization, if indicated and feasible, for high-risk acute coronary syndromes is clinically more beneficial than the classic conservative strategy guided by the presence of ischemia, even in diabetic patients.

The findings of the other study (TACTIS)<sup>21</sup> confirm those of the previous one. This clinical trial included 2220 patients with acute coronary syndrome (unstable angina or acute myocardial infarction without ST-segment elevation) and high risk (positive markers of myocardial damage, changes in the ST segment or

T wave, or previous history of coronary artery disease). All were treated with the glycoprotein IIb/IIIa inhibitor tirofiban and then assigned randomly, as in the FRISC II study, to either a non-invasive strategy guided by the development or detection of ischemia or invasive management consisting of catheterization and revascularization in the first 48 h. Of the 1114 patients assigned to the interventional group, 41% underwent angioplasty and 20%, coronary surgery. The incidence of death or reinfarction at 6 months decreased by 26% and the rate of occurrence of the main endpoint (death, reinfarction, or readmission for acute coronary syndrome) also decreased by 22% (OR, 0.78; 95% CI, 0.62-0.97). Of the 2220 patients included, 28% were diabetic. In these patients, the incidence of death, reinfarction, or readmission for acute coronary syndrome decreased by 28% in the patients assigned to the interventional strategy (28% in the invasive group and 20% in the non-invasive group;  $P < .05$ ). In non-diabetic patients, the reduction was not significant (16.4% vs 14.2%). That is to say, the benefits of the study were observed mainly in diabetic patients. The conclusions of this study are consistent with the above results and confirm that diabetics with acute coronary syndrome benefit from a strategy that includes early appropriate revascularization. It is more complicated to decide on the type of revascularization, surgery, or angioplasty in the context of multivessel disease, as we will see below. In general, and with the present techniques, angioplasty is recommended in single-vessel disease and surgery in multivessel disease, assuming that at least the anterior descending coronary can be revascularized with mammary artery and the surgical risk is not raised by coexistent comorbidity factors.

### Indications for revascularization in diabetic patients with acute coronary syndrome and ST-segment elevation

The results of the DANAMI II study,<sup>22</sup> which have been recently communicated, have confirmed the findings of the meta-analyses of Weaver et al<sup>23</sup> and Michels et al,<sup>24</sup> which demonstrated that primary angioplasty is the reperfusion method of choice in patients with acute myocardial infarction and ST-segment elevation whenever it can be performed quickly (within 3 h) by an experienced team. This and other studies,<sup>25,26</sup> like the recently published CADILLAC study,<sup>25</sup> have clarified doubts about the effectiveness of stents that emerged with the results of the Stent-PAMI study,<sup>27</sup> in which stent implantation was not found to be more beneficial than balloon angioplasty. The results of primary angioplasty with stent implantation in the DANAMI II study were better than those of thrombolytic treatment. In the CADILLAC study, angioplasty with a stent was associated with a lower need for new revascularization procedures during fo-

llow-up.

There is little information about the results of primary angioplasty in diabetic patients. The reason for this probably lies in the fact that most of studies that have compared primary angioplasty with thrombolysis have been small, which has precluded the analysis of subgroups. The only study that has evaluated primary angioplasty in diabetics is the angioplasty substudy of GUSTO IIb.<sup>28</sup> In this clinical trial, 177 diabetics and 961 non-diabetics with acute myocardial infarction were distributed randomly to primary angioplasty or thrombolysis with the accelerated alteplase schedule. The diabetics had clinical and angiographic characteristics (more severe stenosis and worse flow) that were less favorable than those observed in non-diabetics. Nonetheless, the success rate of the procedure in patients assigned to angioplasty was similar (diabetics 70%, non-diabetics 72%). The clinical results at one month of diabetics and non-diabetics were similar, so the incidence of death, reinfarction, or disabling stroke was reduced by 38% (OR, 0.62; 95% CI, 0.41-0.96) in non-diabetics and 30% in diabetics (OR, 0.70; 95% CI, 0.29-1.72). Therefore, the available data, although scant, suggest that primary angioplasty is as effective in diabetics as in non-diabetics, and better than thrombolysis.

Although diabetics treated with primary angioplasty and stent had a less favorable clinical evolution at 6 months<sup>29,30</sup> than non-diabetics, and diabetes has been demonstrated to be an independent predictive factor of thrombotic stent occlusion,<sup>31</sup> the data available for primary angioplasty with stent confirm the usefulness of this device in diabetics with acute myocardial infarction. The incidence of cardiac events (death, reinfarction, disabling stroke, and need for revascularization of the treated vessel) at 6 months in the 135 diabetic patients included in the STENT-PAMI study<sup>29</sup> was lower in the patients assigned to stent implantation than in those assigned to balloon angioplasty (20% vs 30%), especially in non-insulin-dependent patients (12% vs 28%;  $P = .04$ ).

Therefore, given the information available, it can be said that primary angioplasty with stent implantation is the reperfusion strategy of choice in diabetic patients with acute myocardial infarction and ST-segment elevation. In these cases, aside from exceptions, there is no doubt that angioplasty should be performed after coronary angiography, if possible with stent implantation in the causal lesion. Surgery should only be considered in patients in which the artery responsible for infarction is not susceptible to angioplasty, has a large dependent myocardial territory, and can be performed immediately. Scant information is available regarding the best strategy in cases in which angioplasty of the causal artery only achieves partial revascularization. Depending on individualized analysis of each case and the experience of each group, revasculariza-

tion may be completed by angioplasty in the same act or in a second stage, patients may be controlled with medical treatment, or revascularization may be completed surgically in a second intervention.

### GLYCOPROTEIN IIB/IIIA INHIBITORS IN ANGIOPLASTY IN DIABETICS


The development and availability of drugs that block platelet IIB/IIIA receptors is an important advance in the pharmacological treatment of angioplasty. The administration of these potent platelet antiaggregants in angioplasty has substantially improved its short and long-term results, which is why its use in angioplasty is increasing. The group of patients that probably have benefited most from glycoprotein IIB/IIIA inhibitors are diabetics, both in the case of angioplasty in a stable context and in acute coronary syndrome. Currently, three drugs of this group are used: abciximab, the best known and most often used, eptifibatide, and tirofiban.

Abciximab was first used in the context of coronary angioplasty and its effectiveness has been confirmed in several studies that have involved a large number of patients in diverse clinical contexts of ischemic heart disease. The results obtained in diabetic patients have been analyzed in a meta-analysis<sup>32</sup> of the three main clinical trials (EPIC, EPILOG, and EPISTENT). The EPIC Study included 2099 patients in which high-risk angioplasty was indicated. It was considered a high-risk intervention because it was carried out in the context of unstable angina, acute myocardial infarction in evolution, or in lesions with a complex morphology. EPILOG recruited 2792 patients undergoing elective or emergency angioplasty and EPISTENT included 2399 patients who were randomized to treatment with abciximab as well as the angioplasty technique (stent or balloon). The three studies included a total of 1462 diabetics and the results revealed a clear benefit of treatment with abciximab: the 1-year mortality was 4.5% in the patients who were assigned randomly to abciximab and 2.5% in patients assigned to the control group ( $P=.03$ ). That is to say that the use of abciximab in diabetics was associated with a reduction in the 1-year mortality of 45%. This reduction in mortality was greater than that obtained in the overall group (diabetics and non-diabetics), which was 36% when adjusted to a Cox regression model (hazard ratio [HR], 0.64; 95% CI, 0.46-0.90;  $P=.01$ ). A reduction in mortality with abciximab was observed in 462 diabetics, whether insulin-dependent (abciximab, 4.2%, placebo, 8.1%;  $P=.07$ ) or non-insulin-dependent (abciximab, 1.8%, placebo, 2.7%;  $P=.34$ ). Nevertheless, the most interesting finding was that the reduction in mortality was especially important in two diabetic subgroups of interest: diabetics undergoing multivessel percutaneous revascularization (abciximab, 0.9%, placebo, 7.7%;

$P=.02$ ) and diabetics with insulin resistance syndrome (abciximab, 2.3%, placebo, 5.1%;  $P=.04$ ). Two findings underline the special benefit of abciximab in diabetics: in diabetics undergoing angioplasty and treated with abciximab (2.5%), the 1-year mortality equaled that of non-diabetics treated with placebo (2.6%). If diabetics are excluded from the meta-analysis, the difference in mortality between those treated with placebo or abciximab is not significant (2.6% vs 1.9%;  $P=.1$ ). Therefore, it seems that the reduction in mortality in the patients included in the meta-analysis treated with abciximab takes place fundamentally as a result of the benefit that it produces in diabetics (Figure 1). Abciximab also reduced the incidence at one year of the endpoint formed by mortality, the incidence of myocardial infarction, or the need for a new revascularization of the treated vessel in 15% (abciximab, 29%, placebo, 34%;  $P=.02$ ). This reduction took place at the expense of a decrease in mortality and the incidence of infarction.

The good results of prophylactic treatment with abciximab in diabetic patients in the context of angioplasty have also been confirmed with glycoprotein IIB/IIIA inhibitors that have a structure different from that of abciximab, like tirofiban, eptifibatide, or even lamifiban. In the context of acute coronary syndrome without ST-segment elevation, a meta-analysis of six studies that analyzed the influence of the use of glycoprotein IIB/IIIA inhibitors on the clinical evolution at 30 days has been published recently.<sup>33</sup> Of the 6458 diabetic patients included in the six studies, 1279 were treated with coronary angioplasty. The associated use of glycoprotein IIB/IIIA inhibitors in these patients with acute coronary syndrome without ST-segment elevation treated with coronary angioplasty reduced the 30-day mortality from 4% to 1.2%, that is to say, by 70% (OR, 0.30; 95% CI, 0.14-0.69). The results were homogeneous in all the studies, which all showed that the use of these antiaggregant drugs in diabetics treated with angioplasty was associated with a decrease in mortality, as can be seen in Table 1.

The use of glycoprotein IIB/IIIA inhibitors, specifically abciximab, in the context of acute myocardial infarction treated with primary stent angioplasty in diabetic patients has not been specifically analyzed in adequate studies. In the general population, the results of a recent meta-analysis<sup>34</sup> of the three main studies of angioplasty with stent implantation, associated or not with abciximab (CADILLAC, ISAR II, and ADMIRAL), suggest that their use is beneficial, reducing the incidence of adverse cardiac events in the intermediate term. The incidence of death or reinfarction at 6 months was 3.9% vs 7.1% in those not treated with abciximab (OR, 0.53; 95% CI, 0.34-0.84). It is likely that the results in the subgroup of diabetics are at least similar to those of the general population with acute myocardial infarction.



**Fig. 1.** One-year mortality in diabetic and non-diabetic patients treated with angioplasty, depending on whether or not they received abciximab and stent implantation (taken from Bhatt et al<sup>32</sup>).

In the context of stent angioplasty, eptifibatide has demonstrated its effectiveness in reducing events in the short, intermediate, and long term in both acute coronary syndrome and stable ischemic heart disease. The meta-analysis of Bhatt et al,<sup>32</sup> extensively commented above, compiled information about the diabetics in the EPIC, EPILOG and EPISTENT studies. This meta-analysis demonstrated that the association of abciximab to angioplasty with stent implantation reduced the 1-year mortality in diabetics from 4.6% to 1.3% ( $P=.04$ ), which equalized the mortality in diabetics and non-diabetics (1.3%) (Figure 1). In the context of scheduled stent angioplasty, the ESPRIT study (Enhanced Suppression of the Platelet IIb/IIIa Receptor with Integrilin Therapy) demonstrated the effectiveness of eptifibatide in reducing events at 48 h, 30 days, 6 months, and 1 year.<sup>35-37</sup> This study included

2064 patients in which angioplasty was scheduled, who were assigned randomly to placebo or eptifibatide treatment (which began before stent implantation and was maintained for 48 h). The results were consistent with those of previous studies, so that the incidence of death or reinfarction decreased by 37% at 1 year in patients treated with eptifibatide (HR, 0.63; 95% CI, 0.48-0.83). The incidence of the combined endpoint of mortality, non-fatal infarction, or revascularization of the treated lesion was significantly lower in patients treated with glycoprotein IIb/IIIa inhibitors (17.5% vs 22.1%; HR, 0.76; 95% CI, 0.63-0.93;  $P=.007$ ). This reduction in events was greater in diabetics (a 32% reduction at 1 year in the incidence of death or reinfarction in non-diabetics and of 42% in diabetics). However, because of the sample size (only 419 patients were diabetics) the difference was not statisti-

**Table 1. Thirty-day mortality in diabetic patients with acute coronary syndrome treated with coronary angioplasty with or without glycoprotein IIb/IIIa inhibitors associated**

Study	No. of patients	Randomized	Randomized to GP IIb/IIIa	P
PURSUIT	457	3.3%	2.4%	.57
PRISM	147	2.5%	0%	.5
PRISM-PLUS	107	1.8%	0%	1
GUSTO IV	239	6.5%	1.2%	.04
PARAGON A	45	7.1%	0%	.31
PARAGON B	284	4.3%	0.7%	.06
Total	1274	4.0%	1.2%	.002

GP IIb/IIIa indicates glycoprotein IIb/IIIa inhibitors. Taken from Roffi et al.<sup>33</sup>

cally significant.

The combination of the data currently available demonstrates that the use of glycoprotein IIb/IIIa inhibitors in the context of coronary angioplasty is especially beneficial in diabetic patients, improving the clinical results in the short, intermediate, and long term. In diabetics, they reduce the incidence of events at 1 year of angioplasty almost twice as much (78% more) as in non-diabetics, so there now seems to be no doubt about the need for its routine use in diabetic patients. The spread of the use of these drugs has undoubtedly been one of the most important advances to help improve the results of percutaneous revascularization in diabetics.

## RESULTS OF CORONARY ANGIOPLASTY IN DIABETICS

### Results of coronary balloon angioplasty

Percutaneous revascularization in diabetics has been an object of study since the technique was first used in 1977. The results of all the studies have coincided. In the main studies that have analyzed balloon angioplasty in diabetics, it was observed homogeneously that the clinical and angiographic profile of the diabetics was unfavorable in relation to that of non-diabetics. These less favorable clinical and angiographic characteristics do not seem to influence the angiographic success rates, which were similar in diabetics and non-diabetics. Nevertheless, the initial results and, above all, the long-term results were worse in diabetic patients.

In the Registry of the National Heart, Lung and Blood Institute,<sup>38</sup> the results of balloon angioplasty were specifically compared between 281 diabetics and 1833 non-diabetics. The diabetics were older, had more comorbidity, were more often women, and had more extensive and diffuse coronary disease. The initial angiographic success rate and degree of revascularization (complete vs incomplete) were similar, but the incidence of the endpoint formed by mortality, infarction, or the need for urgent revascularization during hospitalization was 11% in diabetics and only 6.7% in non-diabetics ( $P < .001$ ). The mortality and incidence of infarction in the hospital phase were greater in diabetics, especially diabetic women, who had a mortality of 8%. The long-term evolution (Table 2) was clearly unfavorable in diabetics: the adjusted mortality at 9 years was twice as high in diabetics (36% vs 18%; RR, 1.82; 95% CI, 1.41-2.34), the incidence of infarction was 60% higher (29% vs 19%), and the need for new revascularization procedures was greater in diabetics (33% more required new angioplasty and 19% more needed coronary surgery in the follow-up than in non-diabetics). The second classic study analyzed the experien-

ce of Emory University.<sup>39</sup> At this center, in the period between 1980 and 1990, elective angioplasty was performed in 1133 diabetic and 9300 non-diabetic patients. As in the previous study, the diabetics were older and more frequently women than the non-diabetics; likewise, their functional class was worse and the history of previous infarction, multivessel disease, hypertension, obesity, or heart failure were more frequent in the diabetics. The clinical and angiographic results after angioplasty were good and similar in diabetics and non-diabetics. There was a low incidence of major adverse events (3%), although a tendency to a greater mortality and the development of infarction was detected in patients with insulin-dependent diabetes. The situation was clearly different in the long term (5 years). Survival was significantly lower in diabetics (89% vs 93%), as was the percentage of patients who did not develop infarction (81% vs 89%) or require coronary surgery (77% vs 86%) or new angioplasty (57% vs 68%) during follow-up. At 5 years, only 36% of the diabetics remained alive without presenting an infarction or requiring revascularization. The studies made in patients with multivessel disease have demonstrated similar findings: a similar mortality and rate of intrahospital events between diabetic and non-diabetic patients, but a greater long-term incidence of cardiac events and mortality (Table 2). Thus, for example, in the BARI study 170 diabetics with multivessel disease and 734 non-diabetics were randomized to angioplasty. The intrahospital mortality was similar, 0.6% for diabetics and 1.2% for non-diabetics, as was the incidence of death or infarction (diabetics, 2.4%; non-diabetics, 3.1%). Nevertheless, at 5 years the mortality was 5 times greater in diabetics (20.6% vs 4.8% in non-diabetics). The rate of myocardial infarction was 37% greater in diabetics during follow-up. It seems clear, therefore, that similar initial results can be expected from angioplasty of diabetics and non-diabetics, but the long-term clinical evolution is much less favorable in terms of the incidence of infarction, need for new revascularization procedures, and mortality.

The question that immediately arises after knowing the natural history of angioplasty in diabetics is: Is this unfavorable evolution due to a very high rate of restenosis or to a more rapid progression of arteriosclerosis in the untreated segments than in non-diabetics? In the light of the present evidence, both phenomena are involved in the mechanisms that explain the less favorable prognosis of angioplasty in diabetics.<sup>3</sup> Different studies have consistently found higher rates of clinical and angiographic restenosis in diabetics. Rozenman et al<sup>44</sup> observed an incidence of restenosis of 35% in non-diabetics and 61% in insulin-dependent diabetics. In the angiographic analysis of 320 patients in the BARI study at 5 years,<sup>45</sup> the incidence of restenosis was 27% in lesions in non-diabetics and

TABLE 2. Long-term prognosis, in terms of mortality, of diabetic patients treated with angioplasty

Study and citation	Extension of CAD (no. of vessels)	Diabetics		Non-diabetics		Follow-up years
		No. of patients	Mortality, %	No. of patients	Mortality, %	
NHLBI <sup>38</sup>	3	78	28	325	11	5
	2	75	16	570	10	5
	1	121	14	904	6	5
	Total	274	18.5	1799	8.4	5
Stein (Emory) <sup>39</sup>	1-3	1133	12	9300	7	5
CABRI <sup>40</sup>	2-3	125	23	929	9	4
EAST <sup>41</sup>	2-3	29	10	169	12	5
BARI <sup>42</sup>	2-3	170	35	734	9.5	5
Duke University <sup>43</sup>	2-3	144	24	560	12	5

CAD indicates coronary artery disease.

43% in patients with diabetes ( $P=.01$ ). In addition, the available data demonstrate that the progression of the disease is greater in healthy or significantly injured segments in diabetics. In the last commented study, Barsness et al<sup>45</sup> detected a greater rate of appearance of significant new injuries at 5 years in diabetics (3% vs 2%;  $P=.002$ ). Analysis of the patients of the CABRI study<sup>46</sup> demonstrated that the degree of revascularization and extension of coronary disease were similar in diabetics and non-diabetics. Consequently, this factor was not responsible for the less favorable evolution of diabetics. The investigators of this study could only attribute the evolution to a greater progression of coronary disease in the native arteries or grafts. The investigators from the EAST study reached similar conclusions.<sup>41</sup> In this study, the survival curves of diabetic and non-diabetics patients treated with angioplasty separated later (as of 5 years), suggesting that the progression of coronary disease was responsible in large measure for the worse prognosis of diabetic patients.

Therefore, balloon angioplasty in diabetic patients is associated with less favorable intermediate and long-term results than angioplasty in non-diabetics. This is due to a greater incidence of restenosis and a greater progression of coronary arteriosclerosis in untreated segments.

### Results of coronary stent angioplasty


The stent is considered the greatest advance in coronary angioplasty since Grüentzig performed the first angioplasty in 1977. The stent has been shown to reduce the incidence of restenosis in a large variety of lesions and to improve the short, intermediate, and long-term clinical results of coronary angioplasty.<sup>47-57</sup> Consequently, stent angioplasty has become the most frequently used percutaneous coronary revascularization technique. The findings of the registry of the Working Group of Hemodynamics of the Sociedad

Española de Cardiología reported for 2000 indicated that stents have been used in Spain in at least 22 580 angioplasties (77% of the total). This expansion is due not only to its good clinical results but, above all, to the fact that it is a very operative technique.

What has the stent contributed to angioplasty in diabetics? The stent has improved the initial and long-term results and has reduced the incidence of restenosis in relation to conventional angioplasty. Nevertheless, it has not managed to achieve the same results as obtained in non-diabetics: the rate of restenosis and incidence of long-term events, although lower than with balloon angioplasty, are still high and the results are less favorable than expected.

The subanalysis of 92 diabetics in the STRESS I and II studies<sup>59</sup> compared the clinical effectiveness and rate of restenosis at 6 months between patients randomized to balloon angioplasty or stent angioplasty. It demonstrated that stents in diabetics are associated to a greater initial rate of angiographic success, with a greater initial gain (1.61 mm vs 1.06 mm;  $P<.001$ ) and success rate (stent, 100%; balloon, 82%;  $P<.01$ ). In the long term, stents were accompanied by a lower incidence of restenosis at 6 months (24% vs 60%;  $P<.01$ ) or need for new revascularization of the treated vessel (13% vs 31%;  $P=.03$ ). The experience obtained in a non-randomized study by the French group of Lille<sup>60</sup> was similar; stent implantation reduced the incidence of restenosis from 63% to 25% and the rate of complete occlusion from 14% to 2%. The findings are similar for Spanish studies. Navarro et al<sup>61</sup> demonstrated that the use of stents reduced the incidence of clinical restenosis more than two-fold. Nevertheless, except in isolated studies,<sup>62</sup> the rate of restenosis in diabetics continued to be high in unselected patients and diabetes is still an independent predictive factor of restenosis.<sup>63-67</sup> Elezi et al<sup>63</sup> observed an incidence of restenosis in diabetics of 37.5% vs 28.3% in non-diabetics ( $P<.001$ ). Kastrati et al<sup>64</sup> demonstrated in a series of 1349 patients that diabetes increased the risk of re-





**Fig. 2.** Incidence of restenosis at 6 months and independent predictive factors of angiographic restenosis after stent angioplasty in the series of Alonso et al.<sup>65</sup>

nosis by 86% (OR, 1.86; 95% CI, 1.56-2.16) and the need for a new revascularization procedure by 45% (OR, 1.45; 95% CI, 1.11-1.80). In our experience,<sup>65</sup> an angiographic re-evaluation was performed at 6 months and patients were clinically followed up (median of 4 years). In the cohort of 981 consecutive patients treated with stent implantation, including 128 diabetics, the adjusted incidence of angiographic restenosis was 38% greater in diabetics, with a restenosis rate of 39% in diabetics and 26% in non-diabetics ( $P=.008$ ). As in the study by Kastrati et al, diabetes was a strong independent clinical predictor of restenosis (Figure 2).


From a clinical point of view, diabetics have a less favorable evolution than non-diabetics in the intermediate and long term. At 5 years, the incidence of clinical events in our series<sup>67</sup> was 39% higher in diabetics (HR, 1.39; 95% CI, 1.02-1.90). Diabetes, along with hypertension and multivessel disease, was an independent predictor of the long-term evolution (Figura 3). The results of other studies are similar. At 1 year, the diabetics analyzed by Elezi et al<sup>63</sup> had a lower event-free survival than non-diabetics (73.1% vs 78.5%;  $P<.01$ ). The mortality after a mean follow-up of 18#±10 months was 4 times greater in diabetics (8.1% vs 2.1%;  $P=.03$ ) and the need for revascularization of the treated lesion was 35% greater in the series of Paschal et al.<sup>68</sup>

The influence of the type of diabetes (type 1 vs type 2) on the results of stent angioplasty has also been analyzed. Although Schofer et al<sup>66</sup> did not find differences in the incidence of angiographic restenosis, the Washington group, in the largest study published,<sup>69</sup> observed that the clinical evolution of non-insulin-dependent diabetic patients is similar in the short and intermediate term to non-diabetics. However, insulin-dependent diabetics had clearly worse short and intermediate results than non-diabetics.

## MECHANISM OF RESTENOSIS IN DIABETICS

Restenosis, together with the frequent impossibility of achieving complete revascularization of all ischemic territories and faster progression of arteriosclerosis, have been identified as factors in the less favorable clinical evolution of diabetics treated by balloon angioplasty or stent implantation. Restenosis has a key role. This complication is conditioned by three phenomena: the immediate elastic recoil, unfavorable late remodeling, and intimal hyperplasia. In particular, hyperplasia is the predominant mechanism of restenosis in angioplasty with stent implantation.<sup>2,70,71</sup> Reduction of the incidence with stenting seems to be due fundamentally to its capacity to prevent immediate elastic recoil and achieve favorable arterial wall remodeling.<sup>70,71</sup> Nevertheless, the implantation of a stent increases intimal hyperplasia, the third mechanism involved in the restenotic process, which becomes the main mechanism responsible for restenosis in lesions with stents. The final balance of the three effects in most patients treated by stent implantation is a larger final and long-term luminal diameter and a lower incidence of restenosis.

Metabolic, hematological, and biological abnormalities inherent to diabetics probably participate in the complex combination of mechanisms that occur after vascular aggression and are involved in restenosis.<sup>2</sup> In diabetics it has been possible to demonstrate by intravascular ultrasound studies that restenosis is due fundamentally to intimal hyperplasia, which is greater than in non-diabetics in both balloon angioplasty and stenting.<sup>72</sup> Therefore, two factors that increase intimal hyperplasia coincide in diabetics: diabetes *per se* and stent implantation, which is why the final result might not be as favorable as in non-diabetics. The increased hyperplasia in diabetic patients can be measured by the hyperinsulinemia, as determined by the insulin resistance of type 2 diabetics,<sup>73</sup> and other metabolic disorders of diabetics that involve the re-



**Fig. 3.** Incidence of major cardiac adverse events (MCAE) in the follow-up angioplasty with stent implantation: mortality, infarction, need for revascularization or admission for acute coronary syndrome. Predictors of MCAE in the follow-up and MCAE-free survival<sup>67</sup>.

nin-angiotensin system, as well as biological disturbances induced by hyperglycemia *per se*. Another mechanism involved is the greater probability of diabetics to develop thrombotic phenomena and occlusion of the treated lesion. Hyperinsulinemia, lipid disorders, and disturbances in the activity of PAI-1, which are common in diabetics, favor the progression of arteriosclerosis, intimal hyperplasia, and thrombotic disturbances.<sup>2</sup> Although much remains to be learned about the pathophysiology of restenosis in diabetics, the available information suggests that stent-based methods that reduce intimal hyperplasia should be sought. This would lead to less restenosis and, undoubtedly, a more favorable clinical evolution than what can now be expected.

### REVASCULARIZATION IN THE DIABETIC PATIENT WITH MULTIVESSEL DISEASE. RESULTS OF ANGIOPLASTY AND CORONARY SURGERY

Advances in angioplasty in the 1980s have made it possible to approach patients with multivessel disease percutaneously. At this time, a series of clinical trials were designed and carried out to compare coronary balloon angioplasty and surgery in patients with 2 or

3-vessel disease or with acute stenosis of the proximal portion of the anterior descending artery.<sup>74-82</sup> The results of these studies, considered together in a meta-analysis published in 1998,<sup>14</sup> demonstrated that the short and long-term results were similar in terms of mortality and the incidence of non-fatal infarction. Nevertheless, the need for new revascularization procedures was much greater in patients assigned randomly to angioplasty. These conclusions seem valid for all the subgroups of patients with multivessel disease, except diabetics.<sup>83</sup> The BARI study<sup>78,83</sup> showed that the 5-year mortality in diabetics assigned to revascularization by angioplasty was almost 4 times greater (20.6% vs 5.8% in those assigned randomly to surgery). Although the design of this analysis in the subgroup of diabetics was carried out at the end of the patient inclusion period, which is a limitation, it ignited an intense controversy that still goes on and leads many cardiologists to advise against angioplasty in diabetics with multivessel disease.<sup>7</sup>

What was learned in the BARI study was used to publish many papers that analyzed the problem parting from the data of clinical trials or cohort studies (Table 3). The majority confirmed that the prognosis was worse in diabetic patients with multivessel disease treated with angioplasty. The largest experiences come from two non-randomized studies made by the University of Emory<sup>84</sup> and the Northern New England Cardiovascular Study Group.<sup>85</sup> In the first study, the adjusted mortality in a 10-year follow-up was 35% greater in insulin-dependent diabetic patients with multivessel disease treated with angioplasty (63%) than in those who underwent coronary surgery (53%; P=.04). In the second study, the mortality was 49% greater (HR, 1.49; 95% CI, 1.02-2.17) in diabetics who underwent angioplasty than in those who underwent surgery. The analysis of the 125 diabetics of the CABRI study<sup>40</sup> demonstrated that the 4-year mortality was almost two-fold greater in the patients randomized to percutaneous revascularization than to surgery (22.6% vs 12.5%; RR, 1.81; 95% CI, 0.80-4.08). In the 8-year follow-up of 90 diabetic patients of the EAST study,<sup>41</sup> it was observed that the survival of diabetics assigned to angioplasty was significantly lower than in those randomized to aortocoronary bypass surgery (60.1% vs 82.6%; P=.02). The data from the 7-year follow-up of the BARI study<sup>42</sup> also confirmed the better prognosis in terms of survival of diabetics assigned to surgery (76.4%) vs angioplasty (55.7%; P=.0011). Only two large cohort studies demonstrated a similar 5-year mortality in diabetics with multivessel disease treated with either angioplasty or surgery. These two studies are the analysis of the Duke University database<sup>43</sup> and the registry of the BARI study,<sup>86</sup> which included patients who met the inclusion criteria of this clinical trial and were registered and followed up. Their treatment was not assigned randomly

Table 3. Long-term mortality in diabetics with multivessel disease treated with angioplasty or coronary surgery

Study	Study type	Angioplasty		Surgery		Follow-up, years
		No.	Mortality, %	No.	Mortality, %	
Barsness et al <sup>43</sup>	Observational	144	31	626	29	5
BARI <sup>42</sup>	Randomized	173	44	180	24	7
BARI <sup>86</sup>	Registry	182	14	117	15	5
EAST <sup>41</sup>	Randomized	29	40	30	25	8
CABRI <sup>40</sup>	Randomized	62	23	63	12	4
Weintraub et al <sup>84</sup>	Observational	834	22	1805	24	5
Niles et al <sup>85</sup>	Registry	1593	6	4066	5	5

but established by the cardiologist and/or surgeon with the patient. The explanation of this apparent paradox is based on the main characteristic that differentiates the design of the two types of studies: patient selection. Whereas in the BARI clinical trial the revascularizing strategy was assigned randomly, in the patients of the BARI registry and Duke University database, the physician and surgeon of each patient selected the treatment that seemed most suitable for each case. This methodological bias explains the difference and is an interesting clinical finding. In at least 2 out of 4 observational studies, adequate patient selection could ensure acceptable long-term results.

Analysis of these data has some limitations, principally related to the angioplasty technique (stents and glycoprotein IIb/IIIa inhibitors were not used) and surgery (little use of arterial grafts). These points could be relevant. For example, in the BARI study, the benefit observed in surgery was obtained only in patients in which internal mammary artery was used.<sup>78,83</sup> Nonetheless, the data available from studies in which effective techniques from current practice have been used do not differ substantially from the information obtained from references for older trials and registries. The ARTS Study<sup>87</sup> of 1205 patients with multivessel coronary disease compared the strategy of stent angioplasty (600 patients, including 112 diabetics) with coronary surgery (605 patients, including 96 diabetics). The intrahospital evolution of 208 diabetics revealed no significant differences between the patients randomized to surgery and angioplasty with stent implantation, which showed a similar intrahospital survival free of coronary events (surgery, 91%; stent angioplasty, 90%;  $P=.7$ ). Nevertheless, at 1 year the need for new revascularization procedures was significantly greater in the patients assigned to stent implantation (22% vs 3.1%;  $P<.001$ ), with a tendency to a greater mortality (6.3% vs 3.1%). The event-free survival of the patients assigned randomly to stent implantation was 63% vs 84% ( $P<.001$ ) for those randomized to surgery. Diabetes continued to be an independent predictor factor of events in this study, which were twice as frequent at 1 year (OR, 2.1;  $P=.001$ ).

While awaiting the long-term results of the ARTS and other studies that include not only stent implantation but also glycoprotein IIb/IIIa inhibitors in the protocol, it seems clear that, in general, the long-term results of coronary revascularization in diabetics are better when surgical procedures are used instead of angioplasty.

### CONTROVERSY REGARDING INDICATIONS FOR CORONARY ANGIOPLASTY IN DIABETICS

Based on the above discussions, in what diabetics can coronary angioplasty be indicated? We know that angioplasty in diabetics has initial results that are very similar to those of non-diabetics. Even in patients with multivessel disease, the short-term results do not differ from those of surgery. Nevertheless, even with stenting, coronary angioplasty in diabetics is associated with a less favorable long-term evolution than in non-diabetics and to worse results than surgery in multivessel patients. These worse results are due, in good measure, to the greater incidence of restenosis. We also know that intra-stent restenosis is greater when other factors are associated, like small vessel size and the presence of residual stenosis and long lesions.<sup>64,65</sup> In general, the sum of the factors that independently influence an event result in an effect of much greater magnitude than the expected sum of the factors. This also occurred in our case. Thus, in the study by Elezi et al<sup>63</sup> the incidence of restenosis in non-diabetics was 28% and in diabetics, 38%. However, if we analyze the rate of restenosis in diabetics, it is evident that it was 49% in those with an arterial diameter of less than 3 mm, but only 27% in diabetics with an arterial diameter of 3 mm or more, which was similar to the overall result in non-diabetics. Another factor that we must be aware of is that the results of angioplasty are always worse in insulin-dependent diabetics.<sup>69,84</sup>

Therefore, it seems reasonable to perform angioplasty in diabetics who have an indication, as described in the second section of this article, and stable di-

sease or unstable single-vessel disease, in which surgery does not seem reasonable except for special anatomic circumstances. In patients with multivessel disease, the decision must be individualized,<sup>2,70</sup> but angioplasty should be considered in three situations. Firstly, it should be considered in diabetic patients, particularly non-insulin-dependent diabetics, preferentially those who have two-vessel disease and do not present another factor that favors restenosis (which, as mentioned, is the main factor responsible for the poor evolution of diabetic patients treated with angioplasty), such as arterial diameter of less than 3 mm, long lesions, or lesions in which residual stenosis is expected after angioplasty despite stent implantation. The second situation is surgery associated with an unacceptable incidence of complications (comorbidity, inadequate anatomy) or if arterial grafts cannot be used. At this point it should be remembered that, even with balloon angioplasty, the initial complication (mortality or infarction) are less with angioplasty than with surgery. Therefore, in patients with a high risk for surgery, acute angina or extensive ischemia, in which revascularization is inevitably indicated, it seems reasonable to use stent angioplasty although we know that the long-term results are going to be limited. The third situation is when the indication must be evaluated in an unstable context (primary angioplasty in the context of acute myocardial infarction or angioplasty in refractory unstable angina), in which treatment of the causal lesion is inevitable, regardless of whether or not other lesions exist. The later management of other lesions will be assessed individually at each center and for each patient. If we indicate angioplasty in a diabetic, it seems obligatory to perform angioplasty under treatment with glycoprotein IIb/IIIa inhibitors and to implement intensive secondary prevention measures during the follow-up.

## THE NEAR FUTURE OF PERCUTANEOUS REVASCULARIZATION IN DIABETICS

Although the present of percutaneous revascularization in diabetics is, in some respects, uncertain, the future is very promising. This review probably will have a new approach in the near future. This promising situation is due fundamentally to the development of intracoronary brachytherapy and, above all, of stents that release drugs that inhibit cell proliferation. Restenosis has been the main limitation of coronary angioplasty, including angioplasty with stent implantation. The basic mechanism of restenosis after stent implantation is intimal hyperplasia, which is especially important in diabetics. Inhibition of intimal hyperplasia would reduce or impede restenosis, resulting in better clinical results. Both intracoronary brachytherapy and stents coated with proliferation inhibitors have managed to effectively prevent restenosis. Its cli-

nical effectiveness in diabetics is in the process of evaluation, with results that sometimes cannot be described in any other way than as spectacular.

## Intracoronary brachytherapy

Intracoronary brachytherapy reduces the incidence of restenosis because of its capacity to inhibit cell proliferation due to the DNA damage it causes.<sup>88-91</sup> Its most important effect takes place during the phase of mitosis and G2, causing cell death or, more frequently, annulling the replication capacity and accelerating apoptosis. This effect is produced on smooth muscle cells, monocytes, and macrophages, determining an inhibition of neointimal proliferation. Clinical trials of intracoronary brachytherapy with gamma radiation (SCRPPS, WRIST, and GAMMA) or beta radiation (START, INHIBIT) have demonstrated that it reduces the incidence of restenosis by 61% to 69%, with very low restenosis rates, from 16% to 25%, which occur mainly within the context of intra-stent restenosis.<sup>88,90,91</sup> Recent studies seem to confirm that, in spite of the presence of very late reocclusion, brachytherapy is effective in the long term.<sup>92</sup> At present, brachytherapy can be considered the treatment of choice for intra-stent restenosis.

The mechanism by which brachytherapy reduces restenosis (reduction of intimal hyperplasia) could be potentially useful in diabetics treated with stent implantation. In fact, recent communications demonstrate that the effectiveness of brachytherapy in reducing restenosis in the context of intra-stent restenosis is especially high in diabetics.<sup>93</sup> The restenosis rate was 15.6% in diabetics and 11% in non-diabetics ( $P=0.3$ ) treated with brachytherapy, meaning that it was very similar. In contrast, the restenosis rate was 64% in diabetics and 48% in non-diabetics randomized to angioplasty without brachytherapy. These results have inspired the design and execution of studies in which brachytherapy is applied with angioplasty in *de novo* lesions in diabetics. If they demonstrate that they substantially improve the clinical evolution, the indications for angioplasty in groups like diabetics with extensive coronary disease may expand.

## Angioplasty with stents that release inhibitors of intimal hyperplasia

The publication in June 2002 of the results of the RAVEL study has probably opened a new chapter in percutaneous coronary revascularization that could be especially beneficial for diabetics.<sup>95,96</sup> The stent controlled two of the three mechanisms responsible for restenosis (the initial elastic recoil and unfavorable late remodeling). However, it did not reduce intimal hyperplasia and, in fact, increased it. The development of stents that release drugs that inhibit intimal hyper-

plasia may make it possible to eliminate or greatly reduce the main limitation of angioplasty: restenosis.<sup>95</sup> This is what was discovered in the RAVEL study.<sup>96</sup> The first experimental studies demonstrated that rapamycin (sirolimus) also impeded T-cell proliferation and the proliferation and migration of smooth muscle cells (a fundamental phenomenon in postangioplasty intimal hyperplasia). Based on these observations, a stent coated with rapamycin that was slowly released was developed. This stent was applied in 30 patients in São Paulo<sup>97</sup> and in 15 patients in Rotterdam.<sup>98</sup> None of the patients developed restenosis and only one patient died as a result of cerebral hemorrhage, one had subacute occlusion, and one had an infarction at 14 months. These spectacular results motivated the RAVEL study. In this study, 238 patients were assigned randomly to angioplasty with a standard stent or with a stent coated with sirolimus. The results impressed the scientific community. None of the patients in the group randomized to sirolimus-coated stents developed restenosis, vs 26% of those that received the standard stent. The rate of major cardiac events at 1 year was 5.8% in the coated stent group vs 29% ( $P>.001$ ) in the control group. In the 19 diabetic patients treated with sirolimus-coated stents, restenosis was not observed. Other studies (ELUTES, TAXUS) with stents coated with inhibitors of myointimal proliferation (taxol, paclitaxel) have demonstrated less favorable, but still encouraging, results. As a result, new questions must be asked: does this type of stent prevent or delay restenosis? Will it increase the incidence of late thrombotic occlusion? Is it safe in acute coronary syndrome? We should have answers to these questions in the near future.

It is very likely that the development of these devices will substantially change and improve the results of angioplasty, particularly in diabetics. This will make new studies necessary, especially in the context of multivessel disease, whose results will require a re-examination of revascularization strategies in diabetics, thus further expanding the use of percutaneous techniques.

## REFERENCES

1. Fein F, Scheur J. Heart diseases in diabetes mellitus: theory and practice. En: Rifkin H, Porte D Jr, editors. New York: Elsevier, 1990; p. 812-23.
2. Boden WE. Management of patients with diabetes and coronary artery disease. En: Sobel BE, Schneider DJ, editors. Medical management of diabetes and Heart disease. New York: Marcel Dekker Inc., 2002; p. 185-209.
3. Douglas JS. Special therapeutic considerations: coronary interventions and coronary surgery. En: Sobel BE, Schneider DJ, editors. Medical management of diabetes and Heart disease. New York: Marcel Dekker Inc., 2002; p. 231-57.
4. National Cholesterol Education Program. Executive summary of the third report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation and treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486-97.
5. Haffner SM, Lehto S, Rönnemaa T, Pyörälä K, Laakso M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;339:229-34.
6. Ellis SG, Narins CR. Problem of angioplasty in diabetic. *Circulation* 1997;96:1707-10.
7. O'Neill WW. Multivessel balloon angioplasty should be abandoned in diabetic patients. *J Am Coll Cardiol* 1998;31:20-2.
8. Haffner SM. Coronary heart disease in patients with diabetes. *N Engl J Med* 2000;342:1040-1.
9. Alonso JJ, Azpitarte J, Bardají A, Cabadés A, Fernández A, Palencia M, et al. Guías de práctica clínica de la Sociedad Española de Cardiología en cirugía coronaria. *Rev Esp Cardiol* 2000; 53:241-66.
10. Esplugas E, Alfonso F, Alonso JJ, Asín E, Elízaga J, Iñiguez A, et al. Guías de práctica clínica de la Sociedad Española de Cardiología en cardiología intervencionista: angioplastia coronaria y otras técnicas. *Rev Esp Cardiol* 2000;53:218-40.
11. Smith SC Jr, Dove JT, Jacobs AK, Kennedy JW, Kereiakes D, Kern MJ, et al. ACC/AHA Guidelines for percutaneous coronary intervention (revision of the 1993 PTCA guidelines: executive summary and recommendations). A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to revise the 1993 guidelines for percutaneous transluminal coronary angioplasty). *J Am Coll Cardiol* 2001;37:2215-39.
12. Eagle KA, Guyton RA, Davidoff R, Ewy GA, Fonger J, Gadner TJ, et al. ACC/AHA guidelines for coronary bypass graft surgery: executive summary and recommendations. A report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Subcommittee on Practice Guidelines). *Circulation* 1999;100:1464-80.
13. Lazaro P, Fitch K, Martín Y. Estándares para el uso apropiado de la angioplastia coronaria transluminal percutánea y cirugía aortocoronaria. *Rev Esp Cardiol* 1998;51:689-715.
14. Rihal CS, Gersh BJ, Yusuf S. Chronic coronary artery disease: coronary artery bypass surgery vs percutaneous transluminal coronary angioplasty vs medical therapy. En: Yusuf S, editor. Evidence based cardiology. London: BMJ Books 1998; p. 368-92.
15. Emanuelsson H, Brorsson H, Bernstein SJ, Werkö L. Coronary artery bypass graft surgery, percutaneous transluminal coronary angioplasty or medical therapy in anginal pain. A literature review for rating indications. Informe del The Swedish Council on Technology Assessment in Health Care, 1998.
16. Barzilay JI, Kronmal RA, Bittnar V, Eaker E, Evans C, Foster ED. Coronary artery disease and coronary artery bypass grafting in diabetics aged  $\geq 65$  years (report from the Coronary Artery Surgery Study [CASS] registry). *Am J Cardiol* 1994;74:334-9.
17. Herlitz J, Wognsen GB, Emanuelsson H, Haglid M, Karlsson BW, Karlsson T, et al. Mortality and morbidity in diabetic and nondiabetic patients during a 2-year period after coronary artery bypass grafting. *Diabetes Care* 1996;19:698-703.
18. FRISC Investigators. Invasive compared with non-invasive treatment in unstable coronary artery disease: FRISC II prospective randomised multicentre study. *Lancet* 1999;354:708-15.
19. Goy JJ. Contemporary approach to management of unstable angina. *Lancet* 1999;354:694-5.
20. Wallentin L, Lagerqvist B, Husted S, Kontny F, Stahle E, Swahn E, for the FRISC investigators. Outcome at 1 year after an invasive compared with a non-invasive strategy in unstable coronary artery disease: the FRISC II invasive randomised trial. *Lancet*

- 2000;356:9-16.
21. Cannon CP, Weintraub WS, Demopoulos LA, Vicari R, Frey MJ, Lakkis N, et al. Comparison of early invasive and conservative strategies in patients with unstable coronary syndromes treated with glycoprotein IIb/IIIa inhibitor tirofiban. *N Engl J Med* 2001;344:1879-87.
  22. Andersen HR. The Danis Multicenter Randomized Trial on Thrombolytic Therapy versus Acute Coronary Angioplasty in acute Myocardial Infarction (DANAMI II). Atlanta: 51<sup>st</sup> Annual Scientific Session of the American College of Cardiology, 17-20 marzo de 2002.
  23. Weaver WD, Simes J, Betriu A, Grines CL, Zijlstra F, García E, et al. Comparison of primary coronary angioplasty and intravenous thrombolytic therapy for acute myocardial infarction. A quantitative review. *JAMA* 1997;278:2093-8.
  24. Michels KB, Yusuf S. Does PTCA in acute myocardial infarction affect mortality and reinfarction rates? A quantitative overview (meta-analysis) of the randomized trials. *Circulation* 1995;91:476-85.
  25. Stone GW, Grines CL, Cox DA, García E, Tchong JE, Griffin JJ, et al. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 2002;346:954-5.
  26. Satio S, Hosokawa G, Tanaka S, Nakamura S. Primary stent implantation is superior to balloon angioplasty in acute myocardial infarction: final results of the primary angioplasty versus stent implantation in acute myocardial infarction (PASTA) trial. *PASTA Trial Investigators. Cathet Cardiovasc Interv* 1999;48:262-8.
  27. Grines CL, Cox DA, Stone GW, García E, Mattos LA, Giambartolomei A, et al. Coronary angioplasty with or without stent implantation for acute myocardial infarction. *Stent Primary Angioplasty in Myocardial Infarction Study Group. N Engl J Med* 1999;341:1949-56.
  28. Hasdai A, Granger CB, Srivatsa SS, Criger DA, Ellis SG, Califf RM, et al. Diabetes mellitus and outcome after primary coronary angioplasty for acute myocardial infarction: lessons from the GUSTO-IIb angioplasty substudy. *J Am Coll Cardiol* 2000;35:1502-12.
  29. Mattos LA, Grines CL, de Sousa E, Sousa AGMR, Stone GW, Cox D, et al. One-year follow-up after primary coronary interventions for acute myocardial infarction in diabetics patients. A substudy of the STENT PAMI trial. *Arq Bras Cardiol* 2001;77:556-61.
  30. Silva JA, Ramee SR, White CJ, Collins TJ, Jenkins JS, Nunez E, et al. Primary stenting in acute myocardial infarction: influence of diabetes mellitus in angiographic results and clinical outcome. *Am Heart J* 1999;138:446-55.
  31. Silva JA, Nunez E, White CJ, Collins TJ, Jenkins JS, Zhang S, et al. Predictors of stent thrombosis after primary stenting for acute myocardial infarction. *Catheter Cardiovasc Interv* 1999;47:415-22.
  32. Bhatt DL, Marso SP, Lincoff AM, Wolski KE, Ellis SG, Topol EJ. Abciximab reduces mortality in diabetics following percutaneous coronary intervention. *J Am Coll Cardiol* 2000;35:922-8.
  33. Roffi M, Chew DP, Mukherjee D, Bhatt DL, White JA, Heesch C, et al. Platelet glycoprotein IIb/IIIa inhibitors reduce mortality in diabetic patients with non-ST-segment-elevation acute coronary syndromes. *Circulation* 2001;104:2767-71.
  34. Masotti M, Betriu A. Optimal reperfusion in evolving myocardial infarction. Does abciximab improve outcomes in stent treated patients. *Eur Heart J* 2001;22(Suppl):475.
  35. The ESPRIT Investigators. Novel dosing regimen of eptifibatid in planned coronary stent implantation (ESPRIT): a randomised, placebo-controlled trial. *Lancet* 2000;356:2037-44.
  36. O'Shea JC, Hafley GE, Greenberg S, Hasselblad V, Lorenz TJ, Kitt MM, et al. Platelet glycoprotein IIb/IIIa integrin blockade with eptifibatid in coronary stent intervention. The ESPRIT trial: a randomized controlled trial. *JAMA* 2001;285:2468-73.
  37. O'Shea JC, Buller CE, Cantor WJ, Chandler AB, Cohen EA, Cohen DJ. Long-term efficacy of platelet glycoprotein IIb/IIIa integrin blockade with eptifibatid in coronary stent intervention. *JAMA* 2002;287:618-21.
  38. Kip KE, Faxon DP, Detre KM, Yeh W, Kelsey SF, Currier JW, for the investigators of the NHLBI PTCA Registry. Coronary angioplasty in diabetic patients: the National Heart, Lung and Blood Institute Percutaneous Transluminal Coronary Angioplasty Registry. *Circulation* 1996;94:1818-25.
  39. Stein B, Weintraub WS, Gebhart SSP, Cohen-Bernstein CI, Grosswald R, Liberman HA, et al. Influence of diabetes mellitus on early and late outcome after percutaneous transluminal coronary angioplasty. *Circulation* 1995;91:979-89.
  40. Kurbaan A, Bawker TJ, Ilesley CD, Sigwart U, Rickards AF, on behalf of the CABRI investigators. Difference in the mortality of the CABRI diabetic and nondiabetic populations and its relation to coronary artery disease and the revascularization mode. *Am J Cardiol* 2001;87:947-50.
  41. King SB III, Kosinki AS, Guyton RA, Lembo NJ, Weintraub WS, for the Emory Angioplasty versus Surgery Trial. Eight-year mortality in the Emory Angioplasty versus Surgery Trial (EAST). *J Am Coll Cardiol* 2003;41:1116-21.
  42. The BARI investigators. Seven-year outcome in the bypass angioplasty revascularization investigation (BARI) by treatment and diabetic status. *J Am Coll Cardiol* 2000;35:1122-9.
  43. Barsness GW, Peterson ED, Ohman EM, Nelson CL, DeLong ER, Reves JG, et al. Relationship between diabetes mellitus and long-term survival after coronary bypass and angioplasty. *Circulation* 1997;96:2551-6.
  44. Rozeman Y, Sapoznikov D, Gotsman MS. Restenosis and progression of coronary disease after balloon angioplasty in patients with diabetes mellitus. *Clin Cardiol* 2000;23:890-4.
  45. Barsness GW, Hardison RM, Detre MK, Alderman EL, Bourassa MG, Bashore TM, et al. Lesion progression and reestenosis in PTCA-treated diabetic and non-diabetic patients with multivessel disease. *Circulation* 2000;102(Suppl II):479.
  46. Kurbaan AS, Bowker TJ, Ilesley CD, Sigwart U, Rickards AF. Is the poorer outcome post revascularization in diabetics related to completeness of revascularization? *Circulation* 2000;102(Suppl II):256.
  47. Serruys PW, de Jaegere P, Kiemeneij F, Macaya C, Rutsch W, Heyndrickx G, et al. A comparison of balloon-expandable-stent implantation with balloon angioplasty in patients with coronary artery disease: Benestent Study Group. *N Engl J Med* 1994;331:489-95.
  48. Fischman DL, Leon MB, Baim DS, Schatz RA, Savage MP, Penn I, et al. A randomized comparison of coronary-stent placement and balloon angioplasty in the treatment of coronary artery disease: Stent Restenosis Study Investigators. *N Engl J Med* 1994;331:496-501.
  49. Serruys PW, Van Hout B, Bonnier H, Legrand V, García E, Macaya C, et al. Randomised comparison of implantation of heparin-coated stents with balloon angioplasty in selected patients with coronary artery disease (Benestent II). *Lancet* 1998;352:673-81.
  50. Macaya C, Serruys PW, Ruygrok P, Suryapranata H, Mast G, Klugmann S, et al. Continued benefit of coronary stenting versus balloon angioplasty: one-year clinical follow-up of Benestent trial: Benestent Study Group. *J Am Coll Cardiol* 1996;27:255-61.
  51. Betriu A, Masotti M, Serra A, Alonso J, Fernández-Avilés F, Gimeno F, et al. Randomized comparison of coronary stent implantation and balloon angioplasty in the treatment of de novo coronary artery lesions (START). A four year follow-up. *J Am Coll Cardiol* 1999;34:1498-506.
  52. Versaci F, Gasparidone A, Tomai F, Crea F, Chiariello L, Giffre PA. A comparison of coronary artery stenting with angioplasty

- for isolated stenosis of the proximal left anterior descending coronary artery. *N Engl J Med* 1997;336:817-22.
53. Sirnes PA, Golf S, Myreng Y, Molstad P, Albertsson P, Mangschau A, et al. Sustained benefit of stenting chronic coronary artery occlusion: long-term clinical follow-up of the Stenting in Chronic Coronary Occlusion (SICCO) study. *J Am Coll Cardiol* 1998;32:305-10.
  54. Rubartelli P, Niccoli L, Verna E, Giachero C, Zimarino M, Fontanelli A, et al. Stent implantation versus balloon angioplasty in chronic coronary occlusions: results from the GISSOC trial. Gruppo Italiano di Studio sullo Stent nelle Occlusioni Coronariche. *J Am Coll Cardiol* 1998;32:90-6.
  55. Hancock J, Thomas MR, Holmberg S, Wainwright RJ, Jewitt DE. Randomised trial of elective stenting after successful percutaneous transluminal coronary angioplasty of occluded coronary arteries. *Heart* 1998;79:18-23.
  56. Erbel R, Haude M, Hopp HW, Franzen D, Rupprecht HJ, Heublein B, et al. Coronary-artery stenting compared with balloon angioplasty for restenosis after initial balloon angioplasty. *N Engl J Med* 1998;339:1672-8.
  57. Savage MP, Douglas JS Jr, Fischman DL, Pepine CJ, King SB 3rd, Werner JA, et al. Stent placement compared with balloon angioplasty for obstructed coronary bypass grafts. Saphenous Vein De Novo Trial Investigators. *N Engl J Med* 1997;337:740-7.
  58. Hernández JM, Goicolea J, Durán JM, Augé JM. Registro de la actividad de la Sección de Hemodinámica y Cardiología Intervencionista de la Sociedad Española de Cardiología del año 2000. *Rev Esp Cardiol* 2001;54:1426-38.
  59. Savage M, Fischman D, Slota P, Rake P, Leon M, Schatz R, et al. Coronary intervention in the diabetic patient: improved outcome following intracoronary stent placement [resumen]. *J Am Coll Cardiol* 1997;29(Suppl A):188 A.
  60. Van Belle E, Bauters C, Hubert E, Bodart JC, Abolmaali K, Meurice T, et al. Restenosis rate in diabetic patients. A comparison of coronary stenting and balloon angioplasty in native coronary vessels. *Circulation* 1997;96:1454-60.
  61. Navarro LF, Íñiguez A, Ibargollín R, García R, Fernández I, Rodríguez RC. Comparación de la evolución clínica y angiográfica de pacientes diabéticos y no diabéticos tratados con angioplastia convencional frente a implantación de stent en arterias coronarias. *Rev Esp Cardiol* 2000;53:172-8.
  62. Bauters C, Hubert E, Prat A, Bougrimi K, Van Belle E, McFadden EP, et al. Predictors of restenosis after coronary stent implantation. *J Am Coll Cardiol* 1998;31:1291-8.
  63. Elezi S, Kastrati A, Pache A, Wehinger A, Hadamitzky M, Dirschinger J, et al. Diabetes mellitus and the clinical and angiographic outcome after coronary stent placement. *J Am Coll Cardiol* 1998;32:1866-73.
  64. Kastrati A, Schömig A, Elezi S, Schühlen H, Dirschinger J, Hadamitzky M, et al. Predictive factors of restenosis after coronary stent placement. *J Am Coll Cardiol* 1997;30:1428-36.
  65. Alonso JJ, Durán JM, Gimeno F, Bermejo J, Garcimartín I, Fuente L, et al. Clinical and angiographic restenosis after coronary stenting. Incidence and predictors. *J Am Coll Cardiol* 1997;27(Suppl A):239A.
  66. Schofer J, Schlüter M, Rau T, Hammer F, Haag N, Mathey DG. Influence of treatment modality on angiographic outcome after coronary stenting in diabetic patients: a controlled study. *J Am Coll Cardiol* 2000;35:1554-9.
  67. Alonso JJ, Durán JM, Gimeno F, Ramos B, Gómez I, de la Fuente L, et al. Coronary stenting in patients with diabetes mellitus: Long-term outcome and predictors. *Eur Heart J* 2000;21(Suppl):171.
  68. Pascual DA, Valdés M, García F, Garzón A, González J, García A, et al. Influencia de la diabetes mellitus en los resultados clínicos tardíos de la revascularización coronaria con stents. *Rev Esp Cardiol* 2001;54:261-8.
  69. Abizaid A, Kornowski R, Mintz GS, Hong MK, Abizaid AS, Mehran R, et al. The influence of diabetes mellitus on acute and late clinical outcomes following coronary stent implantation. *J Am Coll Cardiol* 1998;32:584-9.
  70. Alonso JJ. Diabetes mellitus y revascularización coronaria. La controversia continua. *Rev Esp Cardiol* 2001;54:255-8.
  71. Mudra H, Regar E, Klauss V, Werner F, Hennke KH, Sbarouni E. Serial follow-up after optimized ultrasound-guide deployment of Palmaz-Schatz Stents. In-Stent neointimal proliferation without significant reference segment response. *Circulation* 1997;95:363-70.
  72. Kornowski R, Mintz GS, Kent KM, Pichard AD, Satler LF, Bucher TA. Increased restenosis in diabetes mellitus after coronary interventions is due to exaggerated intimal hyperplasia. A serial intravascular ultrasound study. *Circulation* 1997;95:1366-9.
  73. Takagi T, Yoshida K, Akasaka T, Kaji S, Kawamoto T, Honda Y, et al. Hyperinsulinemia during oral glucose tolerance test is associated with increased neointimal tissue proliferation after coronary stent implantation in nondiabetic patients. *J Am Coll Cardiol* 2000;36:731-8.
  74. Goy JJ, Eeckhout E, Burnand B, Vogt P, Stauffer JC, Humi M, et al. Coronary angioplasty versus left internal mammary artery grafting for isolated proximal left anterior descending artery stenosis. *Lancet* 1994;343:1449-53.
  75. Hueb WA, Bellotti G, de Oliveira SA, Arie S, de Albuquerque CP, Jatene AD, et al. The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses. *J Am Coll Cardiol* 1995;26:1600-5.
  76. Hueb W, Cardoso RH, Soares PR, Albuquerque CP, Cezar LA, Bellotti G et al. The Medicine, Angioplasty and Surgery Study (MASS) a Prospective Randomised Trial of Medical Therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenosis, five years of follow-up. *Circulation* 1998;98(Suppl 1):I-216.
  77. RITA trial participants. Coronary angioplasty versus coronary artery bypass surgery: the Randomised Intervention Treatment of Angina (RITA) trial. *Lancet* 1993;341:573-80.
  78. The Bypass Angioplasty Revascularization Investigation Investigators. Comparison of coronary bypass surgery with angioplasty in patients with multivessel disease. *N Engl J Med* 1996;335:217-25.
  79. CABRI trial participants. First-year results of CABRI (Coronary Angioplasty versus Bypass Revascularization Investigation). *Lancet* 1995;346:1179-84.
  80. King SB 3rd, Lembo NJ, Weintraub WS, Kosinski AS, Barnhart HX, Kutner MH, et al. A randomized trial comparing coronary angioplasty versus Surgery Trial (EAST). *N Engl J Med* 1994;331:1044-50.
  81. Rodríguez A, Mele E, Peyregne E, Bullon F, Pérez-Balino N, Liprandi MI, et al. Three-year follow-up of the Argentine Randomized Trial of Percutaneous Transluminal Coronary Angioplasty Versus Coronary Artery Bypass Surgery in Multivessel Disease (ERACI). *J Am Coll Cardiol* 1996;27:1178-84.
  82. Hamm CW, Reimers J, Ischinger T, Rupprecht HJ, Berger J, Bleifeld W. A randomized study of coronary angioplasty compared with bypass surgery in patients with symptomatic multivessel coronary disease: German Angioplasty Bypass Surgery Investigation (GABI). *N Engl J Med* 1994;331:1037-43.
  83. The BARI investigators. Influence of diabetes on 5-year mortality and morbidity in a randomized trial comparing CABG and PTCA in patients with multivessel disease. The Bypass Angioplasty Revascularization Investigation (BARI). *Circulation* 1997;96:1761-9.
  84. Weintraub WS, Stein B, Kosinski A, Douglas JS, Ghazzal ZMB, Jones EL. Outcome of coronary bypass surgery versus coronary angioplasty in diabetic patients with multivessel coronary artery

- disease. *J Am Coll Cardiol* 1998;31:10-9.
85. Niles NW, McGrath PD, Malenka D, Quinton H, Wennberg D, Shubrooks SJ, et al. Survival of patients with diabetes and multivessel coronary artery disease after surgical or percutaneous coronary revascularization: results of a large regional prospective study. *J Am Coll Cardiol* 2001;37:1008-15.
  86. Detre KM, Guo P, Holubkov R, Califf RM, Sopko G, Bach R, et al. Coronary revascularization in diabetic patients. A comparison of the randomized and observational components of the Bypass Angioplasty Revascularization Investigation (BARI). *Circulation* 1999;99:633-40.
  87. Abizaid A, Costa MA, Centemero M, Abizaid AS, Legrand VMG, Limet RV. Clinical and economic impact of diabetes mellitus on percutaneous and surgical treatment of multivessel coronary disease patients. Insights from the Arterial Revascularization Therapy Study (ARTS) trial. *Circulation* 2001;104:533-8.
  88. Bailey SR. Coronary restenosis: a review of current insights and therapies. *Cathet Cardiovasc Interv* 2002;55:265-71.
  89. Waksman R, Rodríguez JC, Robinson KA, Cipolla GD, Crocker IR, Scott NA, et al. Effect of intravascular irradiation on cell proliferation, apoptosis, and vascular remodeling after balloon overstretch injury of porcine coronary arteries. *Circulation* 1997;96:1944-52.
  90. Ajani AE, Kim HS, Waksman R. Clinical trial of vascular brachytherapy for in-stent restenosis: update. *Cardiovasc Rad Med* 2001;2:107-13.
  91. Raizner AE, Oesterle SN, Waksman R, Serruys PW, Colombo A, Lim YL, et al. Inhibition of restenosis with beta-emitting radiotherapy: report of the Proliferation Reduction with Vascular Energy Trial (PREVENT). *Circulation* 2000;102:951-8.
  92. Grise MA, Massullo V, Jani S, Popma JJ, Russo RJ, Schatz RA, et al. Five-year clinical follow-up after intracoronary radiation. Result of a randomized clinical trial. *Circulation* 2002;105:2737-40.
  93. Williams DO. Intracoronary brachytherapy. Past, present and future. *Circulation* 2001;105:2699.
  94. Gruber L, Waksman R, Ajani AE, Kim H-S, White RL, Chan RC, et al. The effect of intracoronary radiation for recurrent in-stent restenosis in diabetic patients. *Eur Heart J* 2001;22(Suppl):5.
  95. Regar E, Serruys PW. El estudio RAVEL. Reestenosis del cerebro por ciento: un sueño del cardiólogo hecho realidad. *Rev Esp Cardiol* 2001;55:459-62.
  96. Morice MC, Serruys PW, Sousa JE, Fajadet J, Ban Hayasi E, Perin M, et al. A randomized comparison of a sirolimus-eluting stent with a standard stent for coronary revascularization. *N Engl J Med* 2001;346:1773-80.
  97. Sousa JEMR, Costa MA, Abizaid A, Abizaid AS, Feres F, Pinto IMF, et al. Mid- (4 months) and long-term (1 year) QCA and three-dimensional IVUS follow-up after implantation of sirolimus-coated Stent in human coronary arteries. *J Am Coll Cardiol* 2001;37(Supl):8.
  98. Rensing BJ, Vos J, Smits PC, Floey DP, Van den Brand MJ, Van der Giessen WJ, et al. Coronary restenosis elimination with a sirolimus eluting Stent. First European human experience with six month angiographic and intravascular ultrasonic follow-up. *Eur Heart J* 2001;22:2125-30.