

Sports Requiring Stressful Physical Exertion Cause Abnormalities in Plasma Lipid Profile

Jonatan R. Ruiz, José L.M. Mesa, Ignacio Mingorance, Antonio Rodríguez-Cuartero, and Manuel J. Castillo

Departamento de Fisiología Médica, Facultad de Medicina, Universidad de Granada, Granada, Spain.

Objectives. To study the effect of different types of sports on plasma lipid profile.

Subjects and method. Fasting plasma levels of total, LDL- and HDL-cholesterol (TC, LDL-C, HDL-C), triglycerides, phospholipids, apo A-I, apo B-100, and Lp(a) were measured in 28 swimmers, 17 volleyball players and 23 soccer players, and in a control group. All subjects participated in official national competitions except the control group. The results were compared to those in a control group of 26 healthy sedentary subjects. All groups were matched according to age, body mass index and nutritional status. Cutoff points to classify subjects with high plasma concentrations of Lp(a) and LDL-C were 32 mg/dL and 128 mg/dL, respectively. For other variables the cutoff points were the third quartile values recorded in the control group, except for apo A-I and HDL-C, for which we used the first quartile values.

Results. Persons who practice sports involving a high level of physical exertion (volleyball and soccer players) had a less favorable lipid profile compared to control subjects. In contrast, swimmers had a more favorable lipid profile. The odds ratio for lipid profile anomalies was significantly higher in volleyball and soccer players, and lower in swimmers. In the former two groups we observed a clear tendency toward simultaneous elevations in LDL-C, apo B-100 and Lp(a). In contrast, the opposite interaction was seen in swimmers, who had lower values for all lipids.

Conclusion. Our results show that stressful physical exertion can lead to abnormalities in plasma lipid profile.

Full English text available at: www.revvespcardiol.org

SEE EDITORIAL ON PAGES 495-8

Correspondence: Dr. J. Ruiz Ruiz.
Departamento de Fisiología. Facultad de Medicina.
Universidad de Granada.
Avda. de Madrid, s/n. 18012 Granada. España.
E-mail: ruizj@ugr.es

Received 17 September, 2003.
Accepted for publication 1 April, 2004.

Deportes con alto grado de estrés físico afectan negativamente al perfil lipídico plasmático

Objetivos. Estudiar el efecto en el perfil lipídico plasmático de distintos tipos de deporte.

Sujetos y método. Se midieron los valores plasmáticos basales de colesterol total (CT), colesterol unido a lipoproteínas de baja densidad (cLDL), colesterol unido a lipoproteínas de alta densidad (cHDL), triglicéridos, fosfolípidos, apolipoproteínas (apo) A-I y B100, y lipoproteína (a) (Lp[a]), en 3 grupos de deportistas enrolados en competiciones nacionales (28 nadadores, 17 jugadores de voleibol y 23 futbolistas) así como en un grupo control (26 sujetos sedentarios sanos). Los puntos de corte utilizados para clasificar a los sujetos con cLDL o Lp(a) elevados fueron 128 y 32 mg/dl, respectivamente. Para el resto de las variables, los puntos de corte se establecieron en el tercer cuartil del grupo control, excepto para apo A-1 y cHDL, que se establecieron en el primer cuartil.

Resultados. Los deportistas que practicaban deportes con un alto grado de estrés físico (futbolistas y jugadores de voleibol) presentaron un perfil lipídico menos favorable que los sujetos control. Por el contrario, los nadadores presentaban un perfil lipídico más favorable. La tasa de probabilidad (*odds ratio*) de tener anomalías del perfil lipídico era significativamente mayor para los jugadores de voleibol y los futbolistas, mientras que era inferior para los nadadores. En los primeros 2 grupos se observó una clara tendencia a presentar simultáneamente valores elevados de cLDL, apo B100 y Lp(a). Por el contrario, en los nadadores la interacción se producía en sentido inverso: valores bajos de los distintos parámetros.

Conclusión. Según los resultados de este estudio, el ejercicio físico estresante influye negativamente en el perfil lipídico plasmático.

Palabras clave: Ejercicio. Estrés. Lípidos. Lipoproteínas.

INTRODUCTION

The regular practice of moderate, aerobic exercise is associated with a healthy plasma lipid profile¹ and a reduced risk of coronary artery disease and cardiovascular-related death.^{2,3} The results of cross-sectional^{4,5} and longitudinal⁶ studies suggest that aerobic exercise increases plasma levels of high density lipoprotein cholesterol (HDL-C) and reduces triglycerides and

ABBREVIATIONS

TC: total cholesterol.
 LDL-C: low density lipoprotein cholesterol.
 HDL-C: high density lipoprotein cholesterol.
 apo A-I: apolipoprotein A-I.
 apo B100: apolipoprotein B100.
 Lp(a): lipoprotein (a).

low density lipoprotein cholesterol (LDL-C) levels. However, contradictory results have been reported with respect to certain types of exercise demanding strenuous physical exertion,⁷⁻⁹ especially in terms of lipoprotein (a) [Lp(a)] levels.¹⁰ Several studies report no differences in plasma Lp(a) levels between physically trained and sedentary individuals,¹¹⁻¹³ whereas others report higher levels in trained subjects.^{1,14} Contradictory results have also been obtained in experimental studies.^{12,13,15} There are several possible explanations for these discrepancies, including differences in the methods used to determine the levels of lipids and lipoproteins, the difference in physical fitness of the individuals tested, ethnic factors, or differences in training history or the type, quantity and intensity of exercise undertaken, etc. On their own, increases in Lp(a) as a consequence of high intensity or stressful exercise are given little clinical importance. However, when both Lp(a) and LDL-C levels are high, the risk of cardiovascular disease increases exponentially.¹⁰ Lp(a) has been identified as an atherogenic and thrombogenic factor¹⁶ that can act synergistically with other cardiovascular risk factors.¹⁷⁻¹⁹ In fact, the study of the interactions between apolipoprotein B100 (apo B100),

LDL-C and Lp(a) is of interest in the characterization of cardiovascular risk.^{11,19}

The aims of the present work were: *a)* to compare the plasma lipid profile of persons who practice different types of sport involving different levels of physical stress, and *b)* to characterize the possible interactions between the most important variables of the plasma lipid profile, especially those involving Lp(a).

SUBJECTS AND METHODS

Three groups of sportsmen—swimmers (n=29), volleyball players (n=17), and soccer players (n=23)—were studied, all of whom competed at the regional/national level. For comparison, a fourth, control group was established, composed of healthy, sedentary individuals (n=26) comparable to the sportsmen in terms of age, sex and anthropometric and sociological characteristics. Table 1 summarizes the characteristics of the study subjects. All subjects gave their consent to be included in the study after being informed of its characteristics and scientific interest, and all completed a questionnaire on their health and food habits. All subjects were healthy, took no medication or herbal products that might influence the lipid profile results, and were free of intercurrent disease. All consumed a typical, Spanish Mediterranean diet;²⁰ no significant qualitative differences were recorded between the groups. The subjects were informed of the importance of not embarking on any excessive dietary practices and of not imbibing excessive amounts of alcohol (2 drinks/day maximum, ≤ 30 g alcohol/day) in the 2 weeks before the taking of blood samples. They were also told not to undertake any stressful physical exertion in the preceding 48 hours.

Blood samples were taken after a 12 hour nocturnal fast. Standard biochemical and hematological profiles were determined as well as plasma lipid profiles. A

TABLE 1. Characteristics of the Study Subjects

	Controls (n=26)	Swimmers (n=28)	Volleyball Players (n=17)	Soccer Players (n=23)
Age, years	23.0 \pm 3.1	22.2 \pm 2.4	25.7 \pm 5.4	24.6 \pm 3.4
Smokers, %				
Never smoked	95	97	96	98
Past smokers	5	3	4	2
Current smokers	0	0	0	0
Body mass index	23.1 \pm 2.4	22.3 \pm 2.3	23.9 \pm 1.8	23.7 \pm 1.2
Blood pressure, mm Hg				
Diastolic	88 \pm 4	82 \pm 3	86 \pm 5	83 \pm 6
Systolic	126 \pm 3	122 \pm 4	123 \pm 3	121 \pm 4
Intake of macronutrients, %				
Carbohydrates	61.9 \pm 3.2	61.5 \pm 3.5	62.5 \pm 3.5	61.5 \pm 3.5
Proteins	16.0 \pm 3.6	16.2 \pm 2.5	17.5 \pm 2.2	6.4 \pm 3.1
Lipids	22.1 \pm 3.7	22.3 \pm 3.4	20.0 \pm 3.1	22.1 \pm 3.6

Values are means \pm standard deviation.

number of blood samples were taken at different times during the year; no seasonal differences were seen in the results.

Training History of the Study Subjects

All subjects completed a questionnaire on their physical activity over the last 5 years. Their trainers provided information on the characteristics of the training undertaken (quantity, intensity, number of sessions per day and week) and the frequency of competition.

The 28 swimmers selected had been registered for 10 ± 4.4 years and took part in regional and national competitions. The total time spent in training and competition was 13.4 ± 2.1 h per week. Training was divided into 90% aerobic resistance and 10% elasticity and joint movement. At the beginning of the season, muscular strength was developed using rubber tubing. Table 2 shows the exercise characteristics of the sport.

The 17 volleyball players selected had been registered for 9.4 ± 1.5 years and played in the national premier division. At the time of the study they trained once per day for 5 days per week, and played a match on the weekend. The total time spent in training and competition was 15.3 ± 2.2 h per week. Training included ~55% intermittent high intensity training with prolonged recovery times between series; the remaining time was given over to the development of muscular strength through specific explosive force training (mainly for the legs). Table 2 shows the exercise characteristics of the sport.

The 23 soccer players selected competed in the second division of the national professional league and had practiced this sport for the last 14.5 ± 3.1 years. At the time of the study they trained once per day, 4 days per week, but on Wednesdays they had a double training session. They also played 1 or 2 matches per week (weekends and during the week). The time spent training or in matches was 19.3 ± 2.5 h per week. Training included ~60% intermittent high intensity

training with short recovery times between series; the remaining time was given over to the development of muscular strength through specific explosive force training. Physical contact and falls, etc were common during training and matches—a natural feature of the game. Table 2 shows the exercise characteristics of the sport.

The training and competition characteristics recorded (Table 2) show the swimmers to be subject to a less stressful type of exercise. The training and competition characteristics endured by the soccer players, plus the emotional and environmental factors surrounding this sport,²¹ show these sportsmen undergo more stressful exercise regimens. Training characteristics were considered to be an ordinal variable.

Study of Plasma Lipid Profiles

Plasma total cholesterol (TC), LDL-C, and HDL-C levels were determined by an enzymatic colorimetric method using a Hitachi 911 autoanalyzer (Roche Diagnostics; Indianapolis, USA). Apolipoprotein A-I, apo B100, and Lp(a) levels were determined by nephelometric kinetics (Array 306, Beckman Coulter® Inc.; Fullerton, USA). Since Lp(a) values can change depending on the length of time serum samples are refrigerated,¹² samples were left for no longer than 4 hours at ambient temperature and were refrigerated (2–8 °C) for no longer than 4 days.

Assessment of Lp(a) Interactions

Subjects were grouped into 4 categories depending on their plasma Lp(a) and LDL-C levels (cut-off values 32 mg/dL and 128 mg/dL respectively). They were also grouped into 4 other categories depending on their plasma Lp(a) (cut-off value 32 mg/dL) and TC/HDL-C levels (the cut-off value corresponded to the third quartile value of the control subjects). Two interactions were therefore analyzed: that between Lp(a) and LDL-C, and that between Lp(a) and

TABLE 2. Exercise Characteristics of the Different Sports*

	Swimming	Volleyball	Soccer ^a
Exercise dynamics	Continuous	High intensity, intermittent	High intensity intermittent
Aerobic component	95%	60%	70%
Anaerobic component	5%	40%	30%
Recovery times between series, training	Long	Short	Short
Type of muscular contraction	Concentric/isoconcentric	Concentric/eccentric	Concentric/eccentric
Statistical (S) and dynamic (D) components	S moderate; D high	S low; D moderate	S low; D high
Intensity, VO_{2max} (%)	~60-80%	~50-80%	~75-80%
Joint impact	Low	High	Very high
Muscular impact	Medium	High	Very high
Hours of training-competition	13.4 ± 2.1	15.3 ± 2.2	19.3 ± 2.5

* VO_{2max} indicates maximum oxygen consumption. ^aFrequent falls and collisions, etc.

TC/HDL-C. Finally, an additional category was established to include subjects with high levels of Lp(a), LDL-C, and TC/HDL-C.

Assessment of Apo B100 Interactions

Subjects were grouped into 4 categories depending on their plasma apo B100 (the cut-off value corresponded to the third quartile value of the control subjects) and Lp(a) levels (cut-off value 32 mg/dL). They were also grouped into 4 other categories depending on their plasma apo B100 (cut-off value=third quartile value of the control subjects) and plasma LDL-C levels (cut-off 128 mg/dL). Two interactions were therefore analyzed: that between apo B100 and Lp(a), and that between apo B100 and LDL-C. An additional category was established to include subjects with high apo B100, Lp(a), and LDL-C levels.

Statistical Analysis

Normality was tested using the Shapiro-Wilks test. The Kruskal-Wallis test was used to compare non-normally distributed variables. One way analysis of variance (ANOVA) was used to compare normally distributed variables. To compare the homogeneity of the variance, the Levenne test was used. When variances were homogeneous, comparisons were made using the *post hoc* Tukey test; the *post hoc* Games-Howell test was used for non-homogenous variances. Descriptive values for normally distributed variables are represented as means±standard deviation (SD); non-normally distributed variables with an asymmetry of <-0.5 or >0.5 are represented as geometric means±SD. The odds ratio (OR) for each sport group was calculated using the Mantel-Haenszel method, taking the swimmers, soccer players, and volleyball players to be exposed groups, and the sedentary controls to be non-exposed. Cardiovascular risk factors were considered high when indicators were above the established cut-off points: Lp(a), 32 mg/dL and LDL-C, 128 mg/dL.¹⁷⁻¹⁹ For the remaining variables, the cut-off points were established as the third quartile of the sedentary control values, except for apo A-I and HDL-C, the cut-off values for which were fixed at the first quartile. The test for trends was used to determine the relationship between the type of physical exercise and cardiovascular risk factors. Significance was set at $\alpha=0.05$ in all cases, except for the Shapiro-Wilks test, in which it was set at $\alpha=0.20$. Ninety five percent confidence limits were determined.

RESULTS

Table 1 shows the characteristics of the participating subjects. Owing to the selection criteria, the groups showed no significant differences except with respect

to their exercise regimens. Table 2 shows that the degree of muscular overload or physical stress was greatest for the soccer players and least for the swimmers.

Tables 3 and 4 respectively show the lipid profiles and atherogenic indices of the 3 groups of sportsmen and the control group. Though the lipid profiles of the different groups were within the reference range for the subjects' age, and sex, they show significant differences. The plasma lipid and lipoprotein levels (Table 3) and the atherogenic indices (Table 4) of the swimmers were significantly healthier than those of the soccer and volleyball players. The soccer players had higher plasma levels of Lp(a) than any other group, including the control group (Table 3). The swimmers had the highest apo A-I values; their apo B100 and LDL-C levels were, however, lower than those of the soccer or volleyball players (Table 3). The atherogenic indices showed similar patterns.

The Figure shows the probability—the OR—of having an abnormal lipid profile (always compared to the control group). This analysis clearly shows the swimmers to have healthier lipid profiles, whereas soccer players and volleyball players were at a clear disadvantage, i.e., they had a particularly high probability of their plasma lipid profile being negatively affected. Therefore, the greater the physical stress endured, the greater the risk of having a lipid profile defined as a cardiovascular risk.

Table 5 shows the interactions of Lp(a) and apo B100 with other lipid cardiovascular risk factors. The swimmers showed no tendency towards having high Lp(a) or apo B100 levels associated with high LDL-C or TC/HDL-C levels. In contrast, the volleyball players showed a clear tendency to have high Lp(a) and apo B100 levels associated with high LDL-C and TC/HDL-C levels—a tendency even stronger among the soccer players.

In addition, both the volleyball and the soccer players had a significantly lower OR for having simultaneously low plasma Lp(a) and LDL-C, Lp(a), and TC/HDL-C, or Lp(a) and apo B100 levels. Therefore, the test for trends shows that the OR for having abnormalities with respect to the interactions of lipid profile components is directly related to the increase in stressful physical exertion (Table 5 and Figure 1).

DISCUSSION

The sportsmen subject to the highest levels of physical stress were the soccer players; these showed negative alterations in their atherogenic indices. The interactions of Lp(a) and apo B100 with other cardiovascular risk factors confirmed the possible impairment of the lipid profile among these sportsmen. In addition, these abnormalities were directly and significantly associated with the degree of physical stress to which the different groups of sportsmen were subject.

TABLE 3. Lipid Profile of Sportsmen and Control Groups*

Lipid profile	Controls (A) (n=26)	Swimmers (B) (n=28)	Volleyball Players (C) (n=17)	Soccer Players (D) (n=23)	P
TC	172.8 (159.8-185.9)	163.9 (155.6-172.2)	194.2 (171.0-217.4)	182.1 (168.3-196.0)	B-C: <.05 ^a
LDL-C	98.8 (85.8-111.8)	90.8 (84.4-97.1)	126.2 (107.0-145.8)	113.6 (100.1-127.2)	B-C: <.01b B-D: <.05 ^b
HDL-C	57.3 (52.9-61.6)	61.6 (57.4-65.8)	50 (46.1-53.9)	53.3 (49.9-56.7)	B-C: <.001 ^c B-D: <.05 ^c
Triglycerides	73.0* (63.4-104.6)	55.9* (51.7-63.5)	83.8* (71.2-108.8)	71.1* (63.2-88.5)	B-C: <.05 ^a
apo B100	72.6 (64.7-80.4)	64.1* (55.0-82.9)	87.1 (74.3-99.9)	85.2 (75.1-95.4)	B-C: <.05 ^a B-D: <.05 ^a
apo A-I	138.1 (131.41-144.75)	151.9* (138.30-173.84)	133.1 (124.44-141.67)	134 (126.16-141.84)	B-the rest: <.05 ^a
Lp(a)	9.4* (6.9-31.9)	16.4* (14.9-40.1)	14.9* (12.8-55.9)	45.1 (28.4-61.8)	D-the rest: <.05 ^a

*Apo indicates apolipoprotein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; TC, total cholesterol; Lp(a), lipoprotein (a).
aP value obtained with Kruskal-Wallis test.

^bP value obtained with Games-Howell test and one way ANOVA.

^cP value obtained with Tukey test and one way ANOVA.

Geometric means values are presented as means (95% CI).

The values for HDL-C, LDL-C^{13,22,23}, apo A-I^{22,23}, and Lp(a)²⁴⁻²⁷ recorded in earlier studies agreed with the present results. To our knowledge, this is the first study to analyze the interactions of Lp(a) and apo B100 with other cardiovascular risk factors.

It is difficult to offer an explanation of why the most stressful exercise should impair the lipid profile. However, it is well known that intense exercise, particularly eccentric exercise accompanied by falls and collisions,²⁸ leads to the release of proinflammatory cytokines²⁹⁻³¹ associated with abnormalities in metabolism.^{32,33} In addition, the endotoxin (lipopolysaccha-

ride [LPS]) of the cell walls of Gram-negative bacteria could be involved in the inflammatory response of several cellular disorders. The quantities of circulating LPS in healthy subjects are inappreciable, but they increase after strenuous exercise.³⁴ It has also been shown that LPS can impair lipid metabolism, increasing the activity of 3-hydroxymethylglutaryl coenzyme A reductase and reducing the level and activity of cholesterol 7 α -hydrolase mRNA.^{35,36}

Given the near or total absence of eccentric contractions, contact or falls in swimming, the acute cytokine, and endotoxin responses are probably reduced. Any

TABLE 4. Atherogenic Indices for the Different Groups of Sportsmen and Controls*

Atherogenic Index	Controls (A) (n=26)	Swimmers (B) (n=28)	Volleyball Players (C) (n=17)	Soccer Players (D) (n=23)	P
TC/HDL-C	3.0 ^a (2.7-3.6)	2.7 3.8 ^a (2.5-2.9)	3.5 (3.4-4.4)	B-C: <.001 ^a (3.1-3.9)	B-D: <.01 ^a
LDL-C/HDL-C	1.6 ^b (1.5-2.2)	1.5 (1.4-1.6)	2.6 (2.1-3.0)	2.2 (1.9-2.5)	B-C.D: <.01 ^a C-A: <.05 ^b
[Lp(a)+LDL-C]/HDL-C	1.9 ^b (1.7-2.8)	1.8 ^b (1.7-2.2)	3.0 ^b (2.6-3.9)	3.1 (2.6-3.6)	A-C.D: <.05 ^b B-C.D: <.01 ^b
apo B100/HDL-C	1.2 ^b (1.1-1.6)	1.1 ^b (0.9-1.4)	1.7 ^b (1.5-2.1)	1.6 (1.4-1.9)	B-C.D: <.05 ^b
apo B100/apo A-I	0.5 ^b (0.5-0.6)	0.4 ^b (0.3-0.6)	0.7 (0.6-0.7)	0.6 (0.6-0.7)	B-C: <.05 ^b B-D: <.01 ^b
(apo B100 \times LDL-C)/(apo A-I \times HDL-C)	0.8 ^b (0.7-1.5)	0.6 ^b (0.5-0.9)	1.5 ^b (1.2-2.4)	1.5 (1.1-1.9)	B-C: <.01 ^b B-D: <.01 ^b

*Apo indicates apolipoprotein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; TC, total cholesterol; Lp(a), lipoprotein (a).

^aP value obtained with Kruskal-Wallis test.

^bGeometric means

Values are presented as means (95% CI).




Fig. 1. Odds ratio of having a high lipid cardiovascular risk factor in the indicated groups. The black squares are the odds ratio values; the error bars express the 95% CI. Apo indicates apolipoprotein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; TC, total cholesterol; Lp(a), lipoprotein (a). *P value obtained with the test for trends.

negative changes occasioned by these molecules with respect to lipid metabolism are therefore avoided. In addition, the aerobic training practiced by the swimmers might be responsible for the healthy effects observed in their lipid profiles.

Atherosclerosis is a progressive disease characterized by the accumulation of lipids and fibrous ele-

ments in the major arteries. Its etiology is difficult to define but the risk factors include a poor lipid profile, increased fluid shear stress,³⁷ and increased susceptibility of the lipoproteins to oxidation.^{38,39} Accordingly, the negative lipid profiles of some of the subjects in this and other studies^{8,10,40} might be counteracted by adaptations to physical exercise, especially improved

TABLE 5. Odds Ratio for Interactions Between Lp(a) and Apo B100 and Lipid Cardiovascular Risk Factors*

Interactions	Swimmers (n=28)	Volleyball Players (n=17)	Soccer Players (n=23)	P†
High Lp(a) and high LDL-C	0	5.4 (0.5-56.5)	8.8 (1-80)	.0057
High Lp(a) and high TC/HDL-C	0	5.4 (0.5-56.5)	8.8 (1-80)	.0057
High Lp(a), high LDL-C and High TC/HDL-C	0	5.4 (0.5-56.5)	6.9 (0.7-64.6)	.0148
High apo B100 and high Lp(a)	0	2.6 (0.4-17.3)	5.2 (1-28.6)	.0021
High apo B100 and high LDL-C	0	3.8 (0.9-16.2)	4.2 (1.1-16.3)	<.001
High apo B100, high Lp(a) and high LDL-C	0	3.8 (0.91-16.2)	8.8 (0.97-80).0	.0057
Low Lp(a) and low LDL-C	1.7 (0.5-6.2)	0.3 (0.1-0.9)	0.1 (0-0.5)	<.001
Low Lp(a) and low TC/HDL-C	1.3 (0.4-4.4)	0.1 (0-0.4)	0.1 (0-0.4)	<.001
Low apo B100 and low Lp(a)	1.1 (0.3-3.7)	0.2 (0-0.7)	0.1 (0-0.4)	<.001
Low apo B100 and low LDL-C	4.8 (0.9-25.7)	0.4 (0.1-1.5)	0.3 (0.1-1.1)	<.001

*Apo indicates apolipoprotein; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; TC, total cholesterol; Lp(a), lipoprotein (a).

†P value obtained with the test for trends.

Values are expressed as the odds ratio (95% CI).

antioxidant and fibrinolytic capacities and reduced fluid shear stress.⁴¹⁻⁴³ This suggests the cut-off values for lipids associated with cardiovascular risk should be raised in sportspersons who compete at national level. Further studies are needed to corroborate this hypothesis. At present, no evidence exists to suggest that sportsmen subject to high levels of stressful physical exertion (e.g., soccer or rugby players) are at any greater risk of ischemic heart disease.

The present results take on great importance for occasional practitioners of strenuous sports (whose aim it is to improve their physical fitness on weekends⁴³) involving eccentric contractions, physical contact and falls (e.g., those who play soccer, rugby or basketball). Given the poorer physical fitness characterizing this sector of the population, their antioxidant and fibrinolytic activities and the fluid shear stress they experience may not be optimized. In such persons, the negative lipid profile caused by stressful physical exertion might not be counteracted by protective factors, leading to an increased risk of atherothrombogenesis. Further studies are necessary to corroborate this hypothesis.

CONCLUSIONS

In conclusion, the practice of sports involving a high degree of stressful exertion, such as soccer and volleyball, is accompanied by unfavorable plasma lipid and lipoprotein profiles. In contrast, sports with low levels of stressful exercise, such as swimming, appear to have a favorable effect on plasma lipid profiles.

REFERENCES

- Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, DuBose KD. Blood lipid and lipoprotein adaptations to exercise. A quantitative analysis. *Sports Med* 2001;31:1033-62.
- Balady GJ. Survival of the fittest-more evidence. *N Engl J Med* 2002;346:852-4.
- Shephard RJ, Balady GJ. Exercise as cardiovascular therapy. *Circulation* 1999;99:963-72.
- Williams P. Relationship of distance run per week to coronary heart disease risk factors in 8283 male runners: The National Runner's Health Study. *Arch Intern Med* 1997;157:191-8.
- Williams PT. Relationship of heart disease risk factors to exercise quantity and intensity. *Arch Intern Med* 1998;158:237-45.
- Leon AS, Sanche OA. Response of blood lipids to exercise training alone or combined with dietary intervention. *Med Sci Sports Exerc* 2001;33:S502-15.
- Aellen R, Hollmann W, Boutellier U. Effects of aerobic and anaerobic training on plasma lipoproteins. *Int J Sports Med* 1993;14:396-400.
- Eisenmann JC, Womack CJ, Reeves MJ, Pivarnik JM, Malina RM. Blood lipids in young distance runners. *Med Sci Sports Exerc* 2001;33:1661-6.
- Suárez-Mier MP, Aguilera B. Causas de muerte súbita asociada al deporte en España. *Rev Esp Cardiol* 2002;55:347-58.
- Mackinnon LT, Hubinger LM. Effects of exercise on lipoprotein(a). *Sports Med* 1999;28:11-24.
- Sharrett AR, Ballantyne CM, Coady SA, Heiss G, Sorlie PD, Castelli D, et al. Coronary heart disease prediction from lipoprotein cholesterol levels, triglycerides, lipoprotein(a), apolipoproteins A-I and B, and HDL density subfractions: The Atherosclerosis Risk in Communities (ARIC) Study. *Circulation* 2001;104:1108-13.
- Lovejoy K, Bachorik PS. Lp(a) measurements may be unreliable in stored sera. *Circulation* 1994;90:S504.
- Houmar J, Bruno N, Brune RK, McCammon MR, Israel RG, Barakat HA. Effects of exercise training on the chemical composition of plasma LDL. *Arterioscler Thromb* 1994;14:325-30.
- Stein JH, Rosenson RS. Lipoprotein Lp(a) excess and coronary heart disease. *Arch Intern Med* 1997;157:1170-6.
- Ji LL. Antioxidants and oxidative stress in exercise. *Proc Soc Exp Biol Med* 1999;222:283-92.
- Scanu AM. Atherothrombogenicity of lipoprotein (a): the debate. *Am J Cardiol* 1998;82:26Q-33Q.
- Kronenberg F, Kronenberg MF, Kiechl S, Trenkwalder E, Santer P, Oberhollenzer F, et al. Role of Lipoprotein(a) and Apolipoprotein(a) Phenotype in Atherogenesis. Prospective results from the Bruneck study. *Circulation* 1999;100:1154-60.
- Maher VMG, Brown BG, Marcovina SM, Hillger LA, Zhao XQ, Albers JJ. Effects of lowering elevated LDL cholesterol on the cardiovascular risk of lipoprotein(a). *JAMA* 1995;274:1771-4.
- Hopkins PN, Wu LL, Hunt SC, James BC, Vincent GM, Williams RR. Lipoprotein(a) interactions with lipid and nonlipid risk factors in early familial coronary artery disease. *Arterioscler Thromb Vasc Biol* 1997;17:2783-92.
- Delgado M, Gutiérrez A, Cano MD, Castillo MJ. Elimination of meat, fish and derived products from the Spanish Mediterranean diet: effect on plasma lipid profile. *Ann Nutr Metab* 1996;40:202-11.
- Mitchell JH, Haskell WL, Raven PB. Classifications of sports. *J Am Coll Cardiol* 1994;24:864-6.
- MacAuley D, McCrum EE, Stott G, Evans AE, Duly E, Trinick T, et al. Physical activity, lipids, apolipoproteins, and Lp(a) in the Northern Ireland Health and Activity Survey. *Med Sci Sports Exerc* 1996;28:720-36.
- Santos-Silva A, Rebelo MI, Castro EM, Belo L, Guerra A, Rego C, et al. Leukocyte activation, erythrocyte damage, lipid profile and oxidative stress imposed by high competition physical exercise in adolescents. *Clin Biochim Acta* 2001;306:119-26.
- Cardoso GC, Posadas C, Orvananos OO, Peniche C, Zamora J, Aguilar R, et al. Long distance runners and body-builders exhibit elevated plasma levels of lipoprotein(a). *Chem Phys Lipids* 1994;67-68:207-21.
- Cohen LI, Hartford CG, Rogers GG. Lipoprotein(a) and cholesterol in body builders using anabolic androgenic steroids. *Med Sci Sports Exerc* 1996;28:176-9.
- Holme I, Urdal P, Aerns S, Hjermann I. Exercise-induced increase in lipoprotein(a). *Atherosclerosis* 1996;122:97-104.
- Ponjee GA, Janssen EM, Hermans J, van Wersch JW. Regular physical activity and changes in risk factors for coronary heart disease: a nine months prospective study. *Eur J Clin Chem Clin Biochem* 1996;34:477-83.
- Smith LL. Acute inflammation: the underlying mechanism in delayed onset muscle soreness? *Med Sci Sports Exerc* 1991;23:542-51.
- Ostrowsky K, Rohde T, Asp S, Schjerling P, Pedersen BK. Pro- and anti-inflammatory cytokine balance in stressful exercise in humans. *J Physiol* 1999;515:287-91.
- Pedersen BK, Hoffman-Goetz L. Exercise and the immune system: regulation, integration, and adaptation. *Physiol Rev* 2000;80:1055-81.
- Moldoveanu AI, Shephard RJ, Shek PN. The cytokine response to physical activity and training. *Sports Med* 2001;31:115-44.

32. Hardardottir I, Grunfeld C, Feingold KR. Effects of endotoxin and cytokines on lipid metabolism. *Curr Opin Lipidol* 1994;5: 207-15.
33. Nonogaki K, Fuller GM, Fuentes NL, Moser AH, Staprans I, Grunfeld C, et al. Interleukin-6 stimulates hepatic triglyceride secretion in rats. *Endocrinology* 1995;136:2143-9.
34. Camus G, Poortmans J, Nys M, Deby-Dupont G, Duchateau J, Deby C, et al. Mild endotoxaemia and the inflammatory response induced by a marathon race. *Clin Sci* 1997;92:415-22.
35. Feingold KR, Spady DK, Pollock AS, Moser AH, Grunfeld C. Endotoxin, TNF, and IL-1 decrease cholesterol 7-alpha-hydroxylase mRNA levels and activity. *J Lipid Res* 1996;37:223-8.
36. Memon RA, Holleran WM, Moser AH, Uchida Y, Fuller J, Shigenaga JK, et al. Endotoxin and cytokines increase hepatic sphingolipid biosynthesis and produce lipoproteins enriched in ceramides and sphingomyelin. *Arterioscler Thromb Vasc Biol* 1998;18: 1257-65.
37. Badimon L, Martínez-González J. Endotelio en la protección vascular: nuevos conocimientos. *Rev Esp Cardiol* 2002;55:17-26.
38. Lusis AJ. Atherosclerosis. *Nature* 2000;407:233-41.
39. Sambola A, Fuster V, Badimon JJ. Papel de los factores de riesgo en la trombogenicidad sanguínea y los síndromes coronarios agudos. *Rev Esp Cardiol* 2003;56:1001-9.
40. Aellen R, Hollmann W, Boutellier U. Effects of aerobic and anaerobic training on plasma lipoproteins. *Int J Sports Med* 1993;14:396-400.
41. Kvernmo HD, Osterud B. The effect of physical conditioning suggests adaptation in procoagulant and fibrinolytic potencial. *Thromb Res* 1997;87:559-69.
42. Taylor CA, Hughes TJ, Zarins CK. Effect of exercise on hemodynamic conditions in the abdominal aorta. *J Vasc Surg* 1999;29: 1077-89.
43. Santarlasci PR. Weekend warriors. Common injuries in recreational athletes. *Adv Nurse Pract* 2000;8:42-6.