

Update: Innovation in Cardiology (II)

Telecardiology: Past, Present and Future

Jacob Thorsted Sørensen,^a Peter Clemmensen,^{b,*} and Maria Sejersten^b

^a Department of Cardiology, Aarhus University Hospital, Aarhus, Denmark

^b The Heart Center, Rigshospitalet, Copenhagen, Denmark

Article history:

Available online 19 January 2013

Keywords:

Telemedicine

Electrocardiogram

Acute myocardial infarction

ABSTRACT

Technological advances over the past decades have allowed improved diagnosis and monitoring of patients with acute coronary syndromes as well as patients with advanced heart failure. High-quality digital recordings transmitted wirelessly by cellular telephone networks have augmented the prehospital use of transportable electrocardiogram machines as well as implantable devices for arrhythmia monitoring and therapy. The impact of prehospital electrocardiogram recording and interpretation in patients suspected of acute myocardial infarction should not be underestimated. It enables a more widespread access to rapid reperfusion therapy, thereby reducing treatment delay, morbidity and mortality. Further, continuous electrocardiogram monitoring has improved arrhythmia diagnosis and dynamic ST-segment changes have been shown to provide important prognostic information in patients with acute ST-elevation myocardial infarction. Likewise, remote recording or monitoring of arrhythmias and vital signs seem to improve outcome and reduce the necessity of re-admissions or outpatient contacts in patients with heart failure or arrhythmias. In the future telemonitoring and diagnosis is expected to further impact the way we practice cardiology and provide better care for the patient with cardiovascular disease.

© 2012 Sociedad Española de Cardiología. Published by Elsevier España, S.L. All rights reserved.

Telecardiología: pasado, presente y futuro

RESUMEN

Los avances tecnológicos que se han producido en las últimas décadas han permitido mejorar el diagnóstico y la monitorización de los pacientes con síndromes coronarios agudos y los pacientes con insuficiencia cardiaca avanzada. Los registros digitales de alta calidad transmitidos de manera inalámbrica a través de redes de telefonía móvil han aumentado el uso prehospitalario de aparatos de electrocardiografía transportables y dispositivos implantables para la monitorización y el tratamiento de la arritmia. No se debe subestimar la importancia de los registros de electrocardiogramas prehospitalarios y su interpretación para pacientes con sospecha de infarto agudo de miocardio. Su empleo permite un acceso más amplio a una terapia de reperfusión rápida, con lo que se reducen el retraso en la aplicación del tratamiento, la morbilidad y la mortalidad. Además, la monitorización continua del electrocardiograma ha mejorado el diagnóstico de la arritmia, y se ha demostrado que la valoración de los cambios dinámicos del segmento ST aporta una información pronóstica importante para los pacientes con infarto de miocardio con elevación aguda del ST. De igual modo, parece que el registro o la monitorización a distancia de las arritmias y las constantes vitales mejora los resultados y reduce la necesidad de nuevos ingresos o contactos asistenciales ambulatorios de los pacientes con insuficiencia cardiaca o arritmias. En el futuro, es de prever que la telemonitorización y el diagnóstico influyan aún más en la práctica de la cardiología y aporten una mejor asistencia para el paciente con enfermedad cardiovascular.

© 2012 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Palabras clave:

Telemedicina

Electrocardiograma

Infarto agudo de miocardio

INTRODUCTION

Patients with ST-elevation myocardial infarction (STEMI) should receive reperfusion therapy as soon as possible to reduce mortality and morbidity. Guidelines recommend primary

percutaneous coronary intervention (PCI) as the preferable strategy, if this can be performed in a timely manner.^{1,2} One of the main obstacles to a more widespread use of primary PCI is the limited accessibility to PCI centers operating on a 24-h basis. Thus, to treat STEMI patients within the recommended time limits, early diagnosis is essential in order for patients to be transported to a PCI center in due time. This has led to an increased interest in prehospital diagnosis. The diagnostic modality of choice has been the electrocardiogram (ECG). The ECG is recorded in the field and

* Corresponding author: The Heart Center, Rigshospitalet, Blegdamsvej 9, DK-2100 København Ø, Copenhagen, Denmark.

E-mail address: peter.clemmensen@rh.regionh.dk (P. Clemmensen).

Abbreviations

ECG: electrocardiogram
 ICD: implantable cardioverter-defibrillator
 PCI: percutaneous coronary intervention
 STEMI: ST-elevation myocardial infarction

interpreted by emergency physicians, paramedics or cardiologists receiving the ECG in the hospital by wireless transmission (Figure). This method is very robust and has been successfully implemented in several countries worldwide.^{3–9} Prehospital ECG diagnosis and field triage directly to primary PCI has been shown to decrease treatment delay significantly and is associated with a lower mortality.^{10,11}

The prehospital use of continuous ECG surveillance in the form of ST monitoring has been shown to provide important prognostic information and might further improve triage for tailored treatment regimens.¹² Both ST monitoring and ECG diagnosis rely on optimal equipment for ECG recording. Through the years new hardware and analysis algorithms along with more practical lead positioning systems have been introduced and might help to further expand the indications and use of prehospital ECG recording and diagnosis.

PREHOSPITAL ELECTROCARDIOGRAM RECORDING

In the 1970s Uhley reported their experiences in transmitting single-lead telemetry from the ambulance to the hospital for arrhythmia monitoring in patients with suspected cardiovascular disease.¹³ For several years, however, it was not possible to transmit a diagnostic-quality 12-lead ECG due to technical difficulties and unstable data networks. In 1987 Grim et al. published a paper on successful implementation of diagnostic-quality, 12-lead ECG transmission using new error-correcting digital transmission via cellular phone links.¹⁴ The new

transmission technology was validated in other studies, showing the possible benefit of early and reliable diagnosis of acute myocardial infarction.^{15,16}

In the late 1990s the terms STEMI and non-ST-elevation acute myocardial infarction were introduced to ensure a more clinically and therapeutically relevant triage of acute myocardial infarction patients. This primarily relied on the large thrombolysis trials showing an improved outcome with the immediate use of lytic therapy for patients with STEMI.^{17,18} Early reperfusion therapy thereby became an important objective in the optimal treatment of STEMI and this led to a further interest in early diagnosis and optimal triage. In 2000 Wall et al. reported that prehospital diagnosis of acute myocardial infarction was associated with more rapid reperfusion therapy by thrombolysis,¹⁹ but the focus was still mainly on the in-hospital, emergency department delays in diagnosis.

However, this changed as several randomized trials and meta-analyses proved primary PCI to be a superior reperfusion strategy in STEMI patients, if performed in a timely manner.^{20–23} The limited access to PCI centers increased the focus on early diagnosis, allowing the patient direct admission or transfer to a PCI center as soon as possible.

In 2005 Terkelsen et al. demonstrated the benefit of prehospital ECG diagnosis and field triage to a primary PCI center in patients with STEMI. The study comprised 161 patients. A prehospital ECG was recorded in 106 patients, 21 of them admitted directly to a primary PCI center. The time from first medical contact to balloon was a median 81 min shorter in patients with prehospital diagnosis and field triage compared to patients diagnosed in the hospital.²⁴ Prehospital ECG diagnosis and field triage not only allows the patient to bypass the emergency room of a local, non-PCI-capable hospital, it also reduces in-hospital delays, as the catheterization laboratory is able to prepare for the imminent arrival of the patient. Sejersten et al. have further demonstrated this.⁶ In their study, door-to-PCI delay was significantly reduced in patients with a prehospital diagnosis compared with in-hospital diagnosis (34 vs 97 min, $P<.001$).




Figure. An example of the set-up for prehospital electrocardiogram diagnosis. The electrocardiogram is recorded in the ambulance and sent wirelessly to a primary percutaneous coronary intervention center. The electrocardiogram is interpreted by the on-call cardiologist and after talking to the patient and/or paramedic a decision is made to either redirect the patient for primary percutaneous coronary intervention, in case of ST-elevation myocardial infarction, or send the patient to the nearest local hospital for further diagnosis and initial pharmacological therapy. PCI, percutaneous coronary intervention; STEMI, ST-elevation myocardial infarction.

Table 1
Time From First Medical Contact or Symptom Onset to Balloon in Recent International Publications on Prehospital Electrocardiogram Diagnosis and Triage

	First medical contact to balloon, min		Symptom onset to balloon, min	
	Prehospital diagnosis	No prehospital diagnosis	Prehospital diagnosis	No prehospital diagnosis
Terkelsen et al., ²⁴ 2005	87	168	129	250
Sejersten et al., ⁶ 2008	89*	142*	126*	206*
Le May et al., ⁷ 2008	N/A	N/A	158	230
Pedersen et al., ¹⁰ 2009	90	120	150	200
Dieker et al., ⁸ 2010	89	141	147	206
Martinoni et al., ⁹ 2010	50	85	154	208
Sørensen et al., ¹¹ 2011	90	141	138	199

N/A, not available.

* Recalculated values from original data.

Studies from many countries have reproduced these results despite varying geographic settings, health-care organizations and prehospital logistic set-ups (Table 1).

Recently Sørensen et al. evaluated the routine use of prehospital ECG diagnosis and field triage in a larger region comprising both urban and rural surroundings.¹¹ The authors found that prehospital diagnosis and triage reduces the importance of distance to primary PCI center, in as much as time from first medical contact to balloon inflation only varied 9 min between patients belonging to urban versus rural hospitals, with a median 10 km versus 40 km distance to the PCI center (Table 2). Further the study showed a decreased mortality in patients with prehospital diagnosis and triage. A recent Dutch study further emphasized the importance and value of prehospital diagnosis for patients living far from a PCI center.²⁵

ELECTROCARDIOGRAM BODY-SURFACE MAPPING

One of the limitations of the 12-lead ECG, whether used in-hospital or in the prehospital phase, is the lack of sensitivity. Not all anatomic parts of the left ventricle are equally well covered by conventional leads. The infarct location most commonly missed on 12-lead ECG is the inferoposterior myocardial infarction. It was established many years ago that this limitation can be overcome by using additional leads, like V₇-V₉.²⁶ Actual body-surface mapping often uses an 80-lead ECG, which improves the ability to detect acute myocardial infarction and thus improve the sensitivity of electrocardiography in large clinical series.²⁷ The sensitivity for detection of troponin-confirmed acute myocardial infarction approaches 80% and c-statistics improve with a factor of almost 0.10.²⁷ Right ventricular involvement complicating inferior STEMI can also be better detected by 80-lead body-surface mapping ECG.²⁸ Recently a randomized trial compared the prevalence, clinical care patterns, and clinical outcomes of patients with STEMI identified on 80-lead but not on 12-lead (80-lead-only STEMI). The primary outcome of the trial was door-to-sheath time in patients with 80-lead-only STEMI versus patients with STEMI identified by 12-lead alone (12-lead STEMI). The 80-lead ECG

Table 2
Relationship of Distance to Percutaneous Coronary Intervention Center and Time to Balloon Inflation

Distance to PCI center, km	0-10	11-30	31-50	> 50
FMC to balloon inflation, min	78	89	93	106

FMC, first medical contact; PCI, percutaneous coronary intervention.

provided an incremental 27.5% increase in STEMI detection versus the 12-lead. Patients with 80-lead-only STEMI have adverse outcomes similar to those of 12-lead STEMI patients but are treated with delayed or conservative invasive strategies.²⁹

The 80-lead ECG body-surface map has also been used in the prehospital phase despite being more time-consuming to mount. Even in this setting the sensitivity of the acute myocardial infarction detection method is comparable to findings in the more protected in-hospital setting.³⁰ Despite these findings, body-surface mapping for evaluation of patients with chest pain has not gained widespread usage. This could change with the invention of easy to apply technologies, which are appearing for commercial use.

NOVEL ELECTROCARDIOGRAM RECORDING AND LEAD SYSTEMS

As the prehospital setting often implies working under difficult circumstances with patients in a very poor clinical status, there has been some focus on developing ECG monitoring systems for simple, fast application with a durable design.

One such system (the EASI lead system), using only 4 leads to extract a full, diagnostic “12-lead” ECG, has been compared with a conventional 12-lead ECG by Sejersten et al.³¹ The authors report that the 4-lead EASI system was just as good for detecting acute, myocardial ischemia as the standard 12-lead ECG. Drew et al. also investigated the use of a more simple lead system in the Synthesized Twelve-lead ST Monitoring and Real-time Tele-electrocardiography initiative. The 5-lead ECG system used in this series of studies has proven effective in diagnosing ischemia and has reduced treatment delay in the study region (Northern California, United States).^{32,33}

TELEHEALTHCARE FOR PATIENT AND DEVICE MONITORING

The combination of an aging population and continued developments of modern therapy within cardiology and internal medicine in general have led to an increasing prevalence of chronic diseases, including heart failure,³⁴ and the need for implantable devices.³⁵ This development has increased the need for continued and close contact between the patient and the healthcare system to ensure the optimal treatment for the individual. Consequently, the healthcare system is challenged by the still-increasing need for qualified healthcare professionals, and resources in general, as well as increasing costs. One way to approach this challenge is the use of telehealthcare, allowing the patient to be in close contact with the appropriate healthcare provider and exchanging valuable information needed for managing the patient's disease.

Telehealthcare is defined as personal healthcare over a distance and consists of the following components: *a*) data provided by the patient (eg, blood pressure, pulse, weight, subjective information about health); *b*) electronic transmission of data from the patient to a healthcare professional, and *c*) personalized feedback tailored to the individual patient.³⁶ Telehealthcare can be based on either real-time (synchronous), store-and-forward (asynchronous), or hybrid systems.³⁷ Synchronous telehealthcare requires the availability of patient and healthcare provider at the same time and real-time processing of the patient's data, while asynchronous telehealthcare provides more flexibility for both the patient and healthcare provider.

The technology used in telehealthcare is a composite of portable medical imaging devices (eg, blood pressure device), computer/smart phone, and the wireless communication infrastructure, requiring a stable network for data transmission.³⁸ The transmitted data may then be stored at a receiving station for momentary or later processing and review.

Four generations of telehealthcare for remote management of patients have been proposed³⁸: *a*) 1st generation, a system of nonreactive data collection and analysis, with asynchronous transmission of data to a healthcare provider, eg, from an event recorder; *b*) 2nd generation, a system with a non-immediate analytical or decision-making structure, with synchronous data transfer, allowing the healthcare provider to view data and recognize important changes, but delays may occur if the system is only active during the day; *c*) 3rd generation, a remote patient management system with consistent analytical and decision-making support by a healthcare professional, even outside of office hours, and *d*) 4th generation, a fully integrated remote management system, where data from both invasive and noninvasive telemedical devices are linked in a telemedical platform and available for both the primary care provider and the telemedical center.

A wide variety of both noninvasive and invasive variables are available for telehealthcare within cardiology. Most are recorded manually by the patients via a device (eg, systolic and diastolic blood pressure, pulse, 3-lead ECG, heart rate variability, body-weight, oxygen saturation, blood glucose, natriuretic peptides). Others obtained from invasive devices are recorded automatically (eg, impedance, incidence of arrhythmias, pulmonary and left atrial pressure).³⁸

TELEHEALTHCARE IN HEART FAILURE

Heart failure is associated with substantial morbidity, mortality, and healthcare costs.^{34,39} Patients are often followed closely in heart failure clinics but admittance to a cardiology department is frequently required, and the 30-day readmission rate is high.⁴⁰ Telehealthcare may be a solution, providing a way to identify and monitor subclinical congestion. The earlier identification and treatment of congestion together with improved coordination of care may prevent hospitalization.

Ideally, contact between a patient and a healthcare provider is established with technology enabling the two parties to see and hear each other via a microphone and a computer screen over large distances using a telephone or satellite connection. One device is installed in the patient's home and one at the heart failure clinic. Preferably the patient's device is portable and capable of both videoconference and data transmission, and can be managed by the healthcare professional using a remote control.

According to individual arrangements patients could be instructed to record blood pressure, heart rhythm and weight, and give a subjective description of symptoms every morning, and then transmit data daily, just prior to or during a scheduled remote

follow-up. Data can then be reviewed just before the videoconference with the patient, so treatment can be altered if needed. It is important that the healthcare professional be empowered to change medication regimens so delay is diminished and the correct treatment established in a timely manner.

Several advantages of the use of telehealthcare in heart failure have been reported.^{36,41} Patients get a better understanding of their disease by being responsible for monitoring and being involved in some self-managing, which enhances patient education and empowerment. This supports adherence to the planned treatment, but also ensures preventive care with early detection of disease exacerbations and timely management. Travel and waiting time is reduced or even abolished for the patient, and quality of life improves. Additionally, patients are at home in comfortable and safe surroundings and still in close contact with the hospital. An advantage for the hospital is that the technology leaves room for new and unstable patients, and long admittances to hospital may be prevented.

However, the effectiveness of telehealthcare by reducing morbidity and mortality in heart failure patients has not been established because results have been conflicting. Accordingly, 2 metaanalyses^{42,43} have suggested that telehealthcare reduced morbidity and mortality but 2 prospective randomized clinical trials not including the metaanalyses did not support those findings.^{44,45} A possible explanation for this difference may be that the metaanalyses combined many small dissimilar studies of patients with varying risk profiles followed for different durations of time and with differences between studies in the intervention used. A newly published prospective randomized trial of 3230 patients with either obstructive pulmonary disease or heart failure showed that telehealthcare was associated with lower mortality, emergency admission rates, and shorter length of admission.⁴⁶ An ongoing Danish study randomizes chronic heart failure patients to either traditional treatment in outpatient clinics or monitoring by telehealthcare. Another part of the study randomizes patients admitted with acute heart failure to traditional treatment in hospital versus early discharge and daily contact with the hospital via telehealthcare. The hypothesis is that at-home treatment of heart failure patients is feasible, effective, and safe, and thereby releases resources to other patient groups (personal communication).

In the studies mentioned above telehealthcare was based on noninvasive data, but other studies have investigated the management of heart failure patients using data from implantable devices. An increase in pulmonary vascular congestion is reflected by increased impedance, and thus intrathoracic impedance monitoring may be a valuable tool in the management of heart failure in patients, such as those with an implantable cardioverter-defibrillator (ICD)⁴⁷. This is supported by a study that found significantly greater sensitivity to predict worsening heart failure events for intrathoracic impedance monitoring compared to acute weight increase.⁴⁸ Likewise, left ventricular filling pressures and pulmonary artery pressures are correlated with clinical congestion, functional limitation, and poor prognosis in heart failure patients.⁴⁹ Since the intracardiac and pulmonary artery pressures increase several days to weeks before the onset of clinical symptoms of congestion, close monitoring may provide early warning and assist appropriate management, including changes in medication. Abraham et al.⁵⁰ used a wireless implantable hemodynamic monitoring system to measure pulmonary artery pressure, and reported that the use of the system significantly reduced the duration of hospital stays and the rate of hospital admissions related to heart failure, the latter by 30%.

In a study that monitored left atrial pressure, this monitoring was associated with reduced risk of acute decompensation and death.⁵¹ Additional randomized clinical trials using third- or

fourth-generation telemedical systems that combine noninvasive and invasive variables with daily monitoring and management are needed to fully clarify the value of telehealthcare in cardiology.

TELEHEALTHCARE AND IMPLANTABLE DEVICES

The number of patients with pacemakers and ICDs has increased tremendously over the years as indications have broadened to include more patient groups.³⁵ According to guidelines, patients with a pacemaker or an ICD need a follow-up at 3–12 months and 3–6 months, respectively.⁵² Usually the follow-up visits include both an evaluation of the device function (including battery status and capture thresholds) and a clinical examination, which may result in reprogramming of the device or changes in medical treatment. These time-consuming visits can be a burden to the patient, as they often require long transportation, waiting time, and time off work. For the outpatient clinic, the many visits require both space and a large staff because of the workload associated with every visit.

Remote monitoring and remote follow-up by the use of telehealthcare offers an alternative management strategy.⁵³ Preferably, the system should be able to automatically transmit data stored in the device to the outpatient clinic using the wireless global system for mobile communication network or a landline. Automatic wireless data transmission requires a pacemaker or ICD equipped with a micro-antenna for communication with a transmitter located close to the patient. This setup diminishes the need for patient compliance and increases the frequency of transmissions.

Several companies provide systems for automatic wireless data transmission.²¹ Data stored in the device's memory (eg, battery voltage, lead characteristics, arrhythmias, alerts) can be transmitted. The alerts may be based on a change in device performance (battery status, lead impedance), programming (disabling of ventricular fibrillation therapy, insufficient safety margins for sensing or capture), or medical data (arrhythmias, indication of lung fluid accumulation). Events that trigger an alert can be defined according to the individual patient. Data is usually transmitted to a central database, where it is processed and made available to the physician on a secure webpage. Additional notification by e-mail, SMS, fax, or phone messages may be valuable when critical data are available for consultation.⁵³

Many pacemakers and ICDs can automatically run tests like battery status, lead impedances, sensing and capture thresholds that used to be manually performed at the outpatient clinic.⁵³ Both scheduled follow-up, with transmission of pre-defined data, and unscheduled monitoring can then be performed. Daily monitoring allows transmission of data with any predefined alerts to a physician. Consequently, remote monitoring may improve patient safety and quality of care. Depending on the need for reprogramming, clinic follow-ups are still needed because due to safety issues no system yet allows remote device programming.

The safety and clinical benefits conferred by telehealthcare in patients with implantable devices have been validated in several studies.^{54–56} In the COMPAS trial,⁵⁴ long-term remote monitoring of pacemaker recipients was a safe substitute for conventional follow-ups, decreased the number of ambulatory visits, and enabled early detection of important clinical and device-related adverse events. Another study showed that half of the regularly scheduled visits could be omitted by the use of remote monitoring, without impairing patient safety.⁵⁵ Additionally, only 6% of patients undergo device reprogramming or admittance to hospital during clinic visits, and thus 94% of these visits might as well be executed by remote monitoring.⁵⁶ Remote monitoring could

diagnose 99% of arrhythmia- or device-related problems in patients with an ICD, when combined with a clinical follow-up. In addition to fewer scheduled clinic visits, unscheduled visits following an ICD shock might be prevented by remote follow-up; after such an event the patient may perform manual upload of data, transmitted to the healthcare provider for immediate determination of whether the shock was appropriate or not, and whether device reprogramming or medical modifications are needed.⁵⁷ Finally, several studies have demonstrated that remote monitoring reduces the time to event recognition and diagnosis and the time from the event to a clinical decision for the individual patient.^{58,59}

Remote monitoring is also advantageous in patients with paroxysmic atrial fibrillation because alerts could identify the start-point of atrial fibrillation, resulting in an unscheduled follow-up either in office or by phone, and appropriate interventions (eg, antiarrhythmic medication, anticoagulation or antiplatelet medication, external cardioversion, device reprogramming).⁶⁰ In general, satisfaction is high and both patient and healthcare personals prefer remote follow-up over in-clinic visits.^{61,62}

FUTURE TELEMEDICINE IN CARDIOLOGY

Telemedicine has revolutionized modern cardiology and provided the opportunity for consultations between doctors separated by long distances, even between continents. In the acute phase of heart disease, telemedicine has helped to deliver optimal care faster and in chronic disease consultation between doctor and patient decreases the need for outpatient visits. As these systems are further improved, so are patient outcomes. In the future, it is likely that smaller implantable devices will become available to provide wireless feedback to patients and doctors alike via interfaces that connect through computers or mobile devices.

CONFLICTS OF INTEREST

None declared.

REFERENCES

1. Steg PG, James SK, Atar D, Badano LP, Blömostrom-Lundqvist C, Borger MA, et al. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology (ESC). *Eur Heart J*. 2012;33:2569–619.
2. Kushner FG, Hand M, Smith Jr SC, King 3rd SB, Anderson JL, Antman EM, et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update) a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2009;54:2205–41. <http://dx.doi.org/10.1016/j.jacc.2009.10.015>.
3. Terkelsen CJ, Nørgaard BL, Lassen JF, Gerdes JC, Ankersen JP, Rømer F, et al. Telemedicine used for remote prehospital diagnosing in patients suspected of acute myocardial infarction. *J Intern Med*. 2002;252:412–20.
4. Clemmensen P, Sejersten M, Sillesen M, Hampton D, Wagner GS, Loumann-Nielsen S. Diversion of ST-elevation myocardial infarction patients for primary angioplasty based on wireless prehospital 12-lead electrocardiographic transmission directly to the cardiologist's handheld computer: a progress report. *J Electrocardiol*. 2005;38 Suppl 4:194–8.
5. Adams GL, Campbell PT, Adams JM, Strauss DG, Wall K, Patterson J, et al. Effectiveness of prehospital wireless transmission of electrocardiograms to a cardiologist via hand-held device for patients with acute myocardial infarction (from the Timely Intervention in Myocardial Emergency. NorthEast Experience [TIME-NE]). *Am J Cardiol*. 2006;98:1160–4.
6. Sejersten M, Sillesen M, Hansen PR, Nielsen SL, Nielsen H, Trautner S, et al. Effect on treatment delay of prehospital teletransmission of 12-lead electrocardiogram to a cardiologist for immediate triage and direct referral of patients with ST-segment elevation acute myocardial infarction to primary percutaneous coronary intervention. *Am J Cardiol*. 2008;101:941–6.

7. Le May MR, So DY, Dionne R, Glover CA, Froeschl MP, Wells GA, et al. A citywide protocol for primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med*. 2008;358:231–40.
8. Dieker HJ, Liem SS, El Aidi H, Van Grunsven P, Aengevaeren WR, Brouwer MA, et al. Pre-hospital triage for primary angioplasty: direct referral to the intervention center versus interhospital transport. *JACC Cardiovasc Interv*. 2010;3:705–11. <http://dx.doi.org/10.1016/j.jcin.2010.04.010>.
9. Martinoni A, Politi A, De Servi S. [Role of prehospital electrocardiogram in the early diagnosis of ST-elevation myocardial infarction]. *G Ital Cardiol (Rome)*. 2010;11(10 Suppl 1):S53–6.
10. Pedersen SH, Galatius S, Hansen PR, Mogelvang R, Abildstrom SZ, Sørensen R, et al. Field triage reduces treatment delay and improves long-term clinical outcome in patients with acute ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. *J Am Coll Cardiol*. 2009;54:2296–302.
11. Sørensen JT, Terkelsen CJ, Nørgaard BL, Trautner S, Hansen TM, Bøtker HE, et al. Urban and rural implementation of pre-hospital diagnosis and direct referral for primary percutaneous coronary intervention in patients with acute ST-elevation myocardial infarction. *Eur Heart J*. 2011;32:430–6.
12. Terkelsen CJ, Kalltoft AK, Nørgaard BL, Bottcher M, Lassen JF, Clausen K, et al. ST changes before and during primary percutaneous coronary intervention predict final infarct size in patients with ST elevation myocardial infarction. *J Electrocardiol*. 2009;42:64–72. <http://dx.doi.org/10.1016/j.jelectrocard.2008.08.038>.
13. Uhley HN. Electrocardiographic telemetry from ambulances. A practical approach to mobile coronary care units. *Am Heart J*. 1970;80:838–42.
14. Grim P, Feldman T, Martin M, Donovan R, Nevins V, Childers RW. Cellular telephone transmission of 12-lead electrocardiograms from ambulance to hospital. *Am J Cardiol*. 1987;60:715–20.
15. Kudenchuk PJ, Maynard C, Cobb LA, Wirkus M, Martin JS, Kennedy JW, et al. Utility of the prehospital electrocardiogram in diagnosing acute coronary syndromes: the Myocardial Infarction Triage and Intervention (MITI) Project. *J Am Coll Cardiol*. 1998;32:17–27.
16. Giovas P, Papadoyannis D, Thomakos D, Papazachos G, Rallidis M, Soulis D, et al. Transmission of electrocardiograms from a moving ambulance. *J Telemed Telecare*. 1998;4 Suppl 1:5–7.
17. Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico (GISSI). *Lancet*. 1986;1:397–402.
18. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. *Lancet*. 1988;2:349–60.
19. Wall T, Albright J, Livingston B, Isley L, Young D, Nanny M, et al. Prehospital ECG transmission speeds reperfusion for patients with acute myocardial infarction. *N C Med J*. 2000;61:104–8.
20. Grines CL, Westerhausen DR, Grines LL, Hanlon JT, Logemann TL, Niemela M, et al. A randomized trial of transfer for primary angioplasty versus on-site thrombolysis in patients with high-risk myocardial infarction: the Air Primary Angioplasty in Myocardial Infarction study. *J Am Coll Cardiol*. 2002;39:1713–9.
21. Widimský P, Budesínský T, Vorác D, Groch L, Zelízko M, Aschermann P, et al. Long distance transport for primary angioplasty vs immediate thrombolysis in acute myocardial infarction. Final results of the randomized national multicentre trial—PRAGUE-2. *Eur Heart J*. 2003;24:94–104.
22. Andersen HR, Nielsen TT, Rasmussen K, Thuesen L, Kelbaek H, Thyssen P, et al. A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. *N Engl J Med*. 2003;349:733–42.
23. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet*. 2003;361:13–20.
24. Terkelsen CJ, Lassen JF, Nørgaard BL, Gerdes JC, Poulsen SH, Bendix K, et al. Reduction of treatment delay in patients with ST-elevation myocardial infarction: impact of pre-hospital diagnosis and direct referral to primary percutaneous coronary intervention. *Eur Heart J*. 2005;26:770–7.
25. Atary JZ, De Visser M, Van den Dijk R, Bosch J, Liem SS, Antoni ML, et al. Standardised pre-hospital care of acute myocardial infarction patients: MISSION! guidelines applied in practice. *Neth Heart J*. 2010;18:408–15.
26. Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, et al. Third universal definition of myocardial infarction. *Eur Heart J*. 2012;33:2551–67.
27. Owens C, McClelland A, Walsh S, Smith B, Adgey J. Comparison of value of leads from body surface maps to 12-lead electrocardiogram for diagnosis of acute myocardial infarction. *Am J Cardiol*. 2008;102:257–65.
28. Ornato JP, Menown IB, Peberdy MA, Kontos MC, Riddell JW, Higgins GL, et al. Body surface mapping vs 12-lead electrocardiography to detect ST-elevation myocardial infarction. *Am J Emerg Med*. 2009;27:779–84.
29. Hoekstra JW, O'Neill BJ, Pride YB, Lefebvre C, Diercks DB, Peacock WF, et al. Acute detection of ST-elevation myocardial infarction missed on standard 12-Lead ECG with a novel 80-lead real-time digital body surface map: primary results from the multicenter OCCULT MI trial. *Ann Emerg Med*. 2009;54:779–88. e1.
30. Owens CG, McClelland AJ, Walsh SJ, Smith BA, Tomlin A, Riddell JW, et al. Prehospital 80-LAD mapping: does it add significantly to the diagnosis of acute coronary syndromes? *J Electrocardiol*. 2004;37 Suppl:223–32.
31. Sejersten M, Wagner GS, Pahlm O, Warren JW, Feldman CL, Horáček BM. Detection of acute ischemia from the EASI-derived 12-lead electrocardiogram and from the 12-lead electrocardiogram acquired in clinical practice. *J Electrocardiol*. 2007;40:120–6.
32. Drew B. Pre-hospital synthesized 12-lead ECG ischemia monitoring with trans-telephonic transmission in acute coronary syndromes. Pilot study results of the ST SMART trial. *J Electrocardiol*. 2004;37 Suppl:214–21.
33. Drew BJ, Sommargren CE, Schindler DM, Benedict K, Zegre-Hemsey J, Glancy JP. A simple strategy improves prehospital electrocardiogram utilization and hospital treatment for patients with acute coronary syndrome (from the ST SMART Study). *Am J Cardiol*. 2011;107:347–52.
34. Ho KK, Pinsky JL, Kannel WB, Levy D. The epidemiology of heart failure: the Framingham Study. *J Am Coll Cardiol*. 1993;22 Suppl 4:A6–13.
35. Epstein AE, Dimarco JP, Ellenbogen KA, Estes NA, Freedman RA, Gettes LS, et al. ACC/AHA/HRS 2008 Guidelines for device-based therapy of cardiac rhythm abnormalities. *Heart Rhythm*. 2008;5:e1–62.
36. McLean S, Protti D, Sheikh A. Telehealthcare for long term conditions. *BMJ*. 2011;342:d120. <http://dx.doi.org/10.1136/bmj.d120>.
37. Allely EB. Synchronous and asynchronous telemedicine. *J Med Syst*. 1995;19:207–12.
38. Anker SD, Koehler F, Abraham WT. Telemedicine and remote management of patients with heart failure. *Lancet*. 2011;378:731–9.
39. Bui AL, Fonarow GC. Home monitoring for heart failure management. *J Am Coll Cardiol*. 2012;59:97–104.
40. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med*. 2009;360:1418–28.
41. Sood S, Mbarika V, Jugoo S, Dookhy R, Doarn CR, Prakash N, et al. What is telemedicine? A collection of 104 peer-reviewed perspectives and theoretical underpinnings. *Telemed J E Health*. 2007;13:573–90.
42. Klersy C, De Silvestri A, Gabutti G, Regoli F, Auricchio A. A meta-analysis of remote monitoring of heart failure patients. *J Am Coll Cardiol*. 2009;54:1683–94.
43. Inglis SC, Clark RA, McAlister FA, Ball J, Lewinter C, Cullington D, et al. Structured telephone support or telemonitoring programmes for patients with chronic heart failure. *Cochrane Database Syst Rev*. 2010;8:CD007228.
44. Chaudhry SI, Matterna JA, Curtis JP, Spertus JA, Herrin J, Lin Z, et al. Telemonitoring in patients with heart failure. *N Engl J Med*. 2010;363:2301–9.
45. Koehler F, Winkler S, Schieber M, Sechtem U, Stangl K, Böhm M, et al. Impact of remote telemedical management on mortality and hospitalizations in ambulatory patients with chronic heart failure: the telemedical interventional monitoring in heart failure study. *Circulation*. 2011;123:1873–80.
46. Steventon A, Bardsley M, Billings J, Dixon J, Doll H, Hirani S, et al. Effect of telehealth on use of secondary care and mortality: findings from the Whole System Demonstrator cluster randomised trial. *BMJ*. 2012;344:e3874. <http://dx.doi.org/10.1136/bmj.e3874>.
47. Yu CM, Wang L, Chau E, Chan RH, Kong SL, Tang MO, et al. Intrathoracic impedance monitoring in patients with heart failure: correlation with fluid status and feasibility of early warning preceding hospitalization. *Circulation*. 2005;112:841–8.
48. Abraham WT, Compton S, Haas G, Foreman B, Canby RC, Fishel R, et al. Intrathoracic impedance vs daily weight monitoring for predicting worsening heart failure events: results of the Fluid Accumulation Status Trial (FAST). *Congest Heart Fail*. 2011;17:51–5.
49. Zile MR, Bennett TD, St John Sutton M, Cho YK, Adamson PB, Aaron MF, et al. Transition from chronic compensated to acute decompensated heart failure: pathophysiological insights obtained from continuous monitoring of intracardiac pressures. *Circulation*. 2008;118:1433–41.
50. Abraham WT, Adamson PB, Bourge RC, Aaron MF, Costanzo MR, Stevenson LW, et al. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: a randomised controlled trial. *Lancet*. 2011;377:658–66.
51. Adamson PB, Magalski A, Braunschweig F, Böhm M, Reynolds D, Steinhaus D, et al. Ongoing right ventricular hemodynamics in heart failure: clinical value of measurements derived from an implantable monitoring system. *J Am Coll Cardiol*. 2003;41:565–71.
52. Wilkoff BL, Auricchio A, Brugada J, Cowie M, Ellenbogen KA, Gillis AM, et al. HRS/EHRA Expert Consensus on the Monitoring of Cardiovascular Implantable Electronic Devices (CIEDs): description of techniques, indications, personnel, frequency and ethical considerations: developed in partnership with the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA); and in collaboration with the American College of Cardiology (ACC), the American Heart Association (AHA), the European Society of Cardiology (ESC), the Heart Failure Association of ESC (HFA), and the Heart Failure Society of America (HFSA). Endorsed by the Heart Rhythm Society, the European Heart Rhythm Association (a registered branch of the ESC), the American College of Cardiology, the American Heart Association. *Europace*. 2008;10:707–25.
53. Burri H, Senouf D. Remote monitoring and follow-up of pacemakers and implantable cardioverter defibrillators. *Europace*. 2009;11:701–9.
54. Mabo P, Victor F, Bazin P, Ahres S, Babuty D, Da Costa A, et al. A randomized trial of long-term remote monitoring of pacemaker recipients (the COMPAS trial). *Eur Heart J*. 2012;33:1105–11.
55. Brugada P. What evidence do we have to replace in-hospital implantable cardioverter defibrillator follow-up? *Clin Res Cardiol*. 2006;95 Suppl 3:3–9.
56. Heidbüchel H, Lioen P, Foulon S, Huybrechts W, Ector J, Willems R, et al. Potential role of remote monitoring for scheduled and unscheduled evaluations of patients with an implantable defibrillator. *Europace*. 2008;10:351–7.
57. Santini M, Ricci RP, Lunati M, Landolina M, Perego GB, Marzegalli M, et al. Remote monitoring of patients with biventricular defibrillators through the

- CareLink system improves clinical management of arrhythmias and heart failure episodes. *J Interv Card Electrophysiol.* 2009;24:53–61.
58. Crossley GH, Boyle A, Vitense H, Chang Y, Mead RH; CONNECT Investigators. The CONNECT (Clinical Evaluation of Remote Notification to Reduce Time to Clinical Decision) trial: the value of wireless remote monitoring with automatic clinician alerts. *J Am Coll Cardiol.* 2011;57:1181–9.
59. Varma N, Epstein AE, Irimpen A, Schweikert R, Love C; TRUST Investigators. Efficacy and safety of automatic remote monitoring for implantable cardioverter-defibrillator follow-up: the Lumos-T Safely Reduces Routine Office Device Follow-up (TRUST) trial. *Circulation.* 2010;122:325–32.
60. Ricci RP, Morichelli L, Santini M. Remote control of implanted devices through Home Monitoring technology improves detection and clinical management of atrial fibrillation. *Europace.* 2009;11:54–61.
61. Marzegalli M, Lunati M, Landolina M, Perego GB, Ricci RP, Guenzati G, et al. Remote monitoring of CRT-ICD: the multicenter Italian CareLink evaluation—ease of use, acceptance, and organizational implications. *Pacing Clin Electrophysiol.* 2008;31:1259–64.
62. Raatikainen MJ, Uusimaa P, Van Ginneken MM, Janssen JP, Linnaluoto M. Remote monitoring of implantable cardioverter defibrillator patients: a safe, time-saving, and cost-effective means for follow-up. *Europace.* 2008;10:1145–51.