

Update: Acute Heart Failure (I)

Acute Heart Failure: Epidemiology, Risk Factors, and Prevention



Dimitrios Farmakis, John Parissis, John Lekakis, and Gerasimos Filippatos*

Heart Failure Unit, Department of Cardiology, Attikon University Hospital, Athens, Greece

Article history:

Available online 4 February 2015

Keywords:

Acute heart failure
Epidemiology
Risk factors
Prevention

ABSTRACT

Acute heart failure represents the first cause of hospitalization in elderly persons and is the main determinant of the huge healthcare expenditure related to heart failure. Despite therapeutic advances, the prognosis of acute heart failure is poor, with in-hospital mortality ranging from 4% to 7%, 60- to 90-day mortality ranging from 7% to 11%, and 60- to 90-day rehospitalization from 25% to 30%. Several factors including cardiovascular and noncardiovascular conditions as well as patient-related and iatrogenic factors may precipitate the rapid development or deterioration of signs and symptoms of heart failure, thus leading to an acute heart failure episode that usually requires patient hospitalization. The primary prevention of acute heart failure mainly concerns the prevention, early diagnosis, and treatment of cardiovascular risk factors and heart disease, including coronary artery disease, while the secondary prevention of a new episode of decompensation requires the optimization of heart failure therapy, patient education, and the development of an effective transition and follow-up plan.

© 2014 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Insuficiencia cardiaca aguda: epidemiología, factores de riesgo y prevención

RESUMEN

La insuficiencia cardiaca aguda constituye la primera causa de hospitalización en las personas ancianas y es el principal factor determinante del enorme gasto de asistencia sanitaria asociado a la insuficiencia cardiaca. A pesar de los avances terapéuticos realizados, la insuficiencia cardiaca aguda tiene un mal pronóstico, con una mortalidad hospitalaria que oscila entre el 4 y el 7%, una mortalidad a los 60 a 90 días de entre el 7 y el 11% y una tasa de rehospitalizaciones a los 60 a 90 días que va del 25 al 30%. Hay varios factores, entre los que se encuentran los trastornos cardiovasculares y no cardiovasculares, así como factores relacionados con el paciente y factores iatrogénicos, que pueden desencadenar una progresión rápida o un agravamiento de los signos y síntomas de insuficiencia cardiaca, lo que conduce a un episodio de insuficiencia cardiaca aguda que suele requerir el ingreso hospitalario del paciente. La prevención primaria de la insuficiencia cardiaca aguda se centra principalmente en la prevención, el diagnóstico precoz y el tratamiento de los factores de riesgo cardiovascular y la cardiopatía, incluida la enfermedad coronaria, mientras que la prevención secundaria para evitar nuevos episodios de descompensación requiere la optimización del tratamiento de la insuficiencia cardiaca, la educación sanitaria del paciente y el desarrollo de una transición y un plan de seguimiento efectivos.

© 2014 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Insuficiencia cardiaca aguda
Epidemiología
Factores de riesgo
Prevención

Abbreviations

AHF: acute heart failure

INTRODUCTION

Acute heart failure (AHF) is the rapid development or change of signs and symptoms of heart failure that requires medical attention and usually leads to patient hospitalization.^{1–3} Acute heart failure represents the first cause of hospital admission in elderly persons in the western world and, despite advances in medical and device therapy, it still has unacceptably high morbidity and mortality rates. As a result, AHF represents a major public health issue, an enormous financial burden, and a challenge for current cardiovascular research.³

* Corresponding author: Heart Failure Unit, Athens University Hospital, 12461 Athens, Greece.

E-mail address: geros@otenet.gr (G. Filippatos).

EPIDEMIOLOGY

A number of large-scale registries, such as the ADHERE^{4–6} and OPTIMIZE-HF performed in the United States,^{4–7} the EHFS I and II^{8–10} and the ESC-HF Pilot registry performed in Europe,^{8–11} as well as the international ALARM-HF¹² have provided us with some epidemiological evidence on AHF.

Patients admitted for AHF are aged > 70 years and about half of them are male. Most have a previous history a heart failure, while *de novo* AHF represents only one-fourth to one-third of cases. About 40% to 55% have preserved LVEF (left ventricular ejection fraction). Those patients have a constellation of cardiovascular and noncardiovascular abnormalities. Concerning cardiovascular comorbid conditions, most AHF patients have a history of arterial hypertension, about half have coronary artery disease, and one-third or more have atrial fibrillation. In terms of noncardiovascular comorbidities, about 40% of patients admitted for AHF have a history of diabetes mellitus, about one-fourth to one-third have renal dysfunction and chronic obstructive pulmonary disease, while anemia is also present in 15% to 30% of patients. The main clinical features of AHF patients according to the aforementioned registries are outlined in Table 1.

In most of the published AHF registries, in-hospital mortality ranges from 4% to 7% (Table 2), with the exception of ALARM-HF, in which mortality was as high as 11%, apparently due to the relatively higher proportion of patients with cardiogenic shock (around 12% vs < 4% in the rest of the aforementioned registries). The median length of hospital stay ranged from 4 days to 11 days. Postdischarge mortality up to 3 months was 7% to 11%, while 1-year postdischarge mortality reported by the ADHERE registry was 36%.⁵ Heart failure progression itself represents the cause of death in less than half of patients. According to data from the EVEREST trial,¹³ 41% of AHF patients die of heart failure deterioration, 26% die suddenly, and 13% die of noncardiovascular comorbidities. It should be stressed that, although in-hospital mortality tends to be higher in patients with reduced LEVF compared with those with preserved LVEF, postdischarge morbidity is similar in the 2 groups.¹⁴

Postdischarge rehospitalization rates are quite high, as about one-fourth of patients are readmitted within 3 months, while two-thirds of them are rehospitalized within a year. It has been shown that the readmission rate follows a biphasic course consisting of 2 peaks, an early one during the first 2 to 3 months postdischarge and a late one during the final stage of the syndrome, separated by a long plateau phase with low admission rates.¹⁵

Table 1
Clinical Characteristic of Acute Heart Failure Patients in Different Registries

	ADHERE	OPTIMIZE-HF	E HFS I	EHFS II	ESC-HF Pilot (AHF arm)	ALARM-HF
Patients, No.	105 388	48 612	11 327	3580	1892	4953
Age, mean (SD), y	72.0 (14.0)	73.1 (14.2)	71	69.9 (12.5)	70.0 (13.0)	66–70*
Gender, male, %	48	48	53	61	63	62
History of heart failure, %	75	87	65	63	75	64
Arterial hypertension, %	72.0	71.0	53.0	62.5	61.8	70.2
Coronary artery disease, %	57.0	50.0	68.0	53.6	50.7	30.7
Diabetes mellitus, %	44.0	42.0	27.0	32.8	35.1	45.3
Atrial fibrillation, %	31.0	31.0	43.0	38.7	43.7	24.4
Renal dysfunction, %	30.0	30.0	17.0	16.8	26.0	21.4
COPD, %	31.0	28.0		19.3		24.8
Anemia, %				14.7	31.4	14.4

ADHERE, Acute Decompensated Heart Failure National Registry; AHF, acute heart failure; ALARM-HF, Acute Heart Failure Global Survey of Standard Treatment; COPD, chronic obstructive pulmonary disease; EHFS, EuroHeart Failure Survey; ESC-HF Pilot, European Society of Cardiology-Heart Failure Pilot registry; OPTIMIZE-HF, Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure; SD, standard deviation.

* Median.

Table 2
Acute Heart Failure Outcome in Different Registries

	ADHERE	OPTIMIZE-HF	EHFS I	EHFS II	ESC-HF Pilot (AHF arm)	ALARM-HF
Patients, No.	105 388	48 612	11 327	3580	1892	4953
In-hospital mortality, %	4.0	4.0	6.9	6.7	3.8	11.0
Hospital stay, median, days	4	4	11	9	8	6
30-90-days mortality, %	11.2 (30 days)	9.0 (60-90 days)	6.6 (90 days)			
1-year mortality, %	36					
Readmission (time period),	22.1 (30 days) 65.8 (1 year)	30.0 (60-90 days)	24.0 (90 days)			

ADHERE, Acute Decompensated Heart Failure National Registry; AHF, acute heart failure; ALARM-HF, Acute Heart Failure Global Survey of Standard Treatment; EHFS, EuroHeart Failure Survey; ESC-HF Pilot, European Society of Cardiology-Heart Failure Pilot registry; OPTIMIZE-HF, Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure.

Table 3
Causes and Precipitating Factors of Acute Heart Failure

Cardiovascular	Non cardiovascular	Patient-related or iatrogenic
<ul style="list-style-type: none"> • Acute coronary syndromes • Tachycardias (ie, atrial fibrillation) • Bradycardias (ie, third degree atrioventricular block) • Uncontrolled hypertension or hypertensive crisis • Myocarditis • Acute pulmonary embolism • Acute valvular regurgitation (i.e., endocarditis, myocardial infarction) • Aortic dissection • Cardiac tamponade 	<ul style="list-style-type: none"> • Infections and febrile states • COPD exacerbation or asthma • Renal dysfunction • Anemia • Hyperthyroidism • Hypothyroidism • Strenuous exercise • Emotional stress • Pregnancy (peripartum cardiomyopathy) 	<ul style="list-style-type: none"> • Poor compliance with medication • Increased salt or fluid intake • Surgery • Drugs (ie, NSAID, thiazolidinediones) • Alcohol abuse

COPD, chronic obstructive pulmonary disease; NSAID, non-steroid anti-inflammatory drugs.

RISK FACTORS

Several cardiovascular and noncardiovascular conditions may cause a rapid development or deterioration of signs and symptoms of heart failure leading to hospitalization. A detailed list of the causes and precipitating factors leading to AHF is provided in Table 3. Heart failure accounts for less than half of readmission causes. More specifically, according to the EVEREST trial, 46% of patients are admitted because of heart failure, while 39% are hospitalized because of noncardiovascular comorbid conditions.¹³

A number of clinical and laboratory parameters have been shown to predict AHF hospitalization (Table 4). Gradually aggravating symptoms and signs of congestion, deterioration of biomarkers such as natriuretic peptides or renal function parameters, increasing need for diuretics or intolerance to heart failure medication are the hallmarks of deterioration and thus of an upcoming AHF episode.¹⁵

PREVENTION

Primary prevention of AHF concerns the prevention and early diagnosis and treatment of the causes of heart failure (Table 3) and mainly of cardiovascular risk factors and heart disease. Coronary artery disease is the cause in two-thirds of heart failure patients, particularly those with reduced LVEF, while arterial hypertension is found in approximately 70% of patients, particularly those with

preserved LVEF.³ Therefore, timely and effective treatment of risk factors for atherosclerosis, coronary artery disease, and arterial hypertension is expected to limit the occurrence of heart failure.

The secondary prevention of AHF concerns the prevention of chronic heart failure decompensation that leads to AHF episodes requiring hospitalization. As each hospitalization for AHF is known to confer additional deterioration of cardiac and kidney function, recurrent AHF episodes lead to a gradual decline of the patient's clinical course.¹⁶ As a result, the greater the number of hospital admissions, the worse patient survival.¹⁷ Furthermore, AHF hospitalization represents approximately 70% of the total expenditure on heart failure.¹⁸ Therefore, the secondary prevention of AHF episodes represents an important goal both in medical and socioeconomic terms.

As previously stated, a significant proportion of readmissions occur during the first weeks postdischarge.¹⁵ It has been postulated that early readmissions are preventable in up to 75% of patients and that the main reasons for early readmissions are incomplete decongestion during the preceding hospital stay and a poor transition plan.¹⁹ Therefore, the strategies to prevent readmission mainly concern the optimization of in-hospital management of patients and the development of the patient transition and follow-up plan. These strategies are outlined in Table 5. It should be stressed that patient admission may also provide a chance for implementation or the titration of chronic heart failure therapy, better education, patient training, and the development of a follow-up plan.²⁰

Table 4
Predictors of Postdischarge Rehospitalization for Acute Heart Failure

Predictor type	Examples
Symptoms	Increasing body weight, persisting peripheral edema, dyspnea aggravation
Clinical signs	Increased jugular venous pressure, orthopnea
Comorbid conditions	Chronic renal disease, diabetes mellitus, COPD, anemia
Functional status	Quality of life
Biomarkers	Natriuretic peptides, cardiac troponins, serum sodium, serum creatinine
Echocardiography	Left ventricular filling pattern
Treatment	Increase in diuretics, intolerance to disease-modifying therapy with hypotension or renal impairment
Psychosocial and socioeconomic factors	Living alone, low income

COPD, chronic obstructive pulmonary disease.

Table 5
Strategies to Prevent Postdischarge Rehospitalization for Acute Heart Failure

In-hospital treatment	<ul style="list-style-type: none"> • Decongestion • Identification and treatment of heart failure cause • Treatment and prevention of exacerbating factors • Proper titration of chronic heart failure therapy
Planning and transition	<ul style="list-style-type: none"> • Establishment of specific follow-up plan • Early postdischarge visit (7–10 days) • Collaboration with primary care physician • Patient education and training • Nurse home visits • Telemonitoring

CONFLICT OF INTERESTS

G. Filippatos is a member of steering committees of acute heart failure trials sponsored by Novartis, Cardioentis, the European Union and Bayer. J. Parissis has received fees for conference presentations from Novartis International.

REFERENCES

- McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC Committee for Practice Guidelines. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2012;33:1787–847.
- Gheorghiade M, Filippatos G, Felker M. Diagnosis and management of acute heart failure syndromes. In: Braunwald's heart disease: a textbook of cardiovascular medicine. 9th ed. Philadelphia: Elsevier, Saunders; 2012.
- Farmakis D, Parissis J, Filippatos G. Acute heart failure: epidemiology, classification and pathophysiology. In: ESC Textbook of Intensive and Acute Cardiac Care. 2nd ed. Oxford: Oxford University Press; 2015.
- Adams Jr KF, Fonarow GC, Emerman CL, LeJemtel TH, Costanzo MR, Abraham WT, et al.; ADHERE Scientific Advisory Committee and Investigators. Characteristics and outcomes of patients hospitalized for heart failure in the United States: rationale, design, and preliminary observations from the first 100,000 cases in the Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J*. 2005;149:209–16.
- Kociol RD, Hammill BG, Fonarow GC, Klaskala W, Mills RM, Hernandez AF, et al. Generalizability and longitudinal outcomes of a national heart failure clinical registry: comparison of Acute Decompensated Heart Failure National Registry (ADHERE) and non-ADHERE Medicare beneficiaries. *Am Heart J*. 2010;160:885–92.
- Gheorghiade M, Filippatos G. Reassessing treatment of acute heart failure syndromes: the ADHERE Registry. *Eur Heart J Suppl*. 2005;7 Suppl B:B13–9.
- Fonarow GC, Abraham WT, Albert N, Gattis W, Gheorghiade M, Greenberg B, et al. Impact of evidence-based heart failure therapy use at hospital discharge on treatment rates during follow-up: a report from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure (OPTIMIZE-HF). *J Am Coll Cardiol*. 2005;45:345A.
- Cleland JG, Swedberg K, Follath F, Komajda M, Cohen-Solal A, Aguilar JC, et al.; Study Group on Diagnosis of the Working Group on Heart Failure of the European Society of Cardiology. The EuroHeart Failure Survey programme—a survey on the quality of care among patients with heart failure in Europe. Part 1: patient characteristics and diagnosis. *Eur Heart J*. 2003;24:442–63.
- Komajda M, Follath F, Swedberg K, Cleland J, Aguilar JC, Cohen-Solal A, et al.; Study Group on Diagnosis of the Working Group on Heart Failure of the European Society of Cardiology. The EuroHeart Failure Survey programme—a survey on the quality of care among patients with heart failure in Europe. Part 2: treatment. *Eur Heart J*. 2003;24:464–74.
- Nieminen MS, Brutsaert D, Dickstein K, Drexler H, Follath F, Harjola VP, et al.; EuroHeart Survey Investigators; Heart Failure Association, European Society of Cardiology. EuroHeart Failure Survey II (EHFS II): a survey on hospitalized acute heart failure patients: description of population. *Eur Heart J*. 2006;27:2725–36.
- Maggioni AP, Dahlstrom U, Filippatos G, Chioncel O, Leiro MC, Drozd J, et al.; Heart Failure Association of ESC (HFA). EURObservational Research Programme: the Heart Failure Pilot Survey (ESC-HF Pilot). *Eur J Heart Fail*. 2010;12:1076–84.
- Follath F, Yilmaz MB, Delgado JF, Parissis JT, Porcher R, Gayat E, et al. Clinical presentation, management and outcomes in the Acute Heart Failure Global Survey of Standard Treatment (ALARM-HF). *Intensive Care Med*. 2011;37:619–26.
- O'Connor CM, Miller AB, Blair JE, Konstam MA, Wedge P, Bahit MC, et al.; Efficacy of Vasopressin Antagonism in heart Failure Outcome Study with Tolvaptan (EVEREST) investigators. Causes of death and rehospitalization in patients hospitalized with worsening heart failure and reduced left ventricular ejection fraction: results from Efficacy of Vasopressin Antagonism in Heart Failure Outcome Study with Tolvaptan (EVEREST) program. *Am Heart J*. 2010;159:841–9.
- Fonarow GC, Stough WG, Abraham WT, Albert NM, Gheorghiade M, Greenberg BH, et al. Characteristics, treatments, and outcomes of patients with preserved systolic function hospitalized for heart failure: a report from the OPTIMIZE-HF Registry. *J Am Coll Cardiol*. 2007;50:768–77.
- Desai AS, Stevenson LW. Rehospitalization for heart failure: predict or prevent? *Circulation*. 2012;126:501–6.
- Filippatos G, Zannad F. An introduction to acute heart failure syndromes: definition and classification. *Heart Fail Rev*. 2007;12:87–90.
- Stehlik J, Edwards LB, Kucheryavaya AY, Aurora P, Christie JD, Kirk R, et al. The Registry of the International Society for Heart and Lung Transplantation: twenty-seventh official adult heart transplant report—2010. *J Heart Lung Transplant*. 2010;29:1089–103.
- Stewart S, Jenkins A, Buchan S, McGuire A, Capewell S, McMurray JJ. The current cost of heart failure to the National Health Service in the UK. *Eur J Heart Fail*. 2002;4:361–71.
- Van Walraven C, Bennett C, Jennings A, Austin PC, Forster AJ. Proportion of hospital readmissions deemed avoidable: a systematic review. *CMAJ*. 2011;183:E391–402.
- Gheorghiade M, Vaduganathan M, Fonarow GC, Bonow RO. Rehospitalization for heart failure: problems and perspectives. *J Am Coll Cardiol*. 2013;61:391–403.