

Review article

Adipose-derived Mesenchymal Stem Cells and Their Reparative Potential in Ischemic Heart Disease

Lina Badimon,^{a,b,*} Blanca Oñate,^a and Gemma Vilahur^a^a Centro de Investigación Cardiovascular, CSIC-ICCC, Hospital de la Santa Creu i Sant Pau e IIB-Sant Pau, Barcelona, Spain^b Cátedra de Investigación Cardiovascular, UAB-HSCSP-Fundación Jesús Serra, Barcelona, Spain

Article history:

Available online 29 May 2015

Keywords:

Adipose tissue
Adipose-derived stem cells
Regenerative medicine
Ischemic heart disease
Cardiovascular risk factors

ABSTRACT

Adipose tissue has long been considered an energy storage and endocrine organ; however, in recent decades, this tissue has also been considered an abundant source of mesenchymal cells. Adipose-derived stem cells are easily obtained, show a strong capacity for *ex vivo* expansion and differentiation to other cell types, release a large variety of angiogenic factors, and have immunomodulatory properties. Thus, adipose tissue is currently the focus of considerable interest in the field of regenerative medicine. In the context of coronary heart disease, numerous experimental studies have supported the safety and efficacy of adipose-derived stem cells in the setting of myocardial infarction. These results have encouraged the clinical use of these stem cells, possibly prematurely. Indeed, the presence of cardiovascular risk factors, such as hypertension, coronary disease, diabetes mellitus, and obesity, alter and reduce the functionality of adipose-derived stem cells, putting in doubt the efficacy of their autologous implantation. In the present article, white adipose tissue is described, the stem cells found in this tissue are characterized, and the use of these cells is discussed according to the preclinical and clinical trials performed so far.

© 2015 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Células madre mesenquimales derivadas de tejido adiposo y su potencial reparador en la enfermedad isquémica coronaria

RESUMEN

Se ha considerado al tejido adiposo como de almacenamiento energético y como un órgano endocrino; sin embargo, en las últimas décadas se lo ha considerado como una fuente abundante de células mesenquimales. Las células madre derivadas del tejido adiposo son de fácil obtención, presentan una gran capacidad de expansión *ex vivo* y gran plasticidad a otros tipos celulares, liberan gran variedad de factores angiogénicos y presentan propiedades inmunomoduladoras. Por ello, actualmente constituyen un foco de gran interés en la medicina regenerativa. En el contexto de enfermedad cardíaca coronaria, múltiples estudios experimentales han avalado la seguridad y la eficacia del uso de las células madre derivadas del tejido adiposo en el contexto de infarto de miocardio. Todo ello ha promovido, quizá precozmente, su uso clínico. De hecho, se ha demostrado que la presencia de factores de riesgo cardiovascular como hipertensión, enfermedad coronaria, diabetes mellitus u obesidad, altera y merma la funcionalidad de las células madre derivadas del tejido adiposo, lo que deja en entredicho la eficacia basada en el implante de células madre derivadas del tejido adiposo autólogo. En el siguiente artículo se describe el tejido adiposo blanco, se caracterizan las células madre que lo componen y se discute sobre su uso según los estudios preclínicos y clínicos realizados hasta el momento.

© 2015 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Tejido adiposo
Células madre derivadas del tejido adiposo
Medicina regenerativa
Enfermedad isquémica coronaria
Factores de riesgo cardiovascular

ADIPOSE TISSUE

Adipose tissue is one of the most abundant human tissues. It constitutes between 15% and 20% of the body weight of men and

between 20% to 25% of that of women and is widely distributed throughout various body regions. This specialized tissue is of mesenchymal origin, consisting of a combination of white adipose tissue (WAT) and brown adipose tissue, with each tissue type showing distinct functions, morphologies, and distributions. In both tissues, the predominant cell is the adipocyte, comprising between one- and two-thirds of the total, and the remaining tissue is composed of different types of cells constituting the stromal vascular fraction (SVF).

* Corresponding author: Centro de Investigación Cardiovascular, Sant Antoni M. Claret 167, 08025 Barcelona, Spain.

E-mail address: lbadimon@csic-iccc.org (L. Badimon).

Abbreviations

ADSCs: adipose-derived stem cells
 MSCs: mesenchymal stem cells
 SVF: stromal vascular fraction
 WAT: white adipose tissue

White Adipose Tissue

Although WAT is distributed throughout the body, its principal deposits are subcutaneous, where it acts as an energy storage system, and in the visceral or intra-abdominal region, where it protects against possible trauma. The 2 tissues show different adipokine expression profiles,¹ metabolic functions,² vascular density, and innervation. Visceral adipose tissue shows a higher angiogenic potential and more acute inflammatory profile than subcutaneous tissue.³ The accumulation of subcutaneous adipose tissue represents a physiological response to situations of excessive intake and low energy expenditure (physical inactivity), acting as an “energy sink”. Individuals with peripheral obesity (subcutaneous distribution) do not show the characteristic medical complications of obesity. In contrast, increased visceral adipose tissue (central obesity) is associated with a state of hyperglycemia, hyperinsulinemia, hypertriglyceridemia, hypercholesterolemia, reduced circulating levels of high-density lipoproteins, decreased glucose tolerance, increased apolipoprotein B-rich lipoproteins, and hepatic steatosis. All of these conditions are characteristics of insulin resistance syndrome, which increases the risk of the development of type 2 diabetes mellitus.⁴ Currently, waist size is an important diagnostic component of metabolic syndrome and has been identified as an independent risk factor for other diseases, such as cardiovascular diseases, stroke, hypertension, and nonalcoholic fatty liver disease.^{5–7}

The main function of WAT is to regulate the energy homeostasis of the body, which is under the control of the nervous and endocrine systems. In times of caloric excess, adipose tissue stores fat in the form of triglycerides. These lipids are then released into the blood in times of energy demand to be used as an energy source in other tissues, such as the liver, kidneys, skeletal muscle, and myocardium.⁸ However, WAT is currently considered a multifunctional organ because, besides its energy function, it acts as an endocrine organ and as a reservoir of mesenchymal stem cells (MSCs).

Composition of White Adipose Tissue

White adipose tissue consists of mature adipocytes and intercellular tissue or SVF. Adipocytes, the most abundant cells in adipose tissue, contain a single large cytoplasmic vacuole that mainly stores triglycerides and cholesterol esters. Depending on nutritional status, adipocytes can alter their size to between 25 μm and 200 μm . Adipocytes contain the machinery necessary for lipid metabolism (Figure 1).⁹ These pathways can be altered by an increase in weight, triggering insulin resistance syndrome. Indeed, fatty acids not only show an energy function, but also act as regulatory signals of the gene expression of proteins involved in lipid metabolism,¹⁰ favor a prothrombotic state, and are associated with inflammatory processes.¹¹ Thus, an excess of circulating fatty acids (lipotoxicity) is one of the strongest links between obesity and the development of metabolic syndrome and/or cardiovascular disease. When calorie consumption exceeds energy expenditure, a metabolic state develops that promotes adipocyte

hypertrophy (increased size) and hyperplasia (increased number).¹² The latter involves mobilization of stem cells toward the adipocyte lineage (adipogenesis). New or small adipocytes are more sensitive to insulin and show a marked capacity for the uptake of free fatty acids and triglycerides in the postprandial period.⁴ As adipocytes increase in size (hypertrophy), they become dysfunctional, losing their ability to protect against systemic lipotoxicity, and ectopic fat begins to accumulate. These distended adipocytes become hyperlipolytic and resistant to insulin and its antilipolytic signals. Another important function performed by adipocytes is that of endocrine cells, as discussed below.

The other component of WAT is the SVF. Although the cells constituting the SVF remain to be fully defined, they include adipocyte precursors and vascular and blood cells.¹³ Pericytes form the vasculature of adipose tissue, together with endothelial and smooth muscle cells. The extent and characteristics of this capillary network are crucial for processes such as the growth, function, and development of adipose tissue.¹⁴ Adipocytes and other cells of the SVF secrete proangiogenic factors, ensuring that the tissue has a generous blood supply. In addition, adipose tissue, via its resident immune system cells, strongly controls the metabolism of the body. In nonobese individuals, these cells are involved in elimination of necrotic adipocytes, remodeling of the extracellular matrix, angiogenesis, adipogenesis, and maintaining insulin sensitivity. However, in obese individuals, the number of immune system cells increases; these cells acquire a proinflammatory phenotype and release a huge number of cytokines in charge of recruiting and activating other immune system cells and inducing insulin resistance syndrome in adipose tissue.¹⁵ Macrophages are the cells with the most important role in the acquisition of the low-grade chronic proinflammatory state that characterizes obesity. During adipose tissue expansion, there is a greater recruitment of M1 macrophages (a proinflammatory phenotype). These macrophages secrete most of the proinflammatory cytokines found in obese adipose tissue,¹⁶ whereas the resident M2 macrophages show an anti-inflammatory phenotype.¹⁷ Finally, in the SVF, there are adipose-derived stem cells (ADSCs) and preadipocytes. These cell populations are in charge of maintaining adipocyte population renewal in physiological conditions and play a vital role in the obesity-related expansion of adipose tissue. The differences between these 2 cell groups are poorly defined. Both ADSCs and preadipocytes show a similar morphology. However, whereas ADSCs can differentiate into other lineages and show a large capacity for self-renewal, preadipocytes have lost these differentiation capabilities and can only generate mature adipocytes.¹⁸

Factors Secreted by White Adipose Tissue

The WAT secretes a multitude of bioactive peptides, known under the umbrella term of adipocytokines or adipokines (Table 1).^{19,20} However, many of these factors are not only secreted by adipocytes, but also by the cells constituting the SVF, such as macrophages and ADSCs. Through these secreted factors, adipose tissue participates in the autocrine and paracrine regulation of adipose tissue itself, as well as affecting the function of other organs. In addition, adipose tissue is in charge of regulating energy homeostasis and body weight, insulin sensitivity, and various functions of the immune, vascular, and reproductive systems.²⁰ This endocrine function of adipose tissue explains the pathophysiological relationship between excess body fat and its associated pathological states, because obesity and/or metabolic syndrome trigger a dysregulation of the secreted amounts of these molecules.²¹




Figure 1. Diagram illustrating the lipogenesis and lipolysis processes occurring in mature adipocytes. After eating and an increase in blood insulin, lipogenesis is activated in adipocytes. In this process, adipocytes, via lipoprotein lipase, degrade the triglycerides of chylomicrons and of very-low-density lipoprotein to fatty acids. These molecules enter the adipocyte to be esterified with glycerol-3 phosphate and synthesize the triglycerides that are stored in the lipid vacuole. In the adipocyte, insulin not only stimulates lipoprotein lipase synthesis, but also stimulates uptake and metabolism of glucose to glycerol-3 phosphate. In contrast, during lipolysis, the triglycerides stored are mobilized to produce free fatty acids and glycerol to meet the energy requirements of the body. Catabolic hormones, secreted in response to a low blood concentration of glucose, activate the synthesis and movement of hormone-sensitive lipase from the cytosol to the surface of the lipid vacuole, where hormone-sensitive lipase hydrolyzes triglycerides. The fatty acids produced are secreted as free fatty acids to the circulation, where they will be transported by albumin to the target organs to be oxidized to produce energy. Similarly, the glycerol derived from lipolysis is also released into the circulation to be used by the liver as a source of carbon. AQP7, aquaporin-7; CM, chylomicrons; FABP, fatty acid-binding protein; FAT/CD36, fatty acid translocase; FATP, fatty acid transport protein; GLUT4, glucose transporter type 4; HSL, hormone-sensitive lipase; LPL, lipoprotein lipase; TG, triglycerides; VLDL, very-low-density lipoprotein.

ADIPOSE-DERIVED STEM CELLS

For many years, the hyperplastic growth of adipose tissue was believed to be due to the existence of a unipotent progenitor cell population, the preadipocytes. However, in 2001, Zuk et al²² identified the existence of MSCs with self-renewal and multipotent capacities in adipose tissue. Since then, adipose tissue has been considered a source of MSCs for use in cell therapy.²²

Origin of Adipose-derived Stem Cells

Since adipocytes and their progenitors were found to originate from MSCs,²³ it has been noted that ADSCs could be derived from mesenchymal lineage cells from the bone marrow. Indeed, the cells of the SVF show various similarities with those of the bone marrow. Both stromae contain a heterogeneous population of MSCs with the ability to differentiate into various lineages (adipocytic, chondrocytic, and myogenic) according to culture conditions.²⁴ Mansilla et al²⁵ noted that the bone marrow is the principal producer of the MSCs that supply the populations of MSCs found in other peripheral organs (peripheral reservoirs). Additionally, these authors showed that the cells are maintained in a quiescent and undifferentiated state until they are “called” to proliferate and move to the required tissues.

Indeed, although there are practically no MSCs in the circulation of healthy individuals, these stem cells are mobilized toward damaged regions to participate in tissue repair and regeneration.²⁶ Thus, it can be inferred that obese adipose tissue, as an important source of chemotactic factors, would act as a niche where circulating MSCs could home to and differentiate into adipocytes.²⁵

Characteristics of Adipose-derived Stem Cells

Adipose-derived stem cells show the typical characteristics of MSCs proposed by the Mesenchymal and Tissue Stem Cell Committee of the International Society for Cellular Therapy:²⁷

- They must adhere to plastic while maintained in standard culture conditions.
- They must be able to differentiate into osteogenic, adipogenic, and chondrogenic lineages (Figure 2).
- They must express the surface markers CD105, CD73, and CD90 and not express CD45, CD34, CD14 or CD11b, CD79a, or CD19, or HLA-II surface molecules.

Although ADSCs do not express a single surface marker that enables their identification, they express the characteristic

Table 1
Factors Secreted by Adipose Tissue

Adipokines	Function	Secreting cell	Regulation
11 β -hydroxysteroid dehydrogenase type 1	Steroid metabolism	Adipocytes, preadipocytes	↑ obesity
Free fatty acids	Lipid metabolism	Adipocytes	↑ obesity
Adiponectin	Increases insulin sensitivity, inflammation, and arteriosclerosis	Adipocytes	↓ obesity
Adipsin and acylation stimulating protein (ASP)	Stress and immune response	Adipocytes, M2 macrophages	↑ obesity
Angiotensinogen	Vascular homeostasis	Adipocytes, SVF	↑ obesity
Apelin	IR	Adipocytes, SVF, macrophages	↑ obesity
Aromatase	Lipid metabolism	Adipocytes, ADSCs, macrophages	↑ obesity
IGF-1	Lipid metabolism and IR	Adipocytes, preadipocytes, ADSCs	
TNF α	Inflammation, arteriosclerosis, and IR	Adipocytes, M1 macrophages	↑ obesity
Macrophage migration inhibitory factor (MIF)	Inflammation	Adipocytes, ADSCs, immune system cells	↑ obesity
TGF β	Cell adhesion and migration, growth and differentiation	Adipocytes, SVF, ADSCs	↑ obesity
Steroid hormones	Lipid metabolism and IR	Adipocytes, preadipocytes	↑ obesity
PAI-1	Vascular homeostasis	Adipocytes, SVF	↑ obesity
IL-1	Inflammation and IR	M1 macrophages	↑ obesity
IL-6	Inflammation, arteriosclerosis, and IR	Adipocytes, SVF	↑ obesity
IL-8	Pro-atherogenesis	Adipocytes, SVF	↑ obesity
IL-10	Inflammation and IR	Adipocytes, M2 macrophages	↑ obesity, ↓ MS
Leptin	Dietary intake, reproduction, angiogenesis, and immune system	Adipocytes	↑ obesity
Hormone-sensitive lipase	Lipid metabolism	Adipocytes	↓ obesity
Lipoprotein lipase	Lipid metabolism	Adipocytes	↑ obesity
Metallothionein	Stress and immune response	Adipocytes, SVF	↑ obesity
Monobutyrin	Angiogenesis	Adipocytes	↑ obesity
Omentin	IR	SVF, macrophages	↓ obesity
Perilipin	Lipid metabolism	Adipocytes	↑ obesity
Prostaglandins (PGE ₂ , prostacyclin, PG ₂ F α)	Blood flow, lipolysis, cellular differentiation	Adipocytes, ADSCs	↑ obesity
C-reactive protein	Inflammation, arteriosclerosis, and IR	SVF	↑ obesity
Fatty acid-binding protein (FABP4/aP2)	Lipid metabolism	Adipocytes, macrophages	↑ obesity
MCP-1	Pro-atherogenesis and IR	Adipocytes, M1 macrophages	↑ obesity
CETP	Lipid metabolism	Preadipocytes, adipocytes	↑ obesity
RBP	Lipid metabolism	Adipocytes	Variable in obesity
Resistin	Inflammation and IR	Adipocytes, M2 macrophages	Variable in obesity
Thrombospondin	Angiogenesis	Adipocytes	↑ obesity
Visfatin	IR	Adipocytes, preadipocytes, neutrophils	Variable in obesity
Zinc- α 2-glycoprotein	Lipid metabolism, cancer, and cachexia	Adipocytes, SVF	↓ obesity

ADSCs, adipose-derived stem cells; CETP, cholesteryl ester transfer protein; IGF-1, insulin-like growth factor 1; IR, insulin resistance; IL, interleukin; MCP-1, monocyte chemoattractant protein-1; MS, metabolic syndrome; RBP, retinol-binding protein; PAI-1, plasminogen activator inhibitor-1; SVF, stromal vascular fraction; TGF β , transforming growth factor beta; TNF α , tumor necrosis factor alpha.

Adapted from Ronti et al.²⁰

markers of MSCs in conjunction with some markers expressed by nonprogenitor lines (Table 2).

As metabolically active cells, ADSCs play important roles in the revascularization of damaged tissue, apoptosis inhibition, and immunomodulation. These stem cells secrete a large quantity of extracellular matrix factors and a large number of cytokines and growth, angiogenic, and antiapoptotic factors.²⁸ Indeed, a large part of the beneficial effects of cell therapy with ADSCs is believed to be due to their robust secretion of paracrine factors. Importantly, these

angiogenic and antiapoptotic factors are secreted in bioactive quantities, and this secretion is increased under hypoxia.²⁹

Source-related Differences in Adipose-derived Stem Cells

The metabolic differences between subcutaneous and visceral adipose tissue may be due to the intrinsic characteristics of the cells resident in each tissue, including ADSCs. Indeed, adipocytes




Figure 2. Adipose-derived stem cells differentiation potential. Adipose-derived stem cells are able to differentiate to other types of cells of the same mesodermal lineage (transdifferentiation) or to other types of cells from another lineage (cross-differentiation). ADSCs, adipose-derived stem cells; FABP4, fatty acid-binding protein 4; GFAP, glial fibrillary acidic protein; Ipf-1, insulin promoter factor 1; Isl-1, islet 1; LDL, low-density lipoprotein; LPL, lipoprotein lipase; MAP-2, microtubule-associated protein 2; MyoD1, myogenic differentiation factor 1; NeuN, neuronal nuclear antigen; Ngn-3, neurogenin 3; Nkx2.5, NK2 homeobox 5; OCN, osteocalcin; OPN, osteopontin; Pax-6, paired box protein 6; PECAM-1, platelet endothelial cell adhesion molecule 1; RUNX2, runt-related transcription factor 2; Tie-2, angiopoietin receptor 2; VE, vascular endothelial; VEGFR2, vascular endothelial growth factor 2; vWF, von Willebrand factor.

differentiated *in vitro* from ADSCs derived from the 2 sources show differences inherent to the source tissues.³⁰ These differences are stable and are maintained after the ADSCs have been isolated and cultured *in vitro*.³¹ Various studies have reported differences in the proliferation, differentiation, and apoptotic potentials, as well as gene expression patterns, of ADSCs from different adipose tissues.^{32–34} Adipose-derived stem cells from subcutaneous adipose tissue show greater adipogenic differentiation capacity than ADSCs from visceral adipose tissue.³² The low capacity for differentiation of the visceral ADSCs could partly explain why fat accumulates in already existing adipocytes and, consequently, why the size of their lipid vacuoles increases. In contrast, the greater differentiation capacity of subcutaneous ADSCs would result in lipid accumulation in new adipocytes with smaller vacuoles.³⁵ Accordingly, the size of the lipid vacuoles of visceral adipocytes correlates with the concentrations of circulating lipids, whereas the degree of hyperplasia and size of the subcutaneous adipocytes are more related to the plasma concentrations of glucose and insulin and to insulin sensitivity.³⁶ However, it is still unknown how the ADSCs of each adipose tissue acquire their characteristic phenotypes and at what developmental stage. The regional characteristics of the different ADSCs might be regulated epigenetically, appearing during early developmental phases and being established later by the environment of each adipose tissue and of each individual. Knowledge of the differences between the ADSCs of the different adipose tissues would be of great interest for a better understanding of the biology of the tissue and the development of its different deposits.

Effect of Cardiovascular Risk Factors on Adipose-derived Stem Cells

Various studies have shown that hypercholesterolemia, types 1 and 2 diabetes mellitus, hypertension, and smoking negatively

affect endogenous stem/progenitor cells. Recently, our group has reported that type 2 diabetes mellitus negatively affects the pluripotency and self-renewal capacities of ADSCs, altering the main pathways involved in the maintenance of stem cells and their differentiation and angiogenic potentials.³⁷

Obesity has also been described as a disease that affects ADSCs. Van Harmelen et al³⁸ found that the adipogenic differentiation capacity of ADSCs of subcutaneous mammary adipose tissue is decreased in women with a high body mass index. Subsequently, Nair et al³⁹ reported that ADSCs from subcutaneous adipose tissue of obese Pima Indian individuals had higher expression of proinflammatory genes than those of nonobese individuals. Recently, it has been reported that morbidly obese individuals have ADSCs with impaired proliferation, differentiation, and angiogenic capacities, which negatively affect the regenerative capacity of these cells.⁴⁰ Additionally, the ADSCs of obese patients show lower levels of multipotency markers, increased commitment toward an adipocyte lineage, and higher expression of proinflammatory genes than ADSCs derived from nonobese patients.⁴¹ In addition, the effect of obesity on ADSCs was seen in both those cells derived from subcutaneous adipose tissue and those derived from visceral tissue.⁴²

ADIPOSE-DERIVED STEM CELLS IN CELL THERAPY

Bone marrow-derived MSCs have been used for many years as the main source of stem cells for regenerative medicine and as an alternative to embryonic stem cells.⁴³ However, due to the ease of acquisition and isolation of ADSCs and the large quantity obtained, they have become an important alternative source of stem cells with considerable advantages over bone marrow-derived MSCs.^{44,45} Initially, the reparative/regenerative capacity of ADSCs

Table 2
Characteristic Cell Surface Markers of Adipose-derived Stem Cells

Markers found in ADSCs		Markers not found in ADSCs	
αSMA	α-Smooth muscle actin (ACTA2)	CD104 ^a	β4 integrin
CD10	Neutral endopeptidase (NEP)	CD106 ^a	Component of vascular cell adhesion molecule-1 (VCAM-1)
CD105	Endoglin (SH2)	CD117	c-Kit
CD13	Alanine aminopeptidase	CD11b	αM integrin
CD146	Melanoma cell adhesion molecule (MCAM)	CD11c	αX integrin
CD166	Activated leukocyte cell adhesion molecule (ALCAM)	CD133	Prominin 1
CD24	Heat-stable antigen (HSA)	CD14	
CD29	β1 integrin	CD144	VE-cadherin
CD44	Hyaluronic acid/fibronectin receptor	CD15	Stage-specific embryonic antigen-1 (SSEA-1)
CD49d ^a	α4 integrin	CD16	Fc receptor for IgG
CD49e	α5 integrin	CD19	B lymphocyte surface antigen B4
CD54 ^a	Intracellular adhesion molecule-1 (ICAM-1)	CD3	T cell receptor (TCR)
CD55	Complement decay-accelerating factor (DAF)	CD31	Platelet endothelial cell adhesion molecule 1 (PECAM-1)
CD58		CD33	
CD59	Membrane attack complex inhibition factor (MACIF)	CD34 ^b	
CD71	Transferrin receptor	CD38	
CD73	Ecto-5'-nucleotidase (SH3)	CD4	MHC-II coreceptor
CD9		CD45	Leukocyte common antigen (LCA)
CD90	Thymus cell antigen-1 (Thy-1)	CD56	Neural cell adhesion molecule (NCAM)
HLA-I	Human leukocyte antigen class I (A, B, C)	CD61	β3 integrin
Sca-1	Stem cell antigen 1, ly-6A/E	CD62E	E-selectin
		CD62P	P-selectin
		CD79a	Immunoglobulin-associated α
		CD80	B7.1
		Gly-A	Glycophorin A
		HLA-DR	Human leukocyte antigen class II (DR, DP, DQ)
		Lin ^c	Lineage antigen
		MyD88	Myeloid differentiation primary response gene (88)
		Stro-1 ^a	Stromal cell antigen-1 (at low levels)
		VEGFR2	Vascular endothelial growth factor receptor 2 (Flk-1, KDR)

ADSCs, adipose-derived stem cells; BM-MSCs, bone marrow-derived mesenchymal stem cells.

^a Marker that is oppositely expressed between adipose-derived stem cells and bone marrow-derived mesenchymal stem cells.

^b Marker with controversial ADSC expression results; some authors detect it, whereas others do not.

^c Lin antigens consist of the following group of lineage markers: CD2, CD3, CD4, CD5, CD8, NK1.1, B220, Ter-119, and Gr-1 (granulocyte differentiation antigen 1).

was proposed to be due to their ability to differentiate into other cell types.⁴⁶ However, studies performed in recent years have confirmed that the reparative potential of ADSCs is largely due to their release of paracrine factors.^{44,45}

Adipose-derived Stem Cells in Ischemic Heart Disease

Experimental Studies

In recent years, numerous preclinical studies and some clinical studies have analyzed the safety, behavior, and efficacy of ADSCs in the treatment of ischemic injury, especially that of cardiac origin.^{47–51} The first study was performed in a rat model of heart cryoinjury and involved the injection of recently isolated ADSCs into the left ventricular cavity to simulate intracoronary administration.⁵¹ That study was the first to show that ADSCs home to the myocardium and express specific markers of cardiac cells. Similarly, functional and pathological

analyses of the ADSC-treated animals revealed significantly improved global cardiac function and increased capillary density in the injury border zone compared with controls.⁵¹ Since then, the capacity of ADSCs to generate cardiomyocytes and vascular cells has become a topic of great experimental interest, as shown in Table 3^{51–91} (studies performed in rodents and rabbits) and Table 4^{92–101} (studies performed in porcine models). Notably, controversy surrounds the efficacy of the ADSCs. Whereas some studies have found engrafted ADSCs expressing specific cardiac markers (troponin I and myosin light chain),^{51,102–105} von Willebrand factor, and/or smooth muscle actin, other studies failed to discern the differentiation capacity of ADSCs (Tables 3 and 4).^{52,92} These differences in the *in vivo* differentiation potential of ADSCs could be due to differences in ADSC sources, procurement processes or culture media, animal models, or means of administration, or be due to the limits of histological analysis. Various groups have similarly found a low differentiation capacity of ADSCs in studies *in vivo*.^{52,53,92} All of these observations have led scientists to

Table 3
Experimental Studies Using Rodent and Rabbit Animal Models

Authors	Cell source	Animal model	Lesion	Result
Strem et al ⁵¹	ADSCs from mouse subcutaneous adipose tissue	Mouse	AMI induced by cryolesion	The implanted ADSCs expressed specific markers of cardiomyocytes
Miyahara et al ⁵⁵	Sheets of ADSCs from rat subcutaneous adipose tissue	Rat	AMI induced by LAD ligation	Improved injury and cardiac function
Zhang et al ⁵⁶	Rabbit ADSCs	New Zealand white rabbit	AMI induced by LAD ligation	Improved cardiac function
Mazo et al ⁵⁷	ADSCs from subcutaneous adipose tissue, AD-CMGs, and BM-MNCs from GFP-mice	SD rat	AMI induced by LAD ligation	ADSCs improved cardiac function and tissue viability, increased angiogenesis, and reduced fibrosis
Cai et al ⁵²	ADSCs from human subcutaneous adipose tissue	Rat	AMI induced by LAD ligation	ADSCs improved cardiac function and tissue viability, increased angiogenesis, and reduced fibrosis
Léobon et al ⁵⁸	AD-CMGs from subcutaneous adipose tissue	Mouse	AMI followed by AD-CMG injection	After 4 weeks, the group treated with AD-CMGs showed reduced remodeling and LVEF stability and increased angiogenesis in the peri-infarct zones
Schenke-Layland et al ⁵⁹	ADSCs from rat subcutaneous adipose tissue	Rat	AMI induced by LAD occlusion and reperfusion	Improved cardiac function despite few implanted cells
Van der Bogt et al ⁶⁰	ADSCs from mouse subcutaneous adipose tissue and BM-MSCs	Transgenic FVB mouse	AMI	No improvement detected. Increased apoptosis
Wang et al ⁶¹	ADSCs from rat subcutaneous adipose tissue	Rat	AMI induced by LAD occlusion	After 1 month, improved LVEF, thickening of the cardiac wall, and increased capillary density. Only 0.5% of the ADSCs implanted were positive for specific cardiac cell markers
Zhu et al ⁶²	ADSCs from human subcutaneous adipose tissue overexpressing HGF	SD rat	AMI	ADSCs improved cardiac function and reduced fibrosis
Bai et al ⁵³	ADSCs from human subcutaneous adipose tissue	SCID mouse	AMI	Improved cardiac function, cardiomyogenic differentiation, and increased angiogenesis
Bayes-Genis et al ⁶³	ADSCs from human cardiac adipose tissue	Mouse and rat	AMI	Improved cardiac function, cardiomyogenic differentiation, and increased vasculogenesis
Danoviz et al ⁶⁴	ADSCs from rat subcutaneous adipose tissue. Injected with fibrin α , collagen, or culture medium	Rat	AMI	Inhibition of the negative process of cardiac remodeling
Hwangbo et al ⁶⁵	Human ADSCs	SD rat	AMI induced by permanent ligation of the LAD	After 4 weeks, improved cardiac function and cardiomyogenic differentiation and increased capillary density
Lin et al ⁶⁶	ADSCs from epididymal adipose tissue of rat treated with sildenafil	Lewis rat	Dilated cardiomyopathy	Less apoptosis and fibrosis, improved cardiac function, and increased angiogenesis
Okura et al ⁶⁷	ADSCs from human omental adipose tissue differentiated to cardiomyocytes	Nude rat	AMI	Improved cardiac function and increased cardiomyogenic differentiation
Zhang et al ⁶⁸	ADSCs from the subcutaneous adipose tissue of rats co-injected with fibrin	Rat	AMI	ADSCs + fibrin improved cell implantation, tissue injury, cardiac function, and vascular density
Bai et al ⁶⁹	ADSCs from human subcutaneous adipose tissue	Nude mouse	AMI induced by permanent ligation of the LAD	Implantation of ADSCs; 3.5% of the cells differentiated to cardiomyocytes or endothelial cells
Berardi et al ⁷⁰	Human ADSCs treated with SNAP	Rat	AMI	ADSCs treated with SNAP had improved cardiac function and increased expression of troponin T and von Willebrand factor
Cai et al ⁷¹	ADSCs from rat subcutaneous adipose tissue co-cultured with cardiomyocytes	Rat	AMI	Pretreatment of ADSCs improved their implantation and capacity for repair of cardiac function
Gaebel et al ⁷²	Human ADSCs, BM-MSCs, and MSCs from umbilical cord blood	SCID mouse	LAD ligation	Human MSCs derived from different tissues showed differences in their capacity for repair of cardiac function. CD105+ cells showed better survival in infarcted hearts
Hamdi et al ⁷³	ADSCs from rat subcutaneous adipose tissue in the form of sheets	Rat	Coronary ligation	Rats treated with ADSC sheets survived longer than those that received cell injections. Reduced remodeling of the left ventricle and improved final diastolic volume and cell transplantation

Table 3 (Continued)

Experimental Studies Using Rodent and Rabbit Animal Models

Authors	Cell source	Animal model	Lesion	Result
Li et al ⁵⁴	Human ADSCs	Nude rat	AMI	Improved cardiac function, increased capillary density, no transdifferentiation to cardiac or vascular lineages
Van Dijk et al ⁷⁴	SVF cells and ADSCs from rat subcutaneous adipose tissue	Rat	AMI	SVFs and ADSCs significantly decreased infarct size when they were injected 7 days after the infarct, not the first day
Bagno et al ⁷⁵	ADSCs from rat subcutaneous adipose tissue + matrigel	Wistar rat		After 6 weeks, improved cardiac function and decreased scarring
Beitnes et al ⁷⁶	ADSCs from human subcutaneous adipose tissue and MSCs from human skeletal muscle	Nude rat	AMI	Improved cardiac function and decreased scar size
Fang et al ⁷⁷	Amniotic epithelial cells, MSCs from umbilical cord, and ADSCs from human subcutaneous adipose tissue	Athymic nude rat	AMI	Improved cardiac function and decreased scar size
Hoke et al ⁷⁸	ADSCs from human epicardial adipose tissue treated with phosphodiesterase-5 inhibitor	CD-1 mouse	AMI	Improved cardiac function, increased vascular density, reduced apoptosis, increased secretion of VEGF, FGF-b, and Ang1
Li et al ⁷⁹	Human CDCs, BM-MSCs, ADSCs, and BM-MNCs	Mouse	AMI	After 3 weeks, improved cardiac function, increased engraftment and myogenic lineage differentiation
Liu et al ⁸⁰	ADSCs from rat subcutaneous adipose tissue in chitosan hydrogel	SD rat	AMI induced by LAD ligation	Increased stem cell engraftment, survival, and homing
Paul et al ⁸¹	Human ADSCs injected in genipin cross-linked alginate chitosan microcapsules	Lewis rat	AMI induced by LAD occlusion	Improved retention of implanted cells, decreased infarct zone size, increased vasculogenesis, and improved cardiac function
Paul et al ⁸²	Human ADSCs overexpressing Ang-1	Rat	AMI	Increased cell retention and capillary density, decreased infarct zone, and increased cardiac function
Shi et al ⁸³	ADSCs from rat subcutaneous adipose tissue overexpressing eNOS	Rat	AMI	Decreased infarct zone size
Yang et al ⁸⁴	HO-1-ADSCs from rabbit subcutaneous adipose tissue	Rabbit	AMI	Improved cardiac function, left ventricular size, and cardiomyogenic and angiogenic differentiation
Paul et al ⁸⁵	Human ADSCs and BM-MSCs	Rat	AMI	Improved cardiac function
Wang et al ⁸⁶	Human ADSCs with/without shPHD2 silencing	Mouse	AMI	ADSCs decreased cardiomyocyte apoptosis, fibrosis, and infarct zone size and improved cardiac function. shPHD2-ADSCs further increased these improvements and induced better ADSCs survival. Conditioned medium from the shPHD2-ADSCs decreased cardiomyocyte apoptosis and increased IGF-1
Godier-Furnémont et al ⁸⁷	Patches containing TGFβ-1–conditioned human MSCs	Nude rat	LAD occlusion	Decreased myocyte apoptosis
Karpov et al ⁸⁸	BM-MSCs and ADSCs injected 7 days after infarction	Rat	LAD occlusion and reperfusion	Animals transplanted with BM-MSCs preserved better left ventricular function and had decreased scar size
Jiang et al ⁸⁹	MSCs	Rat	LAD ligation and remote ischemic postconditioning	Remote ischemic postconditioning increased the SDF-1a concentration and increased the injected MSC retention
Hong et al ⁹⁰	ADSCs, EPCs, ADSCs + EPCs	Rat	LAD ligation	Increased LVEF and increased angiogenesis in the peri-infarct zone in the 3 groups
Sun et al ⁹¹	ADSCs embedded in a platelet-rich fibrin scaffold	Rat	AMI induced by LAD occlusion	Improved function and vascular remodeling with ADSCs embedded in the platelet-rich fibrin scaffold instead of directly

AD-CMGs, ADSC-derived cardiomyogenic cells; ADSCs, adipose-derived stem cells; AMI, acute myocardial infarction; Ang-1, angiopoietin-1; b-FGF, basic fibroblast growth factor; BM-MNCs, bone marrow-derived mononuclear cells; BM-MSCs, bone marrow-derived mesenchymal stem cells; CDCs, cardiosphere-derived cells; eNOS, endothelial nitric oxide synthase; EPCs, endothelial progenitor cells; GFP, green fluorescent protein; HGF, hepatic growth factor; HO-1-ADSCs, adipose-derived stem cells transduced with heme oxygenase-1; IGF-1, insulin-like growth factor 1; LAD, left anterior descending coronary artery; LVEF, left ventricular ejection fraction; MSCs, mesenchymal stem cells; SCID, severe combined immunodeficiency; SD, Sprague Dawley; shPHD2, prolyl hydroxylase domain protein 2; SNAP, S-Nitroso-N-acetyl-DL-penicillamine; SVF, stromal vascular fraction; TGFβ-1, transforming growth factor β1; VEGF, vascular endothelial growth factor.

question whether the benefits derived from ADSC administration are directly related to the differentiation processes or if, in contrast, they are conditioned by ADSC secretion of paracrine factors.^{106–108}

Another important function of ADSCs related to ischemic heart disease is derived from their angiogenic potential.^{52,109} Because ADSCs secrete a large number of proangiogenic cytokines and cytoprotective factors, they are an ideal cellular source for angiogenic therapy and apoptosis inhibition.^{29,110–112} An in vivo study showed that intramyocardial injection of human ADSCs significantly promoted angiogenesis and inhibited cell apoptosis in an infarcted heart 4 weeks after their injection.⁵⁴ Moreover, ADSCs showed an increased expression of vascular endothelial and fibroblast growth factors and stromal cell-derived factor 1.⁵⁴ Indeed, the interaction of stromal cell-derived factor 1 with its receptor induces the rapid mobilization of stem/progenitor cells from the bone marrow,¹¹³ which is essential for the revascularization of body systems.¹¹⁴ All of these results indicate that

injected ADSCs cooperate with stem/progenitor cells of the bone marrow via cellular mobilization, which is promoted by stromal cell-derived factor 1 and which boosts angiogenesis and vasculogenesis in myocardial ischemia.⁵⁴

The functional response of ADSCs can also be affected by oxygen concentration.^{29,115} Rehman et al²⁹ found that ADSCs secreted up to 5 times more vascular endothelial growth factors if they were cultured under hypoxic conditions. Additionally, the conditioned supernatant of ADSCs cultured under hypoxic conditions increased the growth of endothelial cells and reduced their apoptosis. Recently, a study reported that hypoxic preconditioning of ADSCs increased their survival and paracrine effects in a hypoxia-inducible factor 1-mediated manner.¹¹⁶ Indeed, our laboratory group has shown that ADSCs cultured under hypoxic conditions had greatly improved proliferative capacity.⁴⁰ These results showed that ADSCs respond to ischemic situations and promote angiogenesis via the secretion of vascular endothelial growth factors. As already mentioned, various experimental studies have

Table 4
Preclinical Studies with Porcine Models

Authors	Cell source	Animal model	Lesion	Result
Watanabe et al ⁹⁶	ADSCs	Pig	AMI induced by LAD occlusion	After 6 months, LVEF increased by 3%
Fotuhi et al ⁹⁷	ADSCs from pig subcutaneous adipose tissue	Pig	AMI induced by LAD ligation	Decreased arrhythmogenesis
Valina et al ⁹³	Pig ADSCs from subcutaneous adipose tissue or BM-MSCs	Pig	AMI induced by LAD angioplasty	After 4 weeks, improved LVEF, capillary density, and thickening of the heart wall
Alt et al ⁹⁴	ADSCs from pig subcutaneous adipose tissue	Pig	AMI induced by LAD occlusion and reperfusion	Better perfusion, LVEF, capillary density, and myocardial recovery
Rigol et al ⁹²	ADSCs from pig subcutaneous adipose tissue via intracoronary and transendocardial administration	Pig	AMI	Increased number of small vessels. Ejection fraction unchanged
Mazo et al ⁹⁵	Pig ADSCs	Minipigs	Ischemia and reperfusion	After 3 months, improved cardiac function, increased angiogenesis and vasculogenesis, and decreased fibrosis and cardiac hypertrophy
Yang et al ¹⁰¹	Human ADSCs immobilized in agarose gel patches	Pig	AMI induced by cryolesion	After 4 weeks, improved perfusion, reduced infarct zone size, and increased cardiac kinetics
Song et al ⁹⁸	MSCs, atorvastatin, and NG-nitrol-L-arginine	Minipigs	LAD ligation and reperfusion	After 4 weeks, atorvastatin + MSCs increased LVEF and decreased the inflammation, fibrosis, and apoptosis indices. There was no improvement in cardiac function with atorvastatin or MSCs alone. NG-nitrol-L-arginine partially blocked the improvements seen
Yin et al ⁹⁹	CMTA + CsA-NP	Minipigs	AMI induced by LAD occlusion	CsA-NP increased ADSC viability and improved LVEF
Rigol et al ¹⁰⁰	ADSCs	Pig	AMI	ADSCs administration immediately after reperfusion is more effective and improves neovascularization

ADSCs, adipose-derived stem cells; AMI, acute myocardial infarction; BM-MSCs, bone marrow-derived mesenchymal stem cells; CsA-NP, cyclosporine A-nanoparticle emulsion; LAD, left anterior descending coronary artery; LVEF, left ventricular ejection fraction; MSCs, mesenchymal stem cells.

shown that ADSCs are safe and efficacious (Tables 3 and 4). As can be seen in Table 4, 2 studies showed an increased capillary density in the area surrounding the infarcted heart and an increased cardiac function 1 month after a myocardial infarction when animals were treated with ADSCs, results similar to those observed after bone marrow-derived MSCs administration.^{93,94} A long-term follow-up study showed that, despite an inability to detect ADSCs in the myocardium 3 months after their injection, ADSC transplantation was associated with increased cardiac function, positive remodeling, and increased angiogenesis and vasculogenesis, confirming the long-term paracrine effect of these cells.⁹⁵ However, these observations indicate that, despite all of the advantages of ADSCs, their low capacity for homing to ischemic tissue is an obstacle to their clinical use.¹¹⁷ Accordingly, various strategies are being explored to resolve the problem of ADSC survival and engraftment in host tissue, such as the administration of ADSCs together with a combination of growth factors,¹¹⁸ injection of genetically modified ADSCs,¹¹⁹ and/or the use of grafts/biomaterial scaffolds.^{120,121} Finally, it is necessary to mention that various studies have indicated that ADSCs, both in vitro and in vivo, can boost the differentiation of the vasculature into pericytes, cells capable of creating microvessels, preventing vascular regression, and promoting long-term microvessel maintenance.^{122,123}

Clinical Trials

The evidence found in the experimental studies of the potential of ADSCs to repair the ischemic myocardium and restore its functional capacity has prompted the performance of clinical trials in this area. However, although some researchers believe that ADSCs will be used in the coming years as a cell therapy aimed at repairing the damaged heart, others believe that there are many unknowns to be resolved before these cells can be clinically used. So far, ADSCs have been satisfactorily used to treat some diseases, such as Crohn's fistula, osteogenesis imperfecta, and breast reconstruction after a partial mastectomy (Table 5). However, the use of ADSCs in the field of ischemic heart disease is still in phase I-II. Various clinical trials have been initiated to determine the feasibility, safety, and efficacy of the use of ADSCs in patients who have had an acute myocardial infarction (APOLLO, ADI-ME-CHF-002, ADVANCE, and ACUTE MI), have chronic ischemic heart disease (PRECISE, MyStromalCell, ATHENA, and ATHENA II), or nonischemic cardiomyopathy (ADI-ME-CHF-002). Of these, the APOLLO¹²⁴, PRECISE,¹²⁵ and MyStromalCell¹²⁶ trials have been completed. These studies showed that the use of ADSCs is safe and feasible. In addition, the results indicated that ADSC use preserves cardiac function, improves cardiac perfusion, and even reduces scar tissue size, thereby reinforcing the findings of the previous preclinical trials.

FUTURE PROSPECTS

Thus, ADSCs are appearing as a viable cell therapy alternative in various medical fields, which will require a better understanding of the mechanisms used by these cells or their paracrine factors in tissue regeneration/recovery, as well as the key molecular factors promoting the differentiation of ADSCs to different lineages. Additionally, it remains to be determined whether therapeutic efficacy is affected by the anatomical source of the cells, the sex and age of the donor, or the presence of comorbidities. The possibility of using both autologous and allogeneic ADSCs should also be considered, because various independent studies have reported that ADSCs have a low immunostimulatory potential.^{127,128}

Table 5

Clinical Trials Performed with Adipose-derived Stem Cells

Disorder	Studies, no.
<i>Metabolic disease</i>	
Lipodystrophy	2
Diabetes mellitus	5
<i>Cardiovascular disease</i>	
Vascular disease	2
Ischemic heart disease	11
Stroke	2
<i>Rheumatic disorders</i>	
Tendon injury	1
Arthritis	6
Degenerative disc disease	1
Necrosis	1
<i>Renal and urological disorders</i>	
Urinary incontinence	4
Renal failure	1
Urethral disorders	1
<i>Endocrine system diseases</i>	
<i>Respiratory disorders</i>	
<i>Nervous system diseases</i>	
Ataxia	1
Facial hemiatrophy	1
Multiple sclerosis	2
Parkinson disease	1
Spinal cord disorders	3
Brain damage	2
<i>Mental disorders</i>	
<i>Digestive system diseases</i>	
Intestinal disorders	2
Cirrhosis	1
Fibrosis	11
<i>Breast diseases</i>	
Graft-versus-host disease	7
Crohn disease	2
Autologous fat grafting	3
Frailty syndrome	1

ACKNOWLEDGMENTS

We thank the *Fundación Jesús Serra* of Barcelona for its continuing support. G. Vilahur is a Ramón y Cajal researcher with a contract with the *Secretaría de Estado de Investigación, Desarrollo e Innovación* from the *Ministerio de Economía y Competitividad* of Spain (RyC-2009-5495).

FUNDING

Part of the work contained in this manuscript was financed by the *Programa Nacional de Salud* (SAF 2013-42962-R, awarded to L. Badimon; SAF 2012-40208 awarded to G. Vilahur), the *Instituto de Salud Carlos III* (TerCel [Red de Terapia Celular] RD12/0019/0026), and the *Fundación Jesús Serra* (FIC-Barcelona).

CONFLICTS OF INTEREST

None declared.

REFERENCES

- Kershaw EE, Flier JS. Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab.* 2004;89:2548-56.
- Gil A, Olza J, Gil-Campos M, Gomez-Llorente C, Aguilera CM. Is adipose tissue metabolically different at different sites? *Int J Pediatr Obes.* 2011;6 Suppl 1: 13-20.
- Villaret A, Galitzky J, Decaunes P, Esteve D, Marques MA, Sengenès C, et al. Adipose tissue endothelial cells from obese human subjects: differences among depots in angiogenic, metabolic, and inflammatory gene expression and cellular senescence. *Diabetes.* 2010;59:2755-63.
- Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obes Rev.* 2010;11:11-8.
- Dobbelsteyn CJ, Joffres MR, MacLean DR, Flowerdew G. A comparative evaluation of waist circumference, waist-to-hip ratio and body mass index as indicators of cardiovascular risk factors. The Canadian Heart Health Surveys. *Int J Obes Relat Metab Disord.* 2001;25:652-61.
- Kanai H, Matsuzawa Y, Kotani K, Keno Y, Kobatake T, Nagai Y, et al. Close correlation of intra-abdominal fat accumulation to hypertension in obese women. *Hypertension.* 1990;16:484-90.
- Ayonrinde OT, Olynyk JK, Beilin LJ, Mori TA, Pennell CE, De Klerk N, et al. Gender-specific differences in adipose distribution and adipocytokines influence adolescent nonalcoholic fatty liver disease. *Hepatology.* 2011;53:800-9.
- Wronska A, Kmiec Z. Structural and biochemical characteristics of various white adipose tissue depots. *Acta Physiol (Oxf).* 2012;205:194-208.
- Jaworski K, Sarkadi-Nagy E, Duncan RE, Ahmadian M, Sul HS. Regulation of triglyceride metabolism. IV. Hormonal regulation of lipolysis in adipose tissue. *Am J Physiol Gastrointest Liver Physiol.* 2007;293:G1-4.
- Duplus E, Glorian M, Forest C. Fatty acid regulation of gene transcription. *J Biol Chem.* 2000;275:30749-52.
- Sheehan MT, Jensen MD. Metabolic complications of obesity. Pathophysiologic considerations. *Med Clin North Am.* 2000;84:363-85. vi.
- Shepherd PR, Gnudi L, Tozzo E, Yang H, Leach F, Kahn BB. Adipose cell hyperplasia and enhanced glucose disposal in transgenic mice overexpressing GLUT4 selectively in adipose tissue. *J Biol Chem.* 1993;268:2243-6.
- Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res.* 2007;100:1249-60.
- Rutkowski JM, Davis KE, Scherer PE. Mechanisms of obesity and related pathologies: the macro- and microcirculation of adipose tissue. *FEBS J.* 2009;276:5738-46.
- Schipper HS, Prakken B, Kalkhoven E, Boes M. Adipose tissue-resident immune cells: key players in immunometabolism. *Trends Endocrinol Metab.* 2012;23:407-15.
- Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante Jr AW. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest.* 2003;112:1796-808.
- Lumeng CN, Bodzin JL, Saltiel AR. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest.* 2007;117:175-84.
- Cawthorn WP, Scheller EL, MacDougald OA. Adipose tissue stem cells meet preadipocyte commitment: going back to the future. *J Lipid Res.* 2012;53: 227-46.
- Kim S, Moustaid-Moussa N. Secretory, endocrine and autocrine/paracrine function of the adipocyte. *J Nutr.* 2000;130:S3110-5.
- Ronti T, Lupattelli G, Mannarino E. The endocrine function of adipose tissue: an update. *Clin Endocrinol (Oxf).* 2006;64:355-65.
- Tesauro M, Canale MP, Rodia G, Di Daniele N, Lauro D, Scuteri A, et al. Metabolic syndrome, chronic kidney, and cardiovascular diseases: role of adipokines. *Cardiol Res Pract.* 2011;653182. <http://dx.doi.org/10.4061/2011/653182>.
- Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, et al. Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Engineering.* 2001;7:211-28.
- Gesta S, Tseng YH, Kahn CR. Developmental origin of fat: tracking obesity to its source. *Cell.* 2007;131:242-56.
- De Ugarte DA, Morizono K, Elbarbary A, Alfonso Z, Zuk PA, Zhu M, et al. Comparison of multi-lineage cells from human adipose tissue and bone marrow. *Cells Tissues Organs.* 2003;174:101-9.
- Mansilla E, Díaz Aquino V, Zambón D, Marin GH, Mártire K, Roque G, et al. Could metabolic syndrome, lipodystrophy, and aging be mesenchymal stem cell exhaustion syndromes? *Stem Cells Int.* 2011;943216. <http://dx.doi.org/10.4061/2011/943216>.
- Kolonin MG, Evans KW, Mani SA, Gomer RH. Alternative origins of stroma in normal organs and disease. *Stem Cell Res.* 2012;8:312-23.
- Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy.* 2006;8:315-7.
- Salgado AJ, Reis RL, Sousa NJ, Gimble JM. Adipose tissue derived stem cells secrete: soluble factors and their roles in regenerative medicine. *Curr Stem Cell Res Ther.* 2010;5:103-10.
- Rehman J, Traktuev D, Li J, Merfeld-Clauss S, Temm-Grove CJ, Bovenkerk JE, et al. Secretion of angiogenic and antiapoptotic factors by human adipose stromal cells. *Circulation.* 2004;109:1292-8.
- Perrini S, Laviola L, Cignarelli A, Melchiorre M, De Stefano F, Caccioppoli C, et al. Fat depot-related differences in gene expression, adiponectin secretion, and insulin action and signalling in human adipocytes differentiated in vitro from precursor stromal cells. *Diabetologia.* 2008;51:155-64.
- Tchkonia T, Giorgadze N, Pirtskhalava T, Thomou T, DePonte M, Koo A, et al. Fat depot-specific characteristics are retained in strains derived from single human preadipocytes. *Diabetes.* 2006;55:2571-8.
- Van Harmelen V, Rohrig K, Hauner H. Comparison of proliferation and differentiation capacity of human adipocyte precursor cells from the omental and subcutaneous adipose tissue depot of obese subjects. *Metabolism.* 2004;53: 632-7.
- Tchkonia T, Lenburg M, Thomou T, Giorgadze N, Frampton G, Pirtskhalava T, et al. Identification of depot-specific human fat cell progenitors through distinct expression profiles and developmental gene patterns. *Am J Physiol Endocrinol Metab.* 2007;292:E298-307.
- Gesta S, Blüher M, Yamamoto Y, Norris AW, Berndt J, Kralisch S, et al. Evidence for a role of developmental genes in the origin of obesity and body fat distribution. *Proc Natl Acad Sci U S A.* 2006;103:6676-81.
- Majka SM, Barak Y, Klemm DJ. Concise review: adipocyte origins: weighing the possibilities. *Stem Cells.* 2011;29:1034-40.
- Hoffstedt J, Arner E, Wahrenberg H, Andersson DP, Qvist V, Lofgren P, et al. Regional impact of adipose tissue morphology on the metabolic profile in morbid obesity. *Diabetologia.* 2010;53:2496-503.
- Ferrer-Lorente R, Bejar MT, Tous M, Vilahur G, Badimon L. Systems biology approach to identify alterations in the stem cell reservoir of subcutaneous adipose tissue in a rat model of diabetes: effects on differentiation potential and function. *Diabetologia.* 2014;57:246-56.
- Van Harmelen V, Skurk T, Rohrig K, Lee YM, Halbleib M, Aprath-Husmann I, et al. Effect of BMI and age on adipose tissue cellularity and differentiation capacity in women. *Int J Obes Relat Metab Disord.* 2003;27:889-95.
- Nair S, Lee YH, Rousseau E, Cam M, Tataranni PA, Baier LJ, et al. Increased expression of inflammation-related genes in cultured preadipocytes/stromal vascular cells from obese compared with non-obese Pima Indians. *Diabetologia.* 2005;48:1784-8.
- Oñate B, Vilahur G, Ferrer-Lorente R, Ybarra J, Díez-Caballero A, Ballesta-López C, et al. The subcutaneous adipose tissue reservoir of functionally active stem cells is reduced in obese patients. *FASEB J.* 2012;26:4327-36.
- Oñate B, Vilahur G, Camino-López S, Díez-Caballero A, Ballesta-López C, Ybarra J, et al. Stem cells isolated from adipose tissue of obese patients show changes in their transcriptomic profile that indicate loss in stemcellness and increased commitment to an adipocyte-like phenotype. *BMC Genomics.* 2013;14:625.
- Roldan M, Macias-Gonzalez M, Garcia R, Tinahones FJ, Martín M. Obesity short-circuits stemness gene network in human adipose multipotent stem cells. *FASEB J.* 2011;25:4111-26.
- Smart N, Riley PR. The stem cell movement. *Circ Res.* 2008;102:1155-68.
- Gimble JM, Bunnell BA, Guilak F. Human adipose-derived cells: an update on the transition to clinical translation. *Regen Med.* 2012;7:225-35.
- Mizuno H, Tobita M, Uysal AC. Concise review: Adipose-derived stem cells as a novel tool for future regenerative medicine. *Stem Cells.* 2012;30:804-10.
- Miranville A, Heeschen C, Sengenès C, Curat CA, Busse R, Bouloumié A. Improvement of postnatal neovascularization by human adipose tissue-derived stem cells. *Circulation.* 2004;110:349-55.
- Makino S, Fukuda K, Miyoshi S, Konishi F, Kodama H, Pan J, et al. Cardiomyocytes can be generated from marrow stromal cells in vitro. *J Clin Invest.* 1999;103:697-705.
- Prockop DJ. Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science.* 1997;276:71-4.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage potential of adult human mesenchymal stem cells. *Science.* 1999;284:143-7.
- Nagaya N, Fujii T, Iwase T, Ohgushi H, Itoh T, Uematsu M, et al. Intravenous administration of mesenchymal stem cells improves cardiac function in rats with acute myocardial infarction through angiogenesis and myogenesis. *Am J Physiol Heart Circ Physiol.* 2004;287:H2670-6.
- Strem BM, Zhu M, Alfonso Z, Daniels EJ, Schreiber R, Beygui R, et al. Expression of cardiomyocytic markers on adipose tissue-derived cells in a murine model of acute myocardial injury. *Cytotherapy.* 2005;7:282-91.
- Cai L, Johnstone BH, Cook TG, Tan J, Fishbein MC, Chen PS, et al. IFATS collection: Human adipose tissue-derived stem cells induce angiogenesis and nerve sprouting following myocardial infarction, in conjunction with potent preservation of cardiac function. *Stem Cells.* 2009;27:230-7.
- Bai X, Yan Y, Song YH, Seidensticker M, Rabinovich B, Metzler R, et al. Both cultured and freshly isolated adipose tissue-derived stem cells enhance cardiac function after acute myocardial infarction. *Eur Heart J.* 2010;31: 489-501.
- Ii M, Horii M, Yokoyama A, Shoji T, Mifune Y, Kawamoto A, et al. Synergistic effect of adipose-derived stem cell therapy and bone marrow progenitor recruitment in ischemic heart. *Lab Invest.* 2011;91:539-52.
- Miyahara Y, Nagaya N, Kataoka M, Yanagawa B, Tanaka K, Hao H, et al. Monolayered mesenchymal stem cells repair scarred myocardium after myocardial infarction. *Nat Med.* 2006;12:459-65.
- Zhang DZ, Gai LY, Liu HW, Jin QH, Huang JH, Zhu XY. Transplantation of autologous adipose-derived stem cells ameliorates cardiac function in rabbits with myocardial infarction. *Chin Med J (Engl).* 2007;120:300-7.
- Mazo M, Planat-Bénard V, Abizanda G, Pelacho B, Léobon B, Gavira JJ, et al. Transplantation of adipose derived stromal cells is associated with functional improvement in a rat model of chronic myocardial infarction. *Eur J Heart Fail.* 2008;10:454-62.

58. Léobon B, Roncalli J, Joffre C, Mazo M, Boisson M, Barreau C, et al. Adipose-derived cardiomyogenic cells: in vitro expansion and functional improvement in a mouse model of myocardial infarction. *Cardiovasc Res*. 2009;83:757-67.
59. Schenke-Layland K, Strem BM, Jordan MC, Deemedio MT, Hedrick MH, Roos KP, et al. Adipose tissue-derived cells improve cardiac function following myocardial infarction. *J Surg Res*. 2009;153:217-23.
60. Van der Bogt KE, Schrepfer S, Yu J, Sheikh AY, Hoyt G, Govaert JA, et al. Comparison of transplantation of adipose tissue- and bone marrow-derived mesenchymal stem cells in the infarcted heart. *Transplantation*. 2009;87:642-52.
61. Wang L, Deng J, Tian W, Xiang B, Yang T, Li G, et al. Adipose-derived stem cells are an effective cell candidate for treatment of heart failure: an MR imaging study of rat hearts. *Am J Physiol Heart Circ Physiol*. 2009;297:H1020-31.
62. Zhu XY, Zhang XZ, Xu L, Zhong XY, Ding Q, Chen YX. Transplantation of adipose-derived stem cells overexpressing hHGF into cardiac tissue. *Biochem Biophys Res Commun*. 2009;379:1084-90.
63. Bayes-Genis A, Soler-Botija C, Farré J, Sepulveda P, Raya A, Roura S, et al. Human progenitor cells derived from cardiac adipose tissue ameliorate myocardial infarction in rodents. *J Mol Cell Cardiol*. 2010;49:771-80.
64. Danoviz ME, Nakamura JS, Marques FL, Dos Santos L, Alvarenga EC, Dos Santos AA, et al. Rat adipose tissue-derived stem cells transplantation attenuates cardiac dysfunction post infarction and biopolymers enhance cell retention. *PLoS One*. 2010;5:e12077.
65. Hwangbo S, Kim J, Her S, Cho H, Lee J. Therapeutic potential of human adipose stem cells in a rat myocardial infarction model. *Yonsei Med J*. 2010;51:69-76.
66. Lin YC, Leu S, Sun CK, Yen CH, Kao YH, Chang LT, et al. Early combined treatment with sildenafil and adipose-derived mesenchymal stem cells preserves heart function in rat dilated cardiomyopathy. *J Transl Med*. 2010;8:88.
67. Okura H, Matsuyama A, Lee CM, Saga A, Kakuta-Yamamoto A, Nagao A, et al. Cardiomyoblast-like cells differentiated from human adipose tissue-derived mesenchymal stem cells improve left ventricular dysfunction and survival in a rat myocardial infarction model. *Tissue Eng Part C Methods*. 2010;16:417-25.
68. Zhang X, Wang H, Ma X, Adila A, Wang B, Liu F, et al. Preservation of the cardiac function in infarcted rat hearts by the transplantation of adipose-derived stem cells with injectable fibrin scaffolds. *Exp Biol Med (Maywood)*. 2010;235:1505-15.
69. Bai X, Yan Y, Coleman M, Wu G, Rabinovich B, Seidensticker M, et al. Tracking long-term survival of intramyocardially delivered human adipose tissue-derived stem cells using bioluminescence imaging. *Mol Imaging Biol*. 2011;13:633-45.
70. Berardi GR, Rebelatto CK, Tavares HF, Ingberman M, Shigunov P, Barchiki F, et al. Transplantation of SNAP-treated adipose tissue-derived stem cells improves cardiac function and induces neovascularization after myocardial infarct in rats. *Exp Mol Pathol*. 2011;90:149-56.
71. Cai A, Zheng D, Dong Y, Qiu R, Huang Y, Song Y, et al. Efficacy of atorvastatin combined with adipose-derived mesenchymal stem cell transplantation on cardiac function in rats with acute myocardial infarction. *Acta Biochim Biophys Sin (Shanghai)*. 2011;43:857-66.
72. Gaebel R, Furlani D, Sorg H, Polchow B, Frank J, Bieback K, et al. Cell origin of human mesenchymal stem cells determines a different healing performance in cardiac regeneration. *PLoS One*. 2011;6:e15652.
73. Hamdi H, Planat-Benard V, Bel A, Puymirat E, Geha R, Pidial L, et al. Epicardial adipose stem cell sheets results in greater post-infarction survival than intramyocardial injections. *Cardiovasc Res*. 2011;91:483-91.
74. Van Dijk A, Naaijken BA, Jurgens WJ, Nalliah K, Sairras S, Van der Pijl RJ, et al. Reduction of infarct size by intravenous injection of uncultured adipose derived stromal cells in a rat model is dependent on the time point of application. *Stem Cell Res*. 2011;7:219-29.
75. Bagno LL, Werneck-de-Castro JP, Oliveira PF, Cunha-Abreu MS, Rocha NN, Kasai-Brunswick TH, et al. Adipose-derived stromal cell therapy improves cardiac function after coronary occlusion in rats. *Cell Transplant*. 2012;21:1985-96.
76. Beitnes JO, Oie E, Shahdadfar A, Karlsten T, Muller RM, Aakhus S, et al. Intramyocardial injections of human mesenchymal stem cells following acute myocardial infarction modulate scar formation and improve left ventricular function. *Cell Transplant*. 2012;21:1697-709.
77. Fang CH, Jin J, Joe JH, Song YS, So BI, Lim SM, et al. In vivo differentiation of human amniotic epithelial cells into cardiomyocyte-like cells and cell transplantation effect on myocardial infarction in rats: comparison with cord blood and adipose tissue-derived mesenchymal stem cells. *Cell Transplant*. 2012;21:1687-96.
78. Hoke NN, Salloum FN, Kass DA, Das A, Kukreja RC. Preconditioning by phosphodiesterase-5 inhibition improves therapeutic efficacy of adipose-derived stem cells following myocardial infarction in mice. *Stem Cells*. 2012;30:326-35.
79. Li TS, Cheng K, Malliaras K, Smith RR, Zhang Y, Sun B, et al. Direct comparison of different stem cell types and subpopulations reveals superior paracrine potency and myocardial repair efficacy with cardiosphere-derived cells. *J Am Coll Cardiol*. 2012;59:942-53.
80. Liu Z, Wang H, Wang Y, Lin Q, Yao A, Cao F, et al. The influence of chitosan hydrogel on stem cell engraftment, survival and homing in the ischemic myocardial microenvironment. *Biomaterials*. 2012;33:3093-106.
81. Paul A, Chen G, Khan A, Rao VT, Shum-Tim D, Prakash S. Genipin-cross-linked microencapsulated human adipose stem cells augment transplant retention resulting in attenuation of chronically infarcted rat heart fibrosis and cardiac dysfunction. *Cell Transplant*. 2012;21:2735-51.
82. Paul A, Nayan M, Khan AA, Shum-Tim D, Prakash S. Angiotensin-1-expressing adipose stem cells genetically modified with baculovirus nanocomplex: investigation in rat heart with acute infarction. *Int J Nanomedicine*. 2012;7:663-82.
83. Shi CZ, Zhang XP, Lv ZW, Zhang HL, Xu JZ, Yin ZF, et al. Adipose tissue-derived stem cells embedded with eNOS restore cardiac function in acute myocardial infarction model. *Int J Cardiol*. 2012;154:2-8.
84. Yang JJ, Yang X, Liu ZQ, Hu SY, Du ZY, Feng LL, et al. Transplantation of adipose tissue-derived stem cells overexpressing heme oxygenase-1 improves functions and remodeling of infarcted myocardium in rabbits. *Tohoku J Exp Med*. 2012;226:231-41.
85. Paul A, Srivastava S, Chen G, Shum-Tim D, Prakash S. Functional assessment of adipose stem cells for xenotransplantation using myocardial infarction immunocompetent models: comparison with bone marrow stem cells. *Cell Biochem Biophys*. 2013;67:263-73.
86. Wang WE, Yang D, Li L, Wang W, Peng Y, Chen C, et al. Prolyl hydroxylase domain protein 2 silencing enhances the survival and paracrine function of transplanted adipose-derived stem cells in infarcted myocardium. *Circ Res*. 2013;113:288-300.
87. Godier-Furnémont AF, Tekabe Y, Kollaros M, Eng G, Morales A, Vunjak-Novakovic G, et al. Noninvasive imaging of myocyte apoptosis following application of a stem cell-engineered delivery platform to acutely infarcted myocardium. *J Nucl Med*. 2013;54:977-83.
88. Karpov AA, Uspenskaya YK, Minasian SM, Puzanov MV, Dmitrieva RI, Bilibina AA, et al. The effect of bone marrow- and adipose tissue-derived mesenchymal stem cell transplantation on myocardial remodeling in the rat model of ischaemic heart failure. *Int J Exp Pathol*. 2013;94:169-77.
89. Jiang Q, Song P, Wang E, Li J, Hu S, Zhang H. Remote ischemic preconditioning enhances cell retention in the myocardium after intravenous administration of bone marrow mesenchymal stromal cells. *J Mol Cell Cardiol*. 2013;56:1-7.
90. Hong SJ, Kihlken J, Choi SC, March KL, Lim DS. Intramyocardial transplantation of human adipose-derived stromal cell and endothelial progenitor cell mixture was not superior to individual cell type transplantation in improving left ventricular function in rats with myocardial infarction. *Int J Cardiol*. 2013;164:205-11.
91. Sun CK, Zhen YY, Leu S, Tsai TH, Chang LT, Sheu JJ, et al. Direct implantation versus platelet-rich fibrin-embedded adipose-derived mesenchymal stem cells in treating rat acute myocardial infarction. *Int J Cardiol*. 2014;173:410-23.
92. Rigol M, Solanes N, Farré J, Roura S, Roqué M, Berrueto A, et al. Effects of adipose tissue-derived stem cell therapy after myocardial infarction: impact of the route of administration. *J Card Fail*. 2010;16:357-66.
93. Valina C, Pinkernell K, Song YH, Bai X, Sadat S, Campeau RJ, et al. Intracoronary administration of autologous adipose tissue-derived stem cells improves left ventricular function, perfusion, and remodeling after acute myocardial infarction. *Eur Heart J*. 2007;28:2667-77.
94. Alt E, Pinkernell K, Scharlau M, Coleman M, Fotuhi P, Nabzdyk C, et al. Effect of freshly isolated autologous tissue resident stromal cells on cardiac function and perfusion following acute myocardial infarction. *Int J Cardiol*. 2010;144:26-35.
95. Mazo M, Hernández S, Gavira JJ, Abizanda G, Arana M, López-Martínez T, et al. Treatment of reperfused ischemia with adipose-derived stem cells in a preclinical swine model of myocardial infarction. *Cell Transplant*. 2012;21:2723-33.
96. Watanabe C. Intracoronary adipose tissue derived stem cells therapy preserves left ventricular function in a porcine infarct model. *Transvascular Cardiovascular Therapeutics Annual Meeting*. Washington, September 2004.
97. Fotuhi P, Song YH, Alt E. Electrophysiological consequence of adipose-derived stem cell transplantation in infarcted porcine myocardium. *Europace*. 2007;9:1218-21.
98. Song L, Yang YJ, Dong QT, Qian HY, Gao RL, Qiao SB, et al. Atorvastatin enhance efficacy of mesenchymal stem cells treatment for swine myocardial infarction via activation of nitric oxide synthase. *PLoS One*. 2013;8:e65702.
99. Yin QX, Wang H, Pei ZY, Zhao YS. [Efficacy of cyclosporine A-nanoparticles emulsion combined with stem cell transplantation therapy for acute myocardial infarction]. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*. 2013;35:404-10.
100. Rigol M, Solanes N, Roura S, Roqué M, Novensà L, Dantas AP, et al. Allogeneic adipose stem cell therapy in acute myocardial infarction. *Eur J Clin Invest*. 2014;44:83-92.
101. Yang Y, Dreesen de Gervai P, Sun J, Glogowski M, Gussakovskiy E, Kupriyanov V. MRI studies of cryoinjury infarction in pig hearts: ii. Effects of intrapericardial delivery of adipose-derived stem cells (ADSC) embedded in agarose gel. *NMR Biomed*. 2012;25:227-35.
102. Bai X, Alt E. Myocardial regeneration potential of adipose tissue-derived stem cells. *Biochem Biophys Res Commun*. 2010;401:321-6.
103. Kim YM, Jeon ES, Kim MR, Jho SK, Ryu SW, Kim JH. Angiotensin II-induced differentiation of adipose tissue-derived mesenchymal stem cells to smooth muscle-like cells. *Int J Biochem Cell Biol*. 2008;40:2482-91.
104. Rodríguez LV, Alfonso Z, Zhang R, Leung J, Wu B, Ignarro LJ. Clonogenic multipotent stem cells in human adipose tissue differentiate into functional smooth muscle cells. *Proc Natl Acad Sci U S A*. 2006;103:12167-72.
105. Ning H, Liu G, Lin G, Yang R, Lue TF, Lin CS. Fibroblast growth factor 2 promotes endothelial differentiation of adipose tissue-derived stem cells. *J Sex Med*. 2009;6:967-79.

106. Hombach-Klonisch S, Panigrahi S, Rashedi I, Seifert A, Alberti E, Pocar P, et al. Adult stem cells and their trans-differentiation potential-perspectives and therapeutic applications. *J Mol Med*. 2008;86:1301–14.
107. Jackson KA, Majka SM, Wang H, Pocius J, Hartley CJ, Majesky MW, et al. Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells. *J Clin Invest*. 2001;107:1395–402.
108. Laflamme MA, Myerson D, Saffitz JE, Murry CE. Evidence for cardiomyocyte repopulation by extracardiac progenitors in transplanted human hearts. *Circ Res*. 2002;90:634–40.
109. Gneocchi M, He H, Liang OD, Melo LG, Morello F, Mu H, et al. Paracrine action accounts for marked protection of ischemic heart by Akt-modified mesenchymal stem cells. *Nat Med*. 2005;11:367–8.
110. Nakagami H, Maeda K, Morishita R, Iguchi S, Nishikawa T, Takami Y, et al. Novel autologous cell therapy in ischemic limb disease through growth factor secretion by cultured adipose tissue-derived stromal cells. *Arterioscler Thromb Vasc Biol*. 2005;25:2542–7.
111. Wang M, Crisostomo PR, Herring C, Meldrum KK, Meldrum DR. Human progenitor cells from bone marrow or adipose tissue produce VEGF, HGF, and IGF-I in response to TNF by a p38 MAPK-dependent mechanism. *Am J Physiol Regul Integr Comp Physiol*. 2006;291:R880–4.
112. Kocher AA, Schuster MD, Szabolcs MJ, Takuma S, Burkhoff D, Wang J, et al. Neovascularization of ischemic myocardium by human bone-marrow-derived angioblasts prevents cardiomyocyte apoptosis, reduces remodeling and improves cardiac function. *Nat Med*. 2001;7:430–6.
113. Liles WC, Broxmeyer HE, Rodger E, Wood B, Hubel K, Cooper S, et al. Mobilization of hematopoietic progenitor cells in healthy volunteers by AMD3100, a CXCR4 antagonist. *Blood*. 2003;102:2728–30.
114. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, et al. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature*. 1998;393:591–4.
115. Rasmussen JG, Probert O, Pilgaard L, Kastrup J, Simonsen U, Zachar V, et al. Prolonged hypoxic culture and trypsinization increase the pro-angiogenic potential of human adipose tissue-derived stem cells. *Cytotherapy*. 2011;13:318–28.
116. Stubbs SL, Hsiao ST, Peshavariya HM, Lim SY, Disting GJ, Dilley RJ. Hypoxic preconditioning enhances survival of human adipose-derived stem cells and conditions endothelial cells in vitro. *Stem Cells Dev*. 2012;21:1887–96.
117. Tateishi-Yuyama E, Matsubara H, Murohara T, Ikeda U, Shintani S, Masaki H, et al. Therapeutic Angiogenesis using Cell Transplantation (TACT) Study Investigators. Therapeutic angiogenesis for patients with limb ischaemia by autologous transplantation of bone-marrow cells: a pilot study and a randomised controlled trial. *Lancet*. 2002;360:427–35.
118. Jay SM, Shepherd BR, Bertram JP, Pober JS, Saltzman WM. Engineering of multifunctional gels integrating highly efficient growth factor delivery with endothelial cell transplantation. *FASEB J*. 2008;22:2949–56.
119. Deuse T, Peter C, Fedak PW, Doyle T, Reichenspurner H, Zimmermann WH, et al. Hepatocyte growth factor or vascular endothelial growth factor gene transfer maximizes mesenchymal stem cell-based myocardial salvage after acute myocardial infarction. *Circulation*. 2009;120 Suppl 1:247–54.
120. Fitzpatrick 3rd JR, Frederick JR, McCormick RC, Harris DA, Kim AY, Muenzer JR, et al. Tissue-engineered pro-angiogenic fibroblast scaffold improves myocardial perfusion and function and limits ventricular remodeling after infarction. *J Thorac Cardiovasc Surg*. 2010;140:667–76.
121. Bhang SH, Cho SW, La WG, Lee TJ, Yang HS, Sun AY, et al. Angiogenesis in ischemic tissue produced by spheroid grafting of human adipose-derived stromal cells. *Biomaterials*. 2011;32:2734–47.
122. Amos PJ, Shang H, Bailey AM, Taylor A, Katz AJ, Peirce SM. IFATS collection: The role of human adipose-derived stromal cells in inflammatory microvascular remodeling and evidence of a perivascular phenotype. *Stem Cells*. 2008;26:2682–90.
123. Zannettino AC, Paton S, Arthur A, Khor F, Itescu S, Gimble JM, et al. Multipotential human adipose-derived stromal stem cells exhibit a perivascular phenotype in vitro and in vivo. *J Cell Physiol*. 2008;214:413–21.
124. Houtgraaf JH, Den Dekker WK, Van Dalen BM, Springeling T, De Jong R, Van Geuns RJ, et al. First experience in humans using adipose tissue-derived regenerative cells in the treatment of patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol*. 2012;59:539–40.
125. Perin EC, Sanz-Ruiz R, Sánchez PL, Lasso J, Pérez-Cano R, Alonso-Farto JC, et al. Adipose-derived regenerative cells in patients with ischemic cardiomyopathy: The PRECISE Trial. *Am Heart J*. 2014;168:88–95. e82.
126. Qayyum AA, Haack-Sørensen M, Mathiasen AB, Jørgensen E, Eklund A, Kastrup J. Adipose-derived mesenchymal stromal cells for chronic myocardial ischemia (MyStromalCell Trial): study design. *Regen Med*. 2012;7:421–8.
127. Kode JA, Mukherjee S, Joglekar MV, Hardikar AA. Mesenchymal stem cells: immunobiology and role in immunomodulation and tissue regeneration. *Cytotherapy*. 2009;11:377–91.
128. Puissant B, Barreau C, Bourin P, Clavel C, Corre J, Bousquet C, et al. Immunomodulatory effect of human adipose tissue-derived adult stem cells: comparison with bone marrow mesenchymal stem cells. *Br J Haematol*. 2005;129:118–29.