

Original article

Factors Associated With Errors in Visual Estimation of the Functional Significance of Coronary Lesions



Ramón López-Palop,* Pilar Carrillo, Pilar Agudo, Alberto Cordero, Araceli Frutos, Samer Mashlab, Rubén Martínez, Amin El Amrani, and David Ramos

Unidad de Hemodinámica, Sección de Cardiología, Hospital Universitario San Juan de Alicante, San Juan de Alicante, Alicante, Spain

Article history:

Received 20 August 2015

Accepted 3 December 2015

Available online 8 April 2016

Keywords:

Coronary heart disease

Coronary angiography

Coronary angioplasty

Fractional flow reserve

ABSTRACT

Introduction and objectives: Visual angiographic assessment continues to be used when decisions are made on whether to revascularize ambiguous coronary lesions. Multiple factors, other than the degree of stenosis, have been associated with the functional significance of a coronary lesion. The aim of this study was to investigate the ability of interventionists to visually predict the functional significance of a coronary lesion and the clinical and angiographic characteristics associated with errors in prediction.

Methods: We conducted a concordance study of the functional significance of coronary lesions predicted by experienced interventionists and fractional flow reserve values measured by intracoronary pressure wire in 665 intermediate lesions (40%-70% diameter stenosis) in 587 patients. We determined which factors were independently associated with errors in prediction.

Results: There was disagreement between the predicted fractional flow reserve value of ≤ 0.80 and the observed value in 30.1% of the lesions (overestimation: 11.3%; underestimation, 18.8%). Stent location in an artery other than the anterior descending artery or in a bifurcation was associated with overestimation. Male sex, severe calcification, and a greater myocardial territory distal to the lesion were significantly associated with the functional significance of the underestimated lesion.

Conclusions: Even when taking into account angiographic and clinical characteristics, there is a high rate of disagreement between visual estimation and direct measurement of intermediate coronary stenosis in relation to its functional significance. Specific angiographic and clinical characteristics are associated with an increased tendency to overestimate or underestimate the significance of lesions.

© 2016 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Factores asociados al error en la estimación visual de la importancia funcional de lesiones coronarias

RESUMEN

Introducción y objetivos: La valoración angiográfica visual sigue utilizándose para decidir la revascularización de lesiones coronarias dudosas. Múltiples factores, distintos del grado de estenosis, se han asociado con la repercusión funcional de una lesión coronaria. El objetivo de este estudio es analizar la capacidad de predecir visualmente la repercusión funcional de una lesión coronaria y los condicionantes clínicos y angiográficos asociados con el error en la predicción.

Métodos: Estudio de concordancia entre la predicción de repercusión funcional realizada por intervencionistas expertos y el valor de reserva fraccional de flujo obtenido mediante guía intracoronaria de presión en 665 lesiones intermedias (estenosis del 40-70% del diámetro) en 587 pacientes. Se determinaron los factores independientemente asociados a un error en la predicción.

Resultados: Se observó una discordancia del 30,1% (sobrestimación, 11,3%; subestimación, 18,8%) entre el valor de reserva fraccional de flujo predicho $\leq 0,80$ y el observado. La localización en el *stent*, en una arteria distinta de la descendente anterior y en una bifurcación se asoció a sobrestimación. El sexo masculino, la calcificación grave y el mayor territorio miocárdico distal a la lesión se asociaron significativamente con importancia funcional de la lesión subestimada.

Conclusiones: Incluso integrando características angiográficas y clínicas, la estimación visual de la importancia funcional de estenosis coronarias intermedias se asocia a una alta tasa de discrepancias respecto a su determinación real. Determinadas características angiográficas y clínicas se asocian específicamente con mayor tendencia a sobrestimar o subestimar la importancia de la lesión.

© 2016 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Enfermedad coronaria

Angiografía coronaria

Angioplastia coronaria

Reserva fraccional de flujo

* Corresponding author: Unidad de Hemodinámica, Hospital Universitario San Juan de Alicante, Ctra. Alicante-Valencia s/n, 03550 San Juan de Alicante, Alicante, Spain.
E-mail address: ramonlopezpalop@gmail.com (R. López-Palop).

Abbreviations

FFR: fractional flow reserve

INTRODUCTION

Although the main purpose of coronary revascularization is to treat myocardial ischemia, the degree of angiographic stenosis is used as the main parameter to guide decisions on whether to revascularize a lesion^{1–3} or to determine the presence of restenosis after treatment. The significance of a coronary lesion is classically defined by the degree of angiographic stenosis, which is obtained by dividing the minimal lumen diameter by the reference diameter in the projection showing the greatest stenosis.^{4,5} Invasive measurement of fractional flow reserve (FFR) has recently become established as the method of choice to determine the functional significance of coronary lesions, especially when their potential to produce ischemia is ambiguous.^{4,5} Several studies have shown the limitations of angiography to define the functional significance of a lesion.^{6–9} These limitations may be due to the difficulty of determining the true degree of stenosis in the presence of certain angiographic characteristics (curvature, calcification, bifurcations, ostial location, etc); another factor is that the functional significance of a lesion is determined by other factors that add to the degree of stenosis. The FFR across a specific lesion is affected, among others factors, by the size of the myocardial territory perfused by the vessel with the lesion,¹⁰ lesion length,^{11,12} the presence of collateral vessels,^{13,14} diffuse disease of the distal bed, or the state of the microcirculation.^{15,16}

Although several studies have found poor correlations between the functional significance of a lesion obtained by visual assessment and by FFR,^{6–9,17} few studies have analyzed the factors associated with this discrepancy. The aim of the present study was to investigate the clinical and angiographic variables that could be associated with a greater degree of error when the functional significance of a coronary lesion is assessed by angiography.

METHODS

Design and Study Patients

We conducted a retrospective observational study of a historic cohort of patients referred to a cardiac catheterization laboratory for coronary angiography for suspected coronary disease between January 1, 2008 and May 31, 2012. The patients were evaluated for revascularization by measuring the FFR across an intermediate coronary lesion (40%–70% diameter stenosis by visual estimation) using a pressure wire. We excluded patients with lesions > 20% in a segment distal or proximal to the target lesion. The FFR was not measured in vessels that perfused akinetic or previously infarcted territory. In patients with acute coronary syndrome, the FFR was only measured in nonculprit vessels.

Procedure

All procedures were performed according to the usual protocol of the center conducting the study. After the decision was made to measure the FFR, the diagnostic catheter used for angiography was replaced by a 6-Fr guide catheter. This catheter was used to repeat the projections providing the best visualization of the lesions, with greater visual stenosis and without overlapping branches

or loss of length because of curvature. All patients received 100 IU/kg intravenous sodium heparin before the procedure if not previously administered. Functional evaluation was performed with a 0.014-inch intracoronary pressure wire (Pressure Wire, Certus or Airis, Radi Medical Systems; Uppsala, Sweden, or Prime-Wire Prestige Pressure Guide Wire, Volcano Corp.; San Diego, California, United States). The pressure wire was calibrated externally and then advanced to the distal end of the guide catheter while equalizing the pressures according to the system used to measure the FFR. After administration of 200 µg to 300 µg intracoronary nitroglycerin, the guide was advanced until the sensor was at least 20 mm distal to the lesion. We followed the standard procedure used in our hospital to obtain the FFR by administering 300 µg to 1200 µg intracoronary adenosine, while taking particular care to avoid wedging the catheter in the coronary ostium after bolus injection of the drug. The beat-to-beat ratio of the mean aortic pressure at the end of the guide catheter and the pressure distal to the lesion, obtained via the pressure wire under maximum hyperemia, were used to measure the FFR. We measured the FFR at least 3 times and used the lowest measurement. We successively administered 300 µg, 600 µg and 1200 µg intracoronary adenosine whenever the previous dose failed to produce a period of asystole ≥ 6 s.

The decision to revascularize was left to the operator's discretion based on the data obtained in the angiographic and functional study.

Angiographic Variables

In our hospital, the routine method to obtain the FFR includes obtaining at least 1 projection that provides the best visualization of the lesion using the guide catheter, after the administration of intracoronary nitroglycerin. The diagnostic angiographic sequences of each procedure were separated from those obtained during the intervention (when applicable). The observers were only provided with diagnostic images and were blinded to the result of the FFR study when performing the digital quantification of lesion stenosis. Data were collected on the following variables: severe calcification (multiple opacification visible in more than 1 projection covering the entire vessel lumen at the site of the lesion); bifurcation (presence of a > 15-mm side branch originating at the site of the lesion); angulation > 45° (target lesion in a segment with angulation > 45°); ostial location (lesion at the origin of the vessel in the aorta); perfused myocardial territory (Duke jeopardy score^{18,19}); and location of the lesion in the stent.

Digital quantification was performed using the QAngio XA version 7.1.43.0 postprocessing software package (Medis Medical Imaging Systems; Leiden, The Netherlands).

The lesions were analyzed by 2 experienced interventional cardiologists (more than 1000 coronary interventions using a pressure wire/measuring the FFR). One of them analyzed the lesions twice (OBS1A and OBS1B, with a 4-month interval between assessments), and the other analyzed them once (OBS2). They predicted whether the result of the pressure wire study was positive (FFR ≤ 0.80). The 2 observers were blinded to the previous predictions, the assessments made by the other, and the FFR results.

Based on their previous experience and published scientific evidence, the 2 observers took into account not only the degree of stenosis when making their predictions but also other parameters that have been associated with the FFR.¹⁶ In cases of disagreement, the mode of the 3 predictions was used. Overestimation was defined by a predicted FFR ≤ 0.80 vs a measured FFR > 0.80. Underestimation was defined by a predicted FFR > 0.80

vs a measured FFR ≤ 0.8 . Disagreement was defined by the underestimation or overestimation of the FFR. We analyzed the factors associated with disagreement, overestimation, or underestimation.

Statistical Analysis

All quantitative variables are expressed as mean \pm standard deviation and were compared using the Student *t* test. Qualitative variables are expressed as absolute values (percentages) and were compared using the chi-squared test. Logistic regression models were used to analyze the angiographic and clinical variables associated with an observed FFR ≤ 0.80 , an estimated FFR ≤ 0.80 , and disagreement (overestimation and underestimation) between the predicted and observed functional significance of the lesions. A P-value of $<.5$ was used as a cutoff for statistical significance. The Hosmer–Lemeshow test and the area under the ROC (receiver operating characteristic) curve were used to assess the model's goodness-of-fit and predictive power, respectively. All analyses were conducted using the SPSS 20.0 statistical package for Windows (SPSS Inc.; Chicago, Illinois, United States). Although this study compared 2 diagnostic tests, rather than assessing the reliability of 1 diagnostic test, the recommendations of the STARD initiative in data presentation were followed as closely as possible.²⁰

RESULTS

We studied 665 lesions in 587 patients. **Table 1** shows the clinical and angiographic characteristics of the patients. Coronary angiography was performed in 56% of the patients in the setting of acute coronary syndrome. In total, 58% of the lesions were located in the anterior descending coronary artery, 26% had severe calcification, and 9% were in-stent restenosis. An FFR of ≤ 0.80 was obtained in 29.9% of the lesions (199).

Table 2 shows the distribution of the clinical and angiographic variables according to the FFR value. After adjustment for other variables, age, male sex, absence of acute coronary syndrome, the presence of calcification, target segment with an angulation $> 45^\circ$, anterior descending coronary artery location, a greater degree of stenosis, greater length, smaller reference diameter, and greater territory distal to the target lesion were associated with FFR ≤ 0.80 (**Table 2**, right).

In relation to the predicted FFR value ≤ 0.80 , intraobserver agreement was 83% ($\kappa = 0.54 \pm 0.039$; $P < .0005$) and interobserver agreement was 79.4% ($\kappa = 0.43 \pm 0.041$; $P < .0005$).

Agreement between the FFR value estimated by the observers (the statistical mode of the estimates made by OBS1A, OBS1B, and OBS2) and the measured FFR value was 69.9% (overestimation, 11.3%; underestimation, 18.8%). Visual estimation of FFR ≤ 0.80 had a sensitivity of 37.2% (95% confidence interval [95%CI], 0.30–0.44), specificity of 83.9% (95%CI, 0.81–0.87), positive predictive value of 49.7% (95%CI, 0.42–0.58), and negative predictive value of 75.8% (95%CI, 0.72–0.79). Observer agreement (OBS1A, OBS1B, and OBS2) on FFR ≤ 0.80 was 67.1%, 67.0%, and 70.1%, respectively.

Table 3 shows the distribution of the clinical and angiographic variables according to the functional significance estimated by the observers.

Univariable analysis showed that the degree of angiographic stenosis, a smaller reference diameter, calcification, angle of the bifurcation, and an ostial location were associated with an estimated FFR ≤ 0.80 . In the unadjusted analysis, the Duke jeopardy score, paradoxically, showed an inverse association with a higher estimated FFR ≤ 0.80 (Duke jeopardy score, 3.60

Table 1

Baseline Characteristics of the Patients

Clinical characteristics of the patients (n = 587)	
Age, y	67.9 \pm 10.6
Women	162 (27.6)
Hypertension	438 (74.6)
Diabetes mellitus	245 (41.7)
Smoking	279 (47.5)
Hyperlipidemia	316 (53.8)
Previous infarction	132 (22.5)
Previous coronary revascularization	158 (26.9)
Percutaneous	138 (23.5)
Surgical	6 (1.1)
Prior stroke	46 (7.8)
Indication for coronary angiography	
STEACS	46 (7.8)
NSTEACS	287 (48.9)
Stable angina/silent ischemia	254 (43.3)
Angiographic characteristics of the lesions (n = 665)	
Target lesion	
Anterior descending	386 (58.0)
Circumflex	135 (20.3)
Right coronary	136 (20.5)
Left main coronary	7 (1.1)
Angulation $> 45^\circ$	37 (5.6)
Calcification	170 (25.6)
Ostium	58 (8.7)
Bifurcation	35 (17.6)
Duke jeopardy score	3.54 \pm 1.50
Location in the stent	62 (9.3)
Stenosis, %	50.3 \pm 9.7
Length, mm	14.0 \pm 7.2
Reference diameter, mm	2.7 \pm 0.6
FFR value	0.83 \pm 0.09
FFR ≤ 0.80	199 (29.9)

FFR, fractional flow reserve; NSTEACS, non–ST-segment elevation acute coronary syndrome; STEACS, ST-segment elevation acute coronary syndrome. Values are expressed as No. (%) or mean \pm standard deviation.

in lesions with an expected FFR > 0.80 vs 3.32 with and estimated FFR ≤ 0.80 ; $P = 0.047$).

Functional significance was overestimated in 75 lesions (16% of which had an FFR > 0.80) and underestimated in 125 lesions (62.8% of which had an FFR ≤ 0.80). **Table 4** shows the distribution of the variables according to agreement, underestimation, or overestimation. Severe calcification, lesion length, and lesion location in the right coronary artery were associated with underestimating the significance of the lesion. Location of the lesion in the stent, ostium, or bifurcation, and angulation were associated with overestimation.

Multivariable analysis showed that severe angiographic calcification, angulation $> 45^\circ$ in the target segment, bifurcation location, greater length, greater diameter stenosis, and a smaller reference diameter were associated with estimation of the functional significance of the lesion (**Table 5**).

After adjustment for other variables, the variables independently associated with incorrectly classifying the functional significance of the lesion by visual estimation were stent location, calcification, and the size of the coronary territory distal to target lesion (**Table 5**).

Table 2
Univariable Analysis of the Observed Fractional Flow Reserve ≤ 0.80 or > 0.80 and Logistic Regression Analysis

	Univariable analysis			Multivariable Analysis	
	FFR > 0.80 (n=466)	FFR ≤ 0.80 (n=199)	P	OR (95%CI)	P
Age, y	68.4 \pm 10.8	67.0 \pm 10.1	.06	0.97 (0.96-0.99)	.005
Women	151 (32.4)	35 (17.6)	<.0005	0.48 (0.30-0.77)	.002
Diabetes mellitus	193 (41.6)	81 (41.1)	.91	0.96 (0.66-1.47)	.96
Hypertension	346 (73.9)	154 (74.4)	.89	1.23 (0.77-1.96)	.37
ACS	281 (60.3)	97 (48.7)	.006	0.47 (0.32-0.70)	<.0005
Anterior descending coronary	231 (49.6)	155 (77.9)	<.0005	2.13 (1.30-3.49)	.003
Angulation $> 45^\circ$	21 (4.5)	16 (8.0)	.069	1.89 (0.83-4.28)	.13
Calcification	85 (18.2)	85 (42.7)	<.0005	2.18 (1.40-3.40)	<.001
Ostium	44 (9.4)	14 (7.0)	.31	1.18 (0.42-3.34)	.75
Bifurcation	58 (12.4)	35 (17.6)	.08	1.15 (0.67-1.96)	.62
Duke jeopardy score (by unit)	3.4 \pm 1.4	4.0 \pm 1.7	<.0005	1.28 (1.10-1.49)	<.002
Location in the stent	42 (9.0)	20 (10.1)	.67	0.99 (0.52-1.88)	.98
Length, mm	12.7 \pm 5.1	16.9 \pm 10.0	<.0005	1.08 (1.05-1.11)	<.0005
Stenosis, %	49.4 \pm 9.3	52.5 \pm 10.2	<.0005	1.04 (1.02-1.06)	<.0005
Reference diameter, mm	2.8 \pm 0.6	2.6 \pm 0.6	.001	0.58 (0.40-0.85)	.005

95%CI, 95% confidence interval; ACS, acute coronary syndrome; FFR, fractional flow reserve; OR, odds ratio.

Values are expressed as No. (%) or mean \pm standard deviation. Hosmer-Lemeshow test: chi-squared = 7.03; degrees of freedom, 8; significance, 0.53. Area under the receiver operating characteristic curve, 0.81 (range, 0.77-0.84; $P < .0005$)

Table 3
Clinical and Angiographic Characteristics of the Lesions According to the Estimated Fractional Flow Reserve

	FFR > 0.80 (n=516)	FFR ≤ 0.80 (n=149)	P
Age, y	67.9 \pm 10.5	67.8 \pm 10.9	.96
Women	149 (28.9)	37 (24.8)	.33
Diabetes mellitus	213 (41.4)	61 (41.5)	.99
Hypertension	387 (75.3)	103 (70.1)	.20
ACS	298 (57.8)	80 (53.7)	.38
<i>Target artery</i>			
Anterior descending	299 (57.9)	87 (58.4)	.92
Circumflex	99 (19.2)	36 (24.2)	.18
Right coronary	110 (21.3)	26 (17.4)	.30
Left main coronary	2 (0.4)	5 (2.5)	.016
<i>Characteristics of the lesion</i>			
Angulation $> 45^\circ$	22 (4.3)	15 (10.1)	.006
Calcification	113 (21.9)	57 (38.3)	<.0005
Ostium	34 (6.6)	24 (16.1)	<.0005
Bifurcation	54 (10.5)	39 (26.2)	<.0005
Duke jeopardy score	3.60 \pm 1.54	3.32 \pm 1.30	.047
Location in the stent	54 (10.5)	8 (5.4)	.059
Length, mm	13.5 \pm 6.7	15.5 \pm 8.2	<.002
Stenosis, %	49.6 \pm 9.6	52.9 \pm 9.6	<.0005
Reference diameter, mm	2.8 \pm 0.6	2.5 \pm 0.5	<.0005

ACS, acute coronary syndrome; FFR, fractional flow reserve. Values are expressed as No. (%) or mean \pm standard deviation.

The variables associated with overestimating the functional significance of the lesions were location in the stent, in an artery other than the anterior descending artery, and in a bifurcation (Table 5).

Male sex, a larger myocardial territory perfused by the vessel with the lesion, and severe calcification were significantly and independently associated with underestimating the functional significance of the lesion by visual estimation (Table 5).

DISCUSSION

Previous studies have shown a weak correlation between the angiographic quantification (visual or automated) of coronary lesions and their functional significance.^{6,9} This study is the first to analyze the variables associated with errors (both overestimation and underestimation) in the visual assessment of the functional significance of coronary lesions.

Table 4

Clinical and Angiographic Characteristics of the Lesions According to Agreement on the Functional Significance of the Lesion Estimated by Visual Assessment and Observed Fractional Flow Reserve

	Agreement (n=465)	Disagreement (n=200)	P	Overestimation (n=75)	P	Underestimation (n=125)	P
Age, y	67.7 ± 10.8	68.0 ± 10.1	.76	70.0 ± 10.8	.17	67.8 ± 10.8	.52
Women	140 (30.1)	46 (23.0)	.06	24 (32.0)	.73	22 (17.6)	.006
Diabetes mellitus	188 (40.8)	86 (43.0)	.59	33 (44.0)	.59	53 (42.4)	.73
Hypertension	338 (73.3)	152 (76.0)	.47	54 (72.0)	.80	98 (78.4)	.25
ACS	263 (56.6)	115 (57.5)	.82	49 (65.3)	.15	66 (52.8)	.47
Anterior descending	256 (55.1)	130 (65.0)	.02	31 (41.3)	.028	99 (79.2)	< .005
Circumflex	96 (20.6)	39 (19.5)	.73	30 (40.0)	< .005	9 (7.2)	< .005
Right coronary	110 (23.7)	26 (13.0)	.02	14 (18.7)	.35	12 (9.6)	.001
Angulation > 45°	20 (4.3)	17 (8.5)	.03	8 (10.7)	.021	9 (7.2)	.18
Calcification	98 (21.1)	72 (36.0)	< .0005	22 (29.3)	.108	50 (40.0)	< .0005
Ostium	19 (4.1)	11 (5.5)	.42	7 (9.3)	.04	4 (3.2)	.65
Bifurcation	55 (11.8)	38 (19.0)	.01	21 (28.0)	< .005	17 (13.6)	.59
Duke jeopardy score	3.45 ± 1.38	3.75 ± 1.71	.02	3.01 ± 1.20	.01	4.32 ± 1.97	< .0005
Location in the stent	46 (9.9)	16 (8.0)	.44	2 (2.79)	.042	14 (11.2)	.66
Length, mm	13.6 ± 6.3	14.7 ± 8.9	.12	12.7 ± 5.5	.2	15.9 ± 10.2	.002
Stenosis, %	49.8 ± 9.7	51.4 ± 9.6	.05	51.4 ± 8.9	.19	51.4 ± 9.9	.12
Reference diameter, mm	2.70 ± 0.59	2.60 ± 0.60	.01	2.60 ± 0.59	.007	2.70 ± 0.60	.09

ACS, acute coronary syndrome; FFR, fractional flow reserve.

Values are expressed as No. (%) or mean ± standard deviation.

The specialists conducting the visual estimation are not only experienced in interventional cardiology, but also have specific experience in the use of FFR for making decisions and in taking into account the degree of stenosis of the lesion and other factors that determine the functional significance of coronary lesions. This study shows that even interventionists who are experienced in the invasive functional assessment of lesions can fail to visually predict the functional significance of lesions.

Overestimation of the functional significance of lesions was associated with the location of the lesion in an artery other than the anterior descending artery, in the stent, and in a bifurcation. Underestimation was associated with male sex, severe calcification, and greater coronary area distal to the lesion (Figure).

Angiographic-functional Agreement

The results of this study are consistent with those obtained in previous studies regarding low intraobserver and interobserver agreement in the visual assessment of the severity of intermediate coronary lesions. Matching values were higher than those of other studies based solely on the degree of stenosis.^{6,8} Despite being members of the same intervention team and using similar criteria for the estimations, $\kappa < 0.6$ was obtained, which indicates the low reliability of the visual assessment of lesions.

Invasive measurement of FFR addresses characteristics of the coronary lesion apart from the degree of stenosis alone. Frequent use of the FFR in the cardiac catheterization laboratory may have an “instructive” effect that would lead the interventionist to modify the assessment of the significance of the lesion by taking into account aspects such as the lesion length, disease of the distal bed, or the size of the myocardium distal to the lesion, in addition to its degree of stenosis. In a recent study, Park et al.⁹ used the degree of angiographic stenosis (52% diameter stenosis) as the only classification parameter. They found an agreement of 66% with a sensitivity of 66%, a specificity of 67%, and positive and negative predictive values of 48% and 81%, respectively.⁹ We used multiple clinical and angiographic parameters to visually

estimate the functional significance of the lesion and observed a slightly higher agreement (70%), lower sensitivity, and greater specificity.

The percentage of lesions with an FFR > 0.80 considered significant by visual estimation was significantly lower than that observed by Park et al. (16% vs 57%). In contrast, a much higher percentage of lesions with an FFR ≤ 0.80 were visually classified as functionally nonsignificant (63% vs 16%). The frequent use of the FFR in intermediate lesions probably leads to the application of a “functional approach” to their assessment, without invasive measurement, which would lead to a more conservative assessment when considering a lesion functionally significant.

Angiographic Characteristics Associated With Visually Overestimating a Lesion

Together with the inability of angiography to predict the functional significance of coronary lesions, our study shows that certain angiographic characteristics of the lesion may be associated with underestimating or overestimating the significance of the stenosis. If these angiographic characteristics are present in an angiographically intermediate stenosis, the probability of error when decisions are made on whether to revascularize without a functional study can be even greater than in the general population.

In our study, moderate in-stent restenosis (40%-70% stenosis) was the factor most strongly associated with overestimating the significance of the lesion. The presence of bare-metal stents makes it difficult to determine the edges of the vessel lumen, which can lead to underestimation of its diameter and overestimation of the degree of restenosis. Although previous studies have demonstrated the limitations of angiography in assessing in-stent restenosis,^{21,22} recent clinical trials have continued to use the degree of angiographic stenosis to determine restenosis or to decide whether to revascularize the treated lesion.^{23–25} The use of FFR for the functional assessment of the lesion during follow-up in these

Table 5
Logistic Regression Analysis of the Variables Associated with a Predicted Fractional Flow Reserve ≤ 0.80 and Disagreement (Underestimation or Overestimation) on the Observed Fractional Flow Reserve

	OR (95%CI)	P
<i>Independent variables associated with a predicted FFR ≤ 0.80</i>		
Reference diameter, mm	0.43 (0.28-0.65)	<.0005
Stenosis (by unit)	1.04 (1.01-1.06)	.002
Length, mm	1.03 (1.00-1.06)	.028
Angulation $> 45^\circ$	2.67 (1.25-5.72)	.011
Calcification	1.58 (1.02-2.51)	.045
Bifurcation	2.94 (1.78-4.85)	<.0005
Location in the stent	0.51 (0.23-1.54)	.09
Hosmer-Lemeshow test: chi-squared=7.03; degrees of freedom, 8; P=.53 Area under the ROC curve, 0.81 (range, 0.77-0.84; P<.0005)		
<i>Variables independently associated with the discrepancy between a predicted and observed FFR ≤ 0.80</i>		
Location in the stent	2.33 (1.29-4.19)	.005
Calcification	2.07 (1.32-3.25)	.002
Duke jeopardy score	1.25 (1.01-1.46)	.003
Hosmer-Lemeshow test: chi-squared=8.1; degrees of freedom, 8; P=.42 Area under the ROC curve 0.75 (range 0.70-0.78; P<.0005)		
<i>Variables independently associated with overestimating the severity of the stenosis (estimated FFR ≤ 0.80 and measured FFR > 0.80)</i>		
Location in the stent	2.67 (1.20-5.92)	.016
Artery other than the anterior descending	2.09 (1.05-4.17)	.036
Bifurcation	2.33 (1.14-4.76)	.02
Hosmer-Lemeshow test: chi-squared=4.6; degrees of freedom, 8; P=.80 Area under the ROC curve, 0.70 (range, 0.65-0.74; P<.0001)		
<i>Variables independently associated with underestimating the severity of the stenosis (estimated FFR > 0.80 and measured FFR ≤ 0.80)</i>		
Male sex	1.93 (1.02-3.64)	.042
Calcification	2.05 (1.19-3.53)	.010
Duke jeopardy score (by unit)	1.36 (1.15-1.61)	<.0005
Hosmer-Lemeshow test: chi-squared=13.6; degrees of freedom, 8; P=.1 Area under the ROC curve, 0.75 (range, 0.70-0.79; P<.0005)		

95%CI, 95% confidence interval; FFR, fractional flow reserve; OR, odds ratio; ROC, receiver operating characteristic.

In all cases, adjustments were made by age, sex, hypertension, diabetes mellitus, acute coronary syndrome, anterior descending artery location, reference diameter, lesion length, degree of stenosis, stent location, bifurcation location, ostial location, angulation $> 45^\circ$, calcification, and Duke jeopardy score.

studies could have led to fewer restenosis events being assigned to patients undergoing percutaneous coronary intervention.

The location of the lesion in an artery other than the anterior descending artery was one of the variables associated with overestimating the functional significance of the lesion. In the multivariable analysis, an adjustment was made for the myocardial territory perfused by the artery with the lesion. This adjustment may have been insufficient, such that the observed overestimation could have been due to a smaller myocardial area in lesions in locations other than the anterior descending artery with a greater FFR value for the same diameter stenosis value. This possibility is consistent with the finding of an association between a larger myocardial territory distal to the lesion and underestimating the functional significance of the lesion. Although the researchers were aware of the effect of perfused myocardial territory on the FFR value for a given lesion, the angiographic characteristics of the lesion continued to take precedence over the other characteristics when estimates were made on its functional significance. The location of the lesion in a bifurcation was also associated with the functional




Figure. Percentages of overestimation, underestimation, and agreement between visual estimation of the functional significance of the coronary lesion (fractional flow reserve ≤ 0.80) and its observed value according to the clinical and angiographic characteristics.

overestimation of the stenosis. Several studies have demonstrated the difficulty of assessing these lesions angiographically and have shown that less functional significance is found when the FFR is used than might be expected by the degree of stenosis observed both before and after treatment of the lesion.^{26,27}

Angiographic Characteristics Associated With Visually Underestimating a Lesion

Male sex and severe calcification were associated with underestimation of the functional significance of lesions. As the researchers were blinded to the sex of the patients when assessing the lesions, they could not have taken this variable into account in their estimates. Our findings are in line with those of several studies that have found an association between female sex and a greater FFR value for the same degree of stenosis in men and women.²⁸⁻³⁰

Calcification hampers the visual and automated angiographic assessment of stenosis. In this study, in contrast to in-stent restenosis, calcification was associated with underestimating the significance of the lesion. Increased radiodensity due to increased vessel thickness can contribute to overestimating lumen diameter, which would lead to systematic errors in visual assessment. Moreover, coronary calcification can be a marker of more extensive and diffuse disease and could lead to diseased coronary segments being used as a reference, and thus lead to underestimation of the degree of stenosis or lesion length.

Limitations

This study was conducted at a single hospital. Although the researchers who visually assessed the lesions were blind to the results of the FFR study and to each other's estimates, local factors may have influenced estimation of the functional significance of the lesions. The generalizability of our findings is supported by the low interobserver agreement found in this study, which is similar to that found in other published studies, and agreement with the literature regarding the variables associated with the predictions. The use of the mode in the 3 estimates gave more weight to the observations of OBS1, who presented slightly worse agreement.

CONCLUSIONS

Visual estimation of the functional impact of intermediate coronary stenosis (40%-70%), even when taking into account factors other than the degree of stenosis, can lead to more than 30% disagreement with the results of invasive measurement of FFR using a cutoff of 0.80. Calcification, male sex, or lesions located in arteries perfusing large myocardial territories can lead to underestimating the functional significance of the lesion. The functional significance of lesions in vessels other than the left anterior descending artery, in-stent restenosis, and lesions in bifurcations tends to be overestimated by visual estimation. The invasive functional study of lesions with these characteristics is strongly indicated to avoid erroneous decisions to revascularize.

CONFLICTS OF INTEREST

None declared.

REFERENCES

- Frye RL, August P, Brooks MM, Hardison RM, Kelsey SF, MacGregor JM, et al. A randomized trial of therapies for type 2 diabetes and coronary artery disease. *N Engl J Med.* 2009;360:2503–15.
- Boden WE, O'Rourke RA, Teo KK, Hartigan PM, Maron DJ, Kostuk WJ, et al. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med.* 2007;356:1503–16.
- Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, et al. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med.* 2009;360:961–72.
- Wijns W, Kolh P, Danchin N, Di MC, Falk V, Folliguet T, et al. Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J.* 2010;31:2501–55.
- Levine GN, Bates ER, Blankenship JC, Bailey SR, Bittl JA, Cercek B, et al. ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *J Am Coll Cardiol.* 2011;58:e44–122.
- Fischer JJ, Samady H, McPherson JA, Sarembock IJ, Powers ER, Gimble LW, et al. Comparison between visual assessment and quantitative angiography versus fractional flow reserve for native coronary narrowings of moderate severity. *Am J Cardiol.* 2002;90:210–5.
- Christou MA, Siontis GC, Katritsis DG, Ioannidis JP. Meta-analysis of fractional flow reserve versus quantitative coronary angiography and noninvasive imaging for evaluation of myocardial ischemia. *Am J Cardiol.* 2007;99:450–6.
- Lindstaedt M, Spiecker M, Perings C, Lawo T, Yazar A, Holland-Letz T, et al. How good are experienced interventional cardiologists at predicting the functional significance of intermediate or equivocal left main coronary artery stenoses? *Int J Cardiol.* 2007;120:254–61.
- Park SJ, Kang SJ, Ahn JM, Shim EB, Kim YT, Yun SC, et al. Visual-functional mismatch between coronary angiography and fractional flow reserve. *JACC Cardiovasc Interv.* 2012;5:1029–36.
- Leone AM, De Caterina AR, Basile E, Gardi A, Laezza D, Mazzari MA, et al. Influence of the amount of myocardium subtended by a stenosis on fractional flow reserve. *Circ Cardiovasc Interv.* 2013;6:29–36.
- López-Palop R, Carrillo P, Cordero A, Frutos A, Mateo I, Mashlab S, et al. Effect of lesion length on functional significance of intermediate long coronary lesions. *Cathet Cardiovasc Intervent.* 2013;81:e186–94.
- Iguchi T, Hasegawa T, Nishimura S, Nakata S, Kataoka T, Ehara S, et al. Impact of lesion length on functional significance in intermediate coronary lesions. *Clin Cardiol.* 2013;36:172–7.
- Sachdeva R, Uretsky BF. The effect of CTO recanalization on FFR of the donor artery. *Catheter Cardiovasc Interv.* 2011;77:367–9.
- Melikian N, Cuisset T, Hamilos M, De BB. Fractional flow reserve—the influence of the collateral circulation. *Int J Cardiol.* 2009;132:e109–10.
- Rodes-Cabau J, Gutierrez M, Courtis J, Larose E, Dery JP, Cote M, et al. Importance of diffuse atherosclerosis in the functional evaluation of coronary stenosis in the proximal-mid segment of a coronary artery by myocardial fractional flow reserve measurements. *Am J Cardiol.* 2011;108:483–90.
- Pijls NHJ, Sels JW. Functional measurement of coronary stenosis. *J Am Coll Cardiol.* 2012;59:1045–57.
- Tonino PA, Fearon WF, De BB, Oldroyd KG, Leeser MA, Ver Lee PN, et al. Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol.* 2010;55:2816–21.
- Dash H, Johnson RA, Dinsmore RE, Harthorne JW. Cardiomyopathic syndrome due to coronary artery disease. I: Relation to angiographic extent of coronary disease and to remote myocardial infarction. *Br Heart J.* 1977;39:733–9.
- Califf RM, Phillips III HR, Hindman MC, Mark DB, Lee KL, Behar VS, et al. Prognostic value of a coronary artery jeopardy score. *J Am Coll Cardiol.* 1985;5:1055–63.
- Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. *BMJ.* 2003;326:41–4.
- López-Palop R, Pinar E, Lozano I, Saura D, Pico F, Valdes M. Utility of the fractional flow reserve in the evaluation of angiographically moderate in-stent restenosis. *Eur Heart J.* 2004;25:2040–7.
- Nam CW, Rha SW, Koo BK, Doh JH, Chung WY, Yoon MH, et al. Usefulness of coronary pressure measurement for functional evaluation of drug-eluting stent restenosis. *Am J Cardiol.* 2011;107:1783–6.
- Haude M, Lee SW, Worthley SG, Silber S, Verheye S, Erbs S, et al. The REMEDEE trial: a randomized comparison of a combination sirolimus-eluting endothelial progenitor cell capture stent with a paclitaxel-eluting stent. *JACC Cardiovasc Interv.* 2013;6:334–43.
- Deftereos S, Giannopoulos G, Raisakis K, Kossyvakis C, Kaoukis A, Panagopoulou V, et al. Colchicine treatment for the prevention of bare-metal stent restenosis in diabetic patients. *J Am Coll Cardiol.* 2013;61:1679–85.
- Byrne RA, Neumann FJ, Mehilli J, Piniček S, Wolff B, Tiroch K, et al. Paclitaxel-eluting balloons, paclitaxel-eluting stents, and balloon angioplasty in patients with restenosis after implantation of a drug-eluting stent (ISAR-DESIRE 3): a randomised, open-label trial. *Lancet.* 2013;381:461–7.
- Ahn JM, Lee JY, Kang SJ, Kim YH, Song HG, Oh JH, et al. Functional assessment of jailed side branches in coronary bifurcation lesions using fractional flow reserve. *JACC Cardiovasc Interv.* 2012;5:155–61.
- Girasis C, Onuma Y, Schuurbiens JCH, Morel Ma, Van Es GA, Van Geuns RJ, et al. Validity and variability in visual assessment of stenosis severity in phantom bifurcation lesions: A survey in experts during the fifth meeting of the European bifurcation club. *Cathet Cardiovasc Intervent.* 2012;79:361–8.
- Fineschi M, Guerrieri G, Orphal D, Palmerini E, Munzel T, Warnholtz A, et al. The impact of gender on fractional flow reserve measurements. *EuroIntervention.* 2013;9:360–6.
- Kim HS, Tonino PA, De BB, Yong AS, Tremmel JA, Pijls NH, et al. The impact of sex differences on fractional flow reserve-guided percutaneous coronary intervention: A FAME (Fractional Flow Reserve Versus Angiography for Multivessel Evaluation) substudy. *JACC Cardiovasc Interv.* 2012;5:1037–42.
- Kang SJ, Ahn JM, Han S, Lee JY, Kim WJ, Park DW, et al. Sex differences in the visual-functional mismatch between coronary angiography or intravascular ultrasound versus fractional flow reserve. *JACC Cardiovasc Interv.* 2013;6:562–8.