

Original article

# Peak Exercise Oxygen Uptake Predicts Recurrent Admissions in Heart Failure With Preserved Ejection Fraction



Patricia Palau,<sup>a,\*</sup> Eloy Domínguez,<sup>a</sup> Eduardo Núñez,<sup>b</sup> José María Ramón,<sup>b</sup> Laura López,<sup>c</sup> Joana Melero,<sup>a</sup> Juan Sanchis,<sup>b,d</sup> Alejandro Bellver,<sup>a</sup> Enrique Santas,<sup>b</sup> Antoni Bayes-Genis,<sup>d,e</sup> Francisco J. Chorro,<sup>b</sup> and Julio Núñez<sup>b,d</sup>

<sup>a</sup> Servicio de Cardiología, Hospital General Universitario de Castellón, Universitat Jaume I, Castellón, Spain

<sup>b</sup> Servicio de Cardiología, Hospital Clínico Universitario, INCLIVA, Universitat de València, Valencia, Spain

<sup>c</sup> Departamento de Fisioterapia, Universitat de València, Valencia, Spain

<sup>d</sup> CIBER Cardiovascular, Madrid, Spain

<sup>e</sup> Servicio de Cardiología, Hospital Universitari Germans Trias i Pujol, Badalona, Departamento de Medicina, Universitat Autònoma de Barcelona, Barcelona, Spain

## Article history:

Received 11 January 2017

Accepted 8 May 2017

Available online 27 June 2017

## Keywords:

Heart failure with preserved ejection fraction

Prognosis

Recurrent hospitalizations

Exercise capacity

## ABSTRACT

**Introduction and objectives:** Heart failure with preserved ejection fraction (HFpEF) is a highly prevalent syndrome with an elevated risk of morbidity and mortality. To date, there is scarce evidence on the role of peak exercise oxygen uptake (peak  $\text{VO}_2$ ) for predicting the morbidity burden in HFpEF. We sought to evaluate the association between peak  $\text{VO}_2$  and the risk of recurrent hospitalizations in patients with HFpEF.

**Methods:** A total of 74 stable symptomatic patients with HFpEF underwent a cardiopulmonary exercise test between June 2012 and May 2016. A negative binomial regression method was used to determine the association between the percentage of predicted peak  $\text{VO}_2$  (pp-peak  $\text{VO}_2$ ) and recurrent hospitalizations. Risk estimates are reported as incidence rate ratios.

**Results:** The mean age was  $72.5 \pm 9.1$  years, 53% were women, and all patients were in New York Heart Association functional class II to III. Mean peak  $\text{VO}_2$  and median pp-peak  $\text{VO}_2$  were  $10 \pm 2.8$  mL/min/kg and 60% (range, 47–67), respectively. During a median follow-up of 276 days [interquartile range, 153–1231], 84 all-cause hospitalizations in 31 patients (41.9%) were registered. A total of 15 (20.3%) deaths were also recorded. On multivariate analysis, accounting for mortality as a terminal event, pp-peak  $\text{VO}_2$  was independently and linearly associated with the risk of recurrent admission. Thus, and modeled as continuous, a 10% decrease of pp-peak  $\text{VO}_2$  increased the risk of recurrent hospitalizations by 32% (IRR, 1.32; 95%CI, 1.03–1.68;  $P = .028$ ).

**Conclusions:** In symptomatic elderly patients with HFpEF, pp-peak  $\text{VO}_2$  predicts all-cause recurrent admission.

© 2017 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

## El consumo máximo de oxígeno predice los ingresos recurrentes por insuficiencia cardiaca con fracción de eyección conservada

## RESUMEN

**Introducción y objetivos:** La insuficiencia cardiaca con fracción de eyección conservada (IC-FEc) es un síndrome muy prevalente con alto riesgo de morbilidad y mortalidad. Hasta la fecha, la evidencia acerca del papel del consumo máximo de oxígeno ( $\text{VO}_{2\text{máx}}$ ) para predecir la carga de morbilidad en la IC-FEc es escasa. El objetivo de este estudio es evaluar la relación entre el  $\text{VO}_{2\text{máx}}$  y el riesgo de ingresos recurrentes de los pacientes con IC-FEc.

**Métodos:** A un total de 74 pacientes con IC-FEc sintomáticos y clínicamente estables, se les realizó una prueba de esfuerzo cardiopulmonar entre junio de 2012 y mayo de 2016. Se utilizó el método de regresión binomial negativa para determinar la asociación entre el porcentaje de  $\text{VO}_{2\text{máx}}$  predicho ( $\% \text{VO}_{2\text{máx-p}}$ ) y los ingresos recurrentes. Las estimaciones del riesgo se informaron como tasas de incidencia.

**Resultados:** La media de edad era  $72,5 \pm 9,1$  años, el 53% eran mujeres y todos los pacientes estaban en clase funcional II-III de la *New York Heart Association*. La media de  $\text{VO}_{2\text{máx}}$  y la mediana de  $\% \text{VO}_{2\text{máx-p}}$  fueron  $10 \pm 2,8$  ml/min/kg y el 60% (47-67) respectivamente. Durante un seguimiento medio de 276 [intervalo intercuartílico, 153-1.231] días, se registraron 84 hospitalizaciones por cualquier causa de 31 pacientes (41,9%). También se determinó un total de 15 muertes (20,3%). En un análisis multivariable, teniendo en

## Palabras clave:

Insuficiencia cardiaca con fracción de eyección conservada

Pronóstico

Ingresos recurrentes

Capacidad funcional

## SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2017.08.012>, *Rev Esp Cardiol.* 2018;71:237-239.

\* Corresponding author: Servicio de Cardiología, Hospital General Universitario de Castellón, Avda. Benicasim s/n, 12004 Castellón, Spain.  
E-mail address: [patri.palau@gmail.com](mailto:patri.palau@gmail.com) (P. Palau).

<http://dx.doi.org/10.1016/j.rec.2017.05.022>

1885-5857/© 2017 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

cuenta la mortalidad como evento terminal, el %VO<sub>2máx-p</sub> mantuvo la asociación independiente y lineal con el riesgo de ingresos recurrentes. Así, y modelado como continuo, una disminución del 10% del %VO<sub>2máx-p</sub> aumentó en un 32% el riesgo de ingresos recurrentes (IRR = 1,32; IC95%, 1,03–1,68; p = 0,028).

**Conclusiones:** En los pacientes de edad avanzada con IC-FEC sintomáticos, el %VO<sub>2máx-p</sub> predice los ingresos recurrentes por todas las causas.

© 2017 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

### Abbreviations

HF: heart failure  
 HFpEF: heart failure with preserved ejection fraction  
 Peak VO<sub>2</sub>: peak exercise oxygen uptake  
 pp-peak VO<sub>2</sub>: percentage of predicted peak exercise oxygen uptake

### INTRODUCTION

Heart failure (HF) is a major public health problem with high associated morbidity and mortality worldwide.<sup>1</sup> More than 50% of patients with HF have preserved left ventricular ejection fraction (HFpEF), which is especially common in elderly people, women, and highly comorbid patients.<sup>2–4</sup> Risk prediction of repeat hospitalizations in HF has often been overlooked and most of the studies endorsing risk prediction of hospitalizations have focused on time-to-first event analysis, ignoring the clinical impact of recurrent hospitalization that frequently occurs in HF.<sup>5</sup> Recent initiatives advocate the inclusion of recurrent hospitalizations in risk stratification<sup>6</sup> to transmit a more realistic picture of the disease burden.

Although the clinical value of peak exercise oxygen uptake (peak VO<sub>2</sub>) in HF with reduced ejection fraction is well-documented,<sup>7,8</sup> the evidence endorsing its prognostic role in HFpEF is scarce<sup>7,9–11</sup> and even absent, especially regarding the risk of recurrent admissions.

The purpose of this study was to evaluate whether the percentage of predicted peak VO<sub>2</sub> (pp-peak VO<sub>2</sub>) is associated with the risk of recurrent admissions in elderly patients with HFpEF.

### METHODS

#### Study Design and Patients

This was a prospective study that included patients with a diagnosis of HFpEF according to the criteria of the European Society of Cardiology<sup>12</sup> and New York Heart Association functional class II–III/IV between 2 periods: June 2012 to May 2013 and June 2015 to May 2016. The study was conducted in a single third-level center in Spain. All patients provided signed informed consent before participation. The protocol was approved by the research ethics committee of our center in accordance with the principles of the Declaration of Helsinki and national regulations.

Candidate patients were selected from the outpatient HF unit.<sup>13</sup> All patients met the following inclusion criteria: a) previous history of symptomatic HF (New York Heart Association functional class ≥ II); b) normal left ventricular ejection fraction (ejection fraction > 0.50 by the Simpson method and end-diastolic diameter < 60 mm); c) structural heart disease (left ventricle hypertrophy/left atrial enlargement) and/or diastolic dysfunction estimated by 2-dimensional echocardiography; d) previous admission for acute HF; and e) clinical stability, without hospital admissions in the past

3 months. Patients were excluded if they could not perform a valid baseline exercise test or showed any previous medical condition such as: unstable angina, myocardial infarction or cardiac surgery within the previous 3 months; chronic metabolic, orthopedic, infectious disease or pulmonary disease (including pulmonary arterial hypertension, chronic thromboembolic pulmonary disease or chronic obstructive pulmonary disease); steroid, hormone, or cancer therapy; acute HF decompensation; any other comorbidity with a life expectancy of less than 1 year. The patient flowchart is depicted in [Figure 1](#).

After signing the informed consent form, a comprehensive medical history, physical examination, anthropometry and examination tests were performed by 2 trained cardiologists.

#### Procedures

Study procedures included electrocardiogram, echocardiography, cardiopulmonary exercise testing, and blood samples for a panel of baseline biomarkers. All of them were performed on the same day.

#### Cardiopulmonary Exercise Testing

Maximal functional capacity was evaluated with an incremental and symptom-limited cardiopulmonary exercise test (CORTEX Metamax 3B) on a bicycle ergometer, beginning with a workload of 10 W and increasing stepwise at 10 W increments every 1 minute. During exercise, patients were continuously monitored with a 12-lead electrocardiogram and blood pressure measurements every 2 minutes. Gas exchange data and cardiopulmonary variables were averaged every 10 seconds. Peak VO<sub>2</sub> was considered the highest value of VO<sub>2</sub> during the last 20 seconds of exercise and pp-peak VO<sub>2</sub> was calculated using the Wasserman equation.<sup>14</sup> Exercise ventilatory efficiency (VE/VCO<sub>2</sub> slope) was determined by measuring the slope across the entire course of the exercise.<sup>15</sup>

#### Echocardiography


Doppler echocardiogram examinations were performed under resting conditions using 2-dimensional echocardiography (iE33, Philips). All parameters, including tissue Doppler parameters, were measured according to the current guidelines of the European Society of Echocardiography.<sup>16</sup>

#### Biomarkers

All blood samples were obtained between 09:00 a.m. and 12:00 p.m. All biomarkers were measured using established commercial essays.

#### Endpoints and Follow-up

The total number of all-cause unplanned hospitalizations was selected as the primary endpoint. Cardiovascular admissions were



**Figure 1.** Flowchart for patient inclusion and follow-up. HF, heart failure; LVEF, left ventricular ejection fraction.

selected as the secondary endpoint. Cardiovascular admissions were considered as those occurring due to acute HF, acute coronary syndrome, arrhythmias, stroke, or other cardiovascular causes, such as rupture of an aneurysm, peripheral ischemia, or aortic dissection. Hospitalizations were identified from the clinical records of patients in the HF unit and hospital wards and from electronic medical records. Fatal events were identified from the clinical records of the HF unit, hospital wards, emergency room, and general practitioners and by contacting the patient's relatives. All patients included were followed up until September 2016. The minimum duration of patient follow-up was 3 months.

### Statistical Analysis

Continuous and categorical variables are presented as the mean  $\pm$  standard deviation, median [interquartile range] or percentages, as appropriate. Crude rates (number of events per 10 person-year) are presented for pp-peak  $\text{VO}_2$  quartiles (pp-peak  $\text{VO}_2\text{Q}$ ). Bivariate negative binomial regression was used to assess the independent association between pp-peak  $\text{VO}_2$  with the endpoint.<sup>17</sup> This method simultaneously models the number of admissions with the mortality event and uses a shared frailty to account for the correlation between these 2 outcomes. Thus, the estimates for recurrent admissions are internally adjusted by mortality as a terminal event (informative dropout). To account for differences in the observation time per patient, we used the natural logarithm of the observation time to offset the model for admission endpoints as well as for mortality. Estimates are reported as incidence rate ratio (IRR). All variables listed in Table 1 were evaluated for prognostic purposes. A backward stepwise selection, with the Akaike information criterion as the

stopping criterion, was used to achieve a parsimonious model. The linearity assumption for continuous variables was simultaneously tested and transformed, if appropriate, with fractional polynomials. The pp-peak  $\text{VO}_2$  was modeled using a fractional polynomial transformation of 0 (natural logarithm), per decrease in 10%, and as pp-peak  $\text{VO}_2\text{Q}$ . The covariates included in the final models for all-cause and cardiovascular rehospitalization are included in the legend of Figure 2. A 2-sided *P*-value of  $< .05$  was considered to be statistically significant. All analyses were performed using Stata 14.0.

### RESULTS

The mean  $\pm$  standard deviation of the patients' age was  $72.5 \pm 9.1$  years, 53% were female, 35.1% were in New York Heart Association functional class III, and the median [interquartile range] for N-terminal pro-B-type natriuretic peptide was 1022 pg/mL [363-2109]. The mean  $\pm$  standard deviation and median [interquartile range] for peak  $\text{VO}_2$  and pp-peak  $\text{VO}_2$  were  $10 \pm 2.8$  mL/min/kg and 60 [47-67], respectively. The remaining baseline characteristics of the sample are summarized in Table 1. There was no serious adverse event during cardiopulmonary exercise testing.

The patients were followed up for a median of 276 days [interquartile range, 153-1231]. During the follow-up, 15 deaths (20.3%) and 84 all-cause hospitalizations in 31 patients (41.9%) were registered, distributed as follows: 1 hospitalization (n = 11 [14.9%]), 2 hospitalizations (n = 9 [12.2%]), 3 hospitalizations (n = 4 [5.4%]), 4 hospitalizations (n = 3 [4.1%]), 5 hospitalizations (n = 5 [6.8%]), and 6 hospitalizations (n = 1 [1.4%]). Most

**Table 1**  
Baseline Characteristics of the Study Population

Variables	Included patients (n = 74)
<i>Demographic, medical history, and vital signs</i>	
Age, y	72.5 ± 9.1
Female	39 (53)
Weight, kg	82 ± 15.1
Body surface area, m <sup>2</sup>	1.9 ± 0.2
Hypertension	67 (90.5)
Dyslipidemia	61 (81)
Diabetes mellitus	37 (50)
Ischemic heart disease	27 (36.5)
Baseline NYHA class III/IV	26 (35.1)
Prior admission for AHF	74 (100)
Previous smoker	26 (35.1)
Atrial fibrillation	44 (59.5)
Systolic blood pressure, mmHg	129 ± 16
Heart rate, bpm	69 ± 14
<i>Laboratory</i>	
Hemoglobin, g/d	12.8 ± 1.5
Ferritin, ng/mL	91 [48-175]
eGFR, mL/min/m <sup>2</sup>	58.4 ± 24.3
NT-proBNP, pg/mL	1021 [363-2109]
<i>Echocardiography</i>	
LVEF, %	68.4 ± 9.8
TAPSE, mm <sup>*</sup>	21.8 ± 4.2
LAVI, mL/m <sup>2</sup>	46 [39-58]
LVMi, g/m <sup>2</sup>	125 [104-125]
E/e' ratio	16.9 [13.4-23.4]
PASP, mmHg	46 [39-53]
<i>Exercise performance</i>	
Peak VO <sub>2</sub> , mL/min/kg	10 ± 2.8
pp-peak VO <sub>2</sub> , %	60 (47-67)
VE/VCO <sub>2</sub> slope	35 ± 7.6
RER	1.04 ± 0.1
METs	2.4 ± 0.9
6MWT, m	262 [198-350]
<i>Treatment</i>	
Beta-blockers	59 (79.7)
ACE inhibitors	14 (18.9)
ARB	36 (48.6)
Antialdosterone	21 (28.3)
Loop diuretics	58 (78.4)
Thiazides	22 (29.7)
Digoxin	5 (6.7)

6MWT, 6-minute walk test; ACE, angiotensin-converting enzyme; AHF, acute heart failure; ARB, angiotensin receptor blocker; eGFR, estimated glomerular filtration rate using the Modification of Diet in Renal Disease formula; LAVI, left atrial volume index; LVEF, left ventricular ejection fraction; LVMi, left ventricular mass index; METs, metabolic equivalents; NT-proBNP, N-terminal pro-B-type natriuretic peptide; NYHA, New York Heart Association; PASP, pulmonary artery systolic pressure; peak VO<sub>2</sub>, peak exercise oxygen uptake; pp-peak VO<sub>2</sub>, percentage of predicted peak exercise oxygen uptake; RER, respiratory exchange ratio; TAPSE, tricuspid annular plane systolic excursion; VE/VCO<sub>2</sub> slope, relationship between minute ventilation and the rate of CO<sub>2</sub> elimination; Continuous and categorical variables are presented as mean ± standard deviation, median [interquartile range] or No. (%), as appropriate.

<sup>\*</sup> Data available in 59 patients.

rehospitalizations were due to cardiovascular causes (n = 62 hospitalizations [73.8% of all-causes]) including acute HF as the most frequent among cardiovascular causes (n = 33 hospitalizations [39.3% of all-causes]). The crude rehospitalization rates across pp-peak VO<sub>2</sub> showed a significant and stepwise increase when moving from higher to lower quartiles (P < .01): 2.9, 7.4, 6.2, and 10 hospitalizations per 10 person-year for Q4 (67.7%-102.8%), Q3 (60%-67.5%), Q2 (47%-60%) and Q1 (26.1%-46.4%), respectively. Likewise, increased rates of cardiovascular and acute HF admissions were found among the lower quartiles of pp-peak VO<sub>2</sub> (Figure 2).


### Percentage of Predicted Peak Exercise Oxygen Uptake and Risk of All-cause Recurrent Admissions

In a univariate setting, pp-peak VO<sub>2</sub> was significantly and inversely associated with the risk of recurrent all-cause admissions. Thus, risk estimates attributable to pp-peak VO<sub>2</sub> evaluated as continuous (per decrease in 10%) and as a natural logarithm (per decrease in 1 log) were (IRR, 1.30; 95% confidence interval [95%CI], 1.06-1.61; P = .014 and IRR, 3.94; 95%CI, 1.40-11.05; P = .009), respectively.

In a multivariate scenario, including well-established prognosticators and potential confounders (age, atrial fibrillation, systolic blood pressure, heart rate, E/e' ratio, hemoglobin and N-terminal pro-B-type natriuretic peptide) in the risk model, the inverse association between pp-peak VO<sub>2</sub> and the risk of repeat hospitalizations remained significant (Table 2). The risk gradient showed an inverse and almost linear relationship between pp-peak VO<sub>2</sub> and the risk of all-cause readmissions (Figure 3). When pp-peak VO<sub>2</sub> was modeled as continuous, a 10% decrease of pp-peak VO<sub>2</sub> increased the risk of recurrent hospitalizations by 32% (IRR, 1.32; 95%CI, 1.03-1.68; P = .028) (Table 2). In a sensitivity analysis, forcing in the multivariate analysis other important cardiopulmonary exercise prognosticators such as the VE/VCO<sub>2</sub> slope, ppVO<sub>2</sub> (per decrease of 10%) remained significantly associated with this endpoint (IRR, 1.31; 95%CI, 1.05-1.65; P = .020) and the VE/VCO<sub>2</sub> slope was borderline associated with the risk of recurrent hospitalizations (IRR, 1.03; 95%CI, 0.99-1.06; P = .088).

### Percentage of Predicted Peak Exercise Oxygen Uptake and Risk of Cardiovascular and Acute Heart Failure Hospitalizations

Univariate estimates showed an inverse relationship between pp-peak VO<sub>2</sub> and cardiovascular admissions. The pp-peak VO<sub>2</sub> (per 10% decrease) and logarithm of pp-peak VO<sub>2</sub> (per decrease by 1 log) were inversely associated with a higher risk of cardiovascular hospitalizations (IRR, 1.28; 95%CI, 0.99-1.66; P = .063 and IRR, 3.73; 95%CI, 1.05-13.24; P = .042, respectively). After a multivariate adjustment, this inverse association was strengthened (Table 2). Regarding acute HF hospitalizations, multivariate analysis showed that pp-peak VO<sub>2</sub> (per 10% decrease) and the logarithm of pp-peak VO<sub>2</sub> (per decrease by 1 log) were not significantly related to the risk of repeat acute HF admissions (IRR, 1.10; 95%CI, 0.25-4.73; P = .583 and IRR, 1.55; 95%CI, 0.36-6.69; P = .556, respectively). Similarly, quartiles of pp-peak VO<sub>2</sub> were not related to the risk of repeat acute HF hospitalizations (Table 2).



**Figure 2.** Rates of all-cause, CV and AHF hospitalizations according to quartiles of percentage of predicted peak exercise oxygen uptake. AHF, acute heart failure; CV, cardiovascular.

Q1 = 26.1% to 46.4% of predicted peak exercise oxygen uptake.

Q2 = 47% to 60% of predicted peak exercise oxygen uptake.

Q3 = 60% to 67.5% of predicted peak exercise oxygen uptake.

Q4 = 67.7% to 102.8% of predicted peak exercise oxygen uptake.

**Table 2**

Risk Estimates of All-cause Hospitalizations and Cardiovascular Hospitalizations

	IRR	95%CI	P
<i>All-cause hospitalizations*</i>			
pp-peak VO <sub>2</sub> , per decrease in 10%	1.32	1.03-1.69	.028
Log pp-peak VO <sub>2</sub> , per decrease by 1 log	3.78	1.48-9.69	.006
pp-peak VO <sub>2</sub> quartiles			
Q4 (67.7%-102.8%)	1		
Q3 (60%-67.5%)	1.73	0.89-3.36	.105
Q2 (47%-60%)	2.13	0.92-4.96	.078
Q1 (26.1%-46.4%)	4.04	1.60-10.22	.003
<i>Cardiovascular hospitalizations*</i>			
pp-peak VO <sub>2</sub> , per decrease in 10%	1.42	1.04-1.94	.027
Log pp-peak VO <sub>2</sub> , per decrease in 1 log	5.90	1.28-27.28	.023
pp-peak VO <sub>2</sub> quartiles			
Q4 (67.7%-102.8%)	1		
Q3 (60%-67.5%)	1.40	0.56-3.49	.470
Q2 (47%-60%)	2.27	0.89-5.79	.085
Q1 (26.1%-46.4%)	5.06	1.87-13.67	.001
<i>Acute heart failure hospitalizations*</i>			
Q4 (67.7%-102.8%)	1		
Q3 (60%-67.5%)	0.95	0.20-4.75	.949
Q2 (47%-60%)	0.88	0.19-4.13	.868
Q1 (26.1%-46.4%)	1.82	0.45-7.30	.390


95%CI, 95% confidence interval; hospitalizations; IRR, incidence rate ratio; Log pp-peak VO<sub>2</sub>, logarithm of percentage of predicted peak exercise oxygen uptake; NT-proBNP, N-terminal pro-B-type natriuretic peptide; pp-peak VO<sub>2</sub>, percentage of predicted peak exercise oxygen uptake.

Risk estimates are reported as IRR.

\* Multivariate estimates adjusted for age, atrial fibrillation, systolic blood pressure, heart rate, E/e' ratio, hemoglobin and NT-proBNP.

**DISCUSSION**

Epidemiological studies have shown that the prevalence of HFpEF compared with HF with reduced ejection fraction is increasing over time; however, the prognosis and optimal pharmacological armamentarium for HFpEF is still somber.<sup>4,12</sup> The main finding of our study was that pp-peak VO<sub>2</sub> was independently and linearly associated with recurrent admissions in a cohort of elderly persons with highly symptomatic HFpEF. To our knowledge, this is the first study that has evaluated the prognostic usefulness of pp-peak VO<sub>2</sub> for predicting recurrent hospitalizations in this type of patient.



**Figure 3.** Relationship between pp-peak VO<sub>2</sub> and the risk of all-cause readmission after multivariate adjustment. pp-peak VO<sub>2</sub>, percentage of predicted peak exercise oxygen uptake. Adjusted for age, atrial fibrillation, systolic blood pressure, heart rate, E/e' ratio, hemoglobin and N-terminal pro-B-type natriuretic peptide.

The evidence available to date has shown that patients with HFpEF have markedly reduced functional capacity as objectively measured by peak  $\text{VO}_2$ <sup>18</sup> but little is known about the relationship between pp-peak  $\text{VO}_2$  and prognosis in HFpEF patients. To date, only 3 studies have evaluated the prognostic value (time-to-first cardiac-related mortality or hospitalization) of peak  $\text{VO}_2$  and other cardiopulmonary exercise testing parameters in patients with HF and diastolic dysfunction.<sup>9–11</sup> In 2005, Guazzi et al.<sup>11</sup> studied 46 young male (mean age  $57.9 \pm 13$  years) HF patients with diastolic dysfunction and a left ventricular ejection fraction  $\geq 50\%$ . These authors found that both the peak  $\text{VO}_2$  and the  $\text{VE}/\text{VCO}_2$  slope were significant predictors of mortality and rehospitalization on univariate analysis; however, after multivariate adjustment, only the  $\text{VE}/\text{VCO}_2$  slope remained independently related to adverse prognosis. In 2011, Yan et al.<sup>10</sup> evaluated 224 predominately male (71%) HFpEF patients with a mean age of  $68.8 \pm 9$  years. Similarly, these authors found that plasma B-type natriuretic peptide and  $\text{VE}/\text{VCO}_2$  slope, but not peak  $\text{VO}_2$ , had independent and incremental prognostic value for all-cause and cardiovascular mortality. More recently, in a retrospective study of 173 young and predominately black male HF patients with a left ventricular ejection fraction  $\geq 50\%$ , Shafiq et al.<sup>9</sup> reported that peak  $\text{VO}_2$  and pp-peak  $\text{VO}_2$  had a strong prognostic role in terms of predicting the composite of all-cause mortality and/or cardiac transplant.

In our opinion, previous and current studies endorse the usefulness of cardiopulmonary exercise testing for risk stratification of patients with HFpEF. Discrepancies about the most accurate parameter for risk stratification may be attributed to the small sample size of the studies, differences in baseline characteristics, the covariates included in the multivariate adjustments, and the methodological approach for evaluating the clinical endpoints (recurrent events vs time-to-first event). The main strength of the present study lies in the clinical characteristics of the patients included (elderly, highly comorbid patients with important functional impairment<sup>19</sup>). Last, in our view, analyzing the risk of recurrent hospitalizations in contrast to time-to-first hospitalizations approaches, adds consistency and validity to our findings.

We believe the present findings endorse the clinical value of peak  $\text{VO}_2$  over other traditional, but more subjective, parameters of disease severity such as New York Heart Association functional classification.<sup>20</sup> Furthermore, peak  $\text{VO}_2$  emerges as a reliable and accurate surrogate endpoint to evaluate new therapeutic strategies in HFpEF. For instance, our group recently found a significant increase of peak  $\text{VO}_2$  after a home-based program of inspiratory muscle training in a small study of patients with HFpEF.<sup>21</sup>

### Limitations

This study has some limitations. First, this is a single center observational study in which there may have been many potential confounders; second, the small number of adverse events registered, such as mortality and acute HF hospitalizations, precluded evaluation of the independent association between pp-peak  $\text{VO}_2$  and those events; third, these findings cannot be directly extrapolated to patients with milder forms of the disease; and fourth, the use of a bicycle exercise protocol rather than treadmill exercise testing could have led to underestimation of functional capacity in some patients.<sup>14</sup>

### CONCLUSIONS

In symptomatic elderly patients with HFpEF, pp-peak  $\text{VO}_2$  was significantly and inversely associated with the risk of long-term repeat hospitalizations. Further studies are needed to confirm these results.

### FUNDING

This work was supported in part by grants from: *Sociedad Española de Cardiología: Investigación Clínica en Cardiología*, Grant SEC 2015, CIBER CV 16/11/00420, 16/11/00403, FEDER and PIE15/00013.

### CONFLICTS OF INTEREST

None declared.

#### WHAT IS KNOWN ABOUT THE TOPIC?

- Heart failure with preserved ejection fraction is the most prevalent form of HF.
- Heart failure with preserved ejection fraction is especially common in the aging population, women, frail persons, and highly comorbid patients.
- The clinical value of peak  $\text{VO}_2$  in HFpEF is not well-documented.
- The evidence endorsing the prognostic role of peak  $\text{VO}_2$  in HFpEF is scarce and even absent, especially regarding the risk of recurrent admissions.

#### WHAT DOES THIS STUDY ADD?

- The main finding of our study was that pp-peak  $\text{VO}_2$  was independently and linearly associated with recurrent admissions in a cohort of elderly persons and highly symptomatic HFpEF patients.
- To our knowledge, this is the first study that has evaluated the prognostic usefulness of pp-peak  $\text{VO}_2$  for predicting recurrent hospitalizations in this type of patient.

### REFERENCES

1. Mosterd A, Hoes AW. Clinical epidemiology of heart failure. *Heart*. 2007;93:1137–1146.
2. Lam CS, Donal E, Kraigher-Krainer E, Vasan RS. Epidemiology and clinical course of heart failure with preserved ejection fraction. *Eur J Heart Fail*. 2011;13:18–28.
3. Abbate A, Arena R, Abouzaki N, et al. Heart failure with preserved ejection fraction: refocusing on diastole. *Int J Cardiol*. 2015;179:430–440.
4. Chan MM, Lam CS. How do patients with heart failure with preserved ejection fraction die? *Eur J Heart Fail*. 2013;15:604–613.
5. Moss AJ, Hall WJ, Cannom DS, et al. Cardiac resynchronization therapy for the prevention of heart-failure events. *N Engl J Med*. 2009;361:1329–1338.
6. Anker SD, McMurray JJ. Time to move on from 'time-to-first': should all events be included in the analysis of clinical trials? *Eur Heart J*. 2012;33:2764–2765.
7. Guazzi M, Arena R, Halle M, Piepoli MF, Myers J, Lavie CJ. 2016 Focused update: clinical recommendations for cardiopulmonary exercise testing data assessment in specific patient populations. *Circulation*. 2016;133:e694–e711.
8. Mancini DM, Eisen H, Kussmaul W, Mull R, Edmunds Jr LH, Wilson JR. Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation*. 1991;83:778–786.
9. Shafiq A, Brawner CA, Aldred HA, et al. Prognostic value of cardiopulmonary exercise testing in heart failure with preserved ejection fraction. The Henry Ford Hospital CardioPulmonary EXercise Testing (FIT-CPX) project. *Am Heart J*. 2016;174:167–172.
10. Yan J, Gong SJ, Li L, et al. Combination of B-type natriuretic peptide and minute ventilation/carbon dioxide production slope improves risk stratification in patients with diastolic heart failure. *Int J Cardiol*. 2013;162:193–198.
11. Guazzi M, Myers J, Arena R. Cardiopulmonary exercise testing in the clinical and prognostic assessment of diastolic heart failure. *J Am Coll Cardiol*. 2005;46:1883–1890.
12. McMurray JJ, Adamopoulos S, Anker SD, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2012;33:1787–1847.

13. Anguita Sánchez M, Lambert Rodríguez JL, Bover Freire R, et al. Classification and quality standards of heart failure units: scientific consensus of the Spanish Society of Cardiology. *Rev Esp Cardiol.* 2016;69:940–950.
14. Wasserman K. *Principles of exercise testing and interpretation: including pathophysiology and clinical applications.* Philadelphia: Lippincott Williams & Wilkins; 2005: 160–182.
15. Arena R, Myers J, Aslam SS, Varughese EB, Peberdy MA. Technical considerations related to the minute ventilation/carbon dioxide output slope in patients with heart failure. *Chest.* 2003;124:720–727.
16. Paulus WJ, Tschöpe C, Sanderson JE, et al. How to diagnose diastolic heart failure: a consensus statement on the diagnosis of heart failure with normal left ventricular ejection fraction by the Heart Failure and Echocardiography Associations of the European Society of Cardiology. *Eur Heart J.* 2007;28:2539–2550.
17. Xu X, Hardin JW. Regression models for bivariate count outcomes. *Stata J.* 2016;16:301–315.
18. Haykowsky M, Brubaker P, Kitzman D. Role of physical training in heart failure with preserved ejection fraction. *Curr Heart Fail Rep.* 2012;9:101–106.
19. Enjuanes C, Bruguera J, Grau M, et al. Iron status in chronic heart failure: impact on symptoms, functional class and submaximal exercise capacity. *Rev Esp Cardiol.* 2016;69:247–255.
20. Raphael C, Briscoe C, Davies J, et al. Limitations of the New York Heart Association functional classification system and self-reported walking distances in chronic heart failure. *Heart.* 2007;93:476–482.
21. Palau P, Domínguez E, Núñez E, et al. Effects of inspiratory muscle training in patients with heart failure with preserved ejection fraction. *Eur J Prev Cardiol.* 2014;21:1465–1473.