

Reperfusion Strategies in Hospitals Without Primary Percutaneous Coronary Intervention. Response



Estrategias de reperusión en hospitales sin intervención coronaria percutánea primaria. Respuesta

To the Editor,

We appreciate the comments by Rosell-Ortiz et al. Indeed, half the patients in our study required transfer between centers for primary angioplasty with the associated delay that this entails.¹ Although some randomized studies have shown noninferiority of fibrinolysis therapy in the first few hours after ST-elevation myocardial infarction compared with primary angioplasty,² in general, primary angioplasty is superior, given its greater reperfusion efficacy and lower frequency of complications and bleeding complications in particular.³ Another recent analysis of patients treated according to the *Codi Infart* protocol within 2 hours of infarction also reported that primary angioplasty was superior to fibrinolysis, except when the delay from first medical contact to reperfusion exceeded 140 minutes.⁴

The *Codi Infart* protocol includes administration of fibrinolysis when the delay between first medical contact and reperfusion is expected to be longer than 120 minutes on the basis of availability of a catheterization laboratory, number of ambulances, and traffic in the area. This strategy is also used in other consolidated care systems such as that in Asturias, with a different orography to that in Catalonia, although also with excellent results.⁵ The geography of Catalonia is not complex, which usually allows primary angioplasty within the recommended timeframe. Fibrinolysis (and subsequent transfer to a referral hospital with a catheterization laboratory) is reserved for situations when the timeframe is truly unrealistic (that is, centers a very long way from the referral hospital). This suggests that most delays could be shortened by much earlier diagnosis and drainage after first contact.

Acknowledgments

We thank all the health care professionals involved in the *Codi Infart* program.

Antoni Carol Ruiz,^{a,*} Josep Masip Utset,^b and Albert Ariza-Solé^c

^aServicio de Cardiología, Unidad de Hospitalización, Hospital Moisès Broggi, Consorci Sanitari Integral (CSI), Sant Joan Despí, Barcelona, Spain

^bServicio de Medicina Intensiva, Hospital Moisès Broggi, Consorci Sanitari Integral (CSI), Sant Joan Despí, Barcelona, Spain

^cServicio de Cardiología, Unidad de Cuidados Agudos Cardiológicos, Hospital Universitario de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain

*Corresponding author:

E-mail address: 35955acr@comb.cat (A. Carol Ruiz).

Available online 16 August 2017

REFERENCES

1. Carol Ruiz A, Masip Utset J, Ariza Solé A. Predictors of late reperfusion in STEMI patients undergoing primary angioplasty. Impact of the place of first medical contact. *Rev Esp Cardiol.* 2017;70:162–169.
2. Armstrong P, Gershlick A, Goldstein P, et al. Fibrinolysis or primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med.* 2013;368:1379–1387.
3. Keeley E, Boura J, Gines C. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet.* 2003;361:13–20.
4. Carrillo X, Fernández-Nofreías E, Rodríguez-Leor O, et al. Early ST elevation myocardial infarction in non-capable percutaneous coronary intervention centres: in situ fibrinolysis vs. percutaneous coronary intervention transfer. *Eur Heart J.* 2016;37:1034–1040.
5. Lozano I, Suárez-Cuervo A, Rondán J, et al. Care network for ST-elevation myocardial infarction: what is the ideal catchment area for primary angioplasty? *Rev Esp Cardiol.* 2015;68:444–445.

SEE RELATED CONTENT:

<http://dx.doi.org/10.1016/j.rec.2017.05.031>

<http://dx.doi.org/10.1016/j.rec.2017.06.025>

1885-5857/

© 2017 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.