

which can be fatal in patients with COVID-19. In the meantime, antiplatelet therapy can be prudently administered to critically ill patients with COVID-19 who are at high risk of thromboinflammation but low risk of bleeding. With more studies available in the future, we could then identify the population of patients with COVID-19 most likely to benefit from the use of antiplatelet therapy to prevent the occurrence of arterial or even venous thrombotic events.

Chia Siang Kow^{a,*} and Syed Shahzad Hasan^b

^aSchool of Postgraduate Studies, International Medical University, Kuala Lumpur, Malaysia

^bDepartment of Pharmacy, University of Huddersfield, Huddersfield, United Kingdom

*Corresponding author:

E-mail address: chiasiang_93@hotmail.com (C.S. Kow).

Available online 28 August 2020

REFERENCES

1. Rey JR, Caro-Codón J, Poveda Pineda D, et al. Arterial thrombotic complications in hospitalized patients with COVID-19. *Rev Esp Cardiol.* 2020;73:769–771.
2. Müller C, Karl N, Ziebuhr J, Pleschka S. Dipyridamole and aspirin acetylsalicylate impairs coronavirus replication. *J Antivir Antiretrovir.* 2016;8:142–150.
3. Liu X, Li Z, Liu S, et al. Potential therapeutic effects of dipyridamole in the severely ill patients with COVID-19 [published online ahead of print, 2020 Apr 20]. *Acta Pharm Sin B.* 2020. <http://doi.org/10.1016/j.apsb.2020.04.008>.
4. Ranucci M, Ballotta A, Di Dedda U, et al. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost.* 2020;18:1747–1751.
5. Viecca M, Radovanovic D, Forleo GB, Santus P. Enhanced platelet inhibition treatment improves hypoxemia in patients with severe Covid-19 and hypercoagulability. A case control, proof of concept study. *Pharmacol Res.* 2020;158:104950.

<https://doi.org/10.1016/j.rec.2020.08.009>

1885-5857/

© 2020 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

A new inflammatory-microthrombotic syndrome as an explanation for thrombotic complications in patients with COVID-19



Un nuevo síndrome inflamatorio-microtrombótico como explicación para las complicaciones trombóticas en pacientes con COVID-19

To the Editor,

We have read with great interest the article by King et al.¹ recently published in *Revista Española de Cardiología*. In this article, the authors discuss the higher incidence of thrombotic events in multiple territories and a higher International Society on Thrombosis and Haemostasis (ISTH) disseminated intravascular coagulation (DIC) score in hospitalized patients with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Although the authors mention that they have not ruled out a pre-existing prothrombotic state, our group, consistent with what the authors described,² has proposed a possible role of endothelial injury, complement, and coagulation in the pathogenesis of coronavirus disease 2019 (COVID-19).³

Our pathogenic scheme is based on the similarity of certain clinical and histopathologic findings of various entities that have thrombotic microangiopathy in common with COVID-19, and it postulates that the damage induced by this disease has an endothelial origin with 2 pathogenic routes: an inflammatory route, with predominance of the “cytokine storm” component, and a microangiopathic route involving the complement system.³

Furthermore, endothelial involvement could lead to platelet activation, thus altering coagulation and causing DIC, as described by the authors of the article. This fact *per se* could increase thrombin and prothrombin and trigger complement activation through C5.⁴

For instance, thrombotic microangiopathy has been reported in a patient with severe COVID-19, indicating a pathogenic relationship between these conditions.⁵ The potential role of ADAMTS13 deficiency in serious forms of the disease is another possible factor, as described by Huisman et al.⁶ in a recent article.

Additionally, the inflammatory route itself could activate complement through certain neutrophil serine proteases and macrophages.⁷ Consequently, we believe that there is a strong relationship between inflammation, complement, thrombotic microangiopathy, and coagulation in the pathogenesis of COVID-19. This may represent a new COVID-19–related inflammatory-microthrombotic syndrome which could explain the authors’ interesting findings.^{2–4}

Francisco Valga,^{a,*} Nicanor Vega-Díaz,^a Manuel Macía,^b and José Carlos Rodríguez-Pérez^a

^aServicio de Nefrología, Hospital Universitario de Gran Canaria Doctor Negrín, Las Palmas de Gran Canaria, Las Palmas, Spain

^bServicio de Nefrología, Hospital Universitario Nuestra Señora de la Candelaria, Santa Cruz de Tenerife, Spain

*Corresponding author:

E-mail address: fvalga@hotmail.com (F. Valga).

Available online 2 October 2020

REFERENCES

1. Rey JR, Caro-Codón J, Poveda D, Merino JL, Iniesta AM, López-Sendón JL. Complicaciones arteriales trombóticas en pacientes hospitalizados con COVID-19. *Rev Esp Cardiol.* 2020;73:769–771.
2. Noris M, Benigni A, Remuzzi G. The case of complement activation in COVID-19 multiorgan impact. *Kidney Int.* 2020;98:314–322.
3. Valga F, Vega-Díaz N, Macía M, Monzón T, Rodríguez-Pérez JC. Targeting complement in severe coronavirus disease 2019 to address microthrombosis. *Clin Kidney J.* 2020;13:477–479.
4. Huber-Lang M, Sarma JV, Zetoune FS, et al. Generation of C5a in the absence of C3: a new complement activation pathway. *Nat Med.* 2006;12:682–687.
5. Jhaveri KD, Meir LR, Flores Chang BS, et al. Thrombotic microangiopathy in a patient with COVID-19. *Kidney Int.* 2020;98:509–512.

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2020.05.008>

<https://doi.org/10.1016/j.rec.2020.09.020>

6. Huisman A, Beun R, Sikma M, Westerink J, Kusadasi N. Involvement of ADAMTS13 and von Willebrand factor in thromboembolic events in patients infected with SARS-CoV-2. *Int J Lab Hematol*. 2020. <http://doi.org/10.1111/ijlh.13244>.
7. Huber-Lang M, Younkin EM, Sarma JV, et al. Generation of C5a by phagocytic cells. *Am J Pathol*. 2002;161:1849–1859.

<https://doi.org/10.1016/j.rec.2020.08.021>
1885-5857/

© 2020 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Arterial thrombotic complications in hospitalized patients with COVID-19. Response to related letters



Complicaciones arteriales trombóticas en pacientes hospitalizados con COVID-19. Respuesta a cartas relacionadas

To the Editor,

We appreciate the interest shown by Kow et al. in our work.¹ We fully agree with the comment that the lower cardiovascular risk profile in the cohort of patients with coronavirus disease 2019 (COVID-19), as well as the simultaneous thrombosis in different territories, supports the hypothesis of a systemic prothrombotic state in close relation to the inflammatory response associated with the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)^{1,2}.

Regarding the potential use of prophylactic antiplatelet therapy for its antithrombotic effect and perhaps even, as noted by the authors, its antiviral effect in patients with COVID-19, we recognize that this is an attractive proposal, but currently there is no clear clinical evidence of its usefulness in SARS-CoV-2 infection. There is no doubt about the importance of antiplatelet therapy in patients with arterial thrombotic complications, but its use in patients with high cardiovascular risk without established disease provides minimal benefit and an increased risk of bleeding complications.³ One might think that with COVID-19 this would be different due to the endothelial dysfunction and inflammatory response it causes, but we must avoid empiricism and not support its de novo use in patients with COVID-19 without a specific cardiovascular reason, except in research studies specifically designed to test its efficacy.

We also thank and congratulate Valga et al.⁴ for their recent publication on the role played by endothelial injury, complement, and coagulation in the pathogenesis of coronavirus disease. In our scientific letter,¹ we focused exclusively on the 1.8% (n = 38) of COVID-19 positive patients with arterial thrombotic complications treated at our hospital in March 2020. Although they had a higher

score according to the International Society on Thrombosis and Haemostasis (ISTH) diagnostic criteria for disseminated intravascular coagulation (DIC), only 3 strictly met the diagnostic criteria. As other authors have noted,⁵ it is likely that patients with COVID-19 have a severe hypercoagulability, more so than a consumption coagulopathy, as is the case of classical DIC. Indeed, the pattern is different, as in patients with COVID-19, fibrinogen is characteristically elevated and thrombocytopenia is uncommon, and if it occurs, it is usually mild or moderate. We agree with the hypothesis of Valga et al. of multiple interactions between the immune system, coagulation (immunothrombosis), and associated endothelial dysfunction as a response to SARS-CoV-2 to explain the prothrombotic state of coronavirus disease.

Juan R. Rey,* José Luis Merino, Ángel M. Iniesta, and Juan Caro-Codón, CARD-COVID investigators
Servicio de Cardiología, Hospital Universitario La Paz, Madrid, Spain

*Corresponding author:

E-mail address: juanr.rey@salud.madrid.org (J.R. Rey).

Available online 12 October 2020

REFERENCES

1. Rey JR, Caro-Codón J, Poveda Pineda D, et al. Arterial thrombotic complications in hospitalized patients with COVID-19. *Rev Esp Cardiol*. 2020;73:769–771.
2. Rey JR, Jiménez Valero S, Poveda Pineda D, et al. COVID-19 and simultaneous thrombosis of two coronary arteries. *Rev Esp Cardiol*. 2020;73:676–677.
3. ASCEND Study Collaborative Group, Bowman L, Mafham M, et al. Effects of aspirin for primary prevention in persons with diabetes mellitus. *N Engl J Med*. 2018;379:1529–1539.
4. Valga F, Vega-Díaz N, Macía M, et al. Targeting complement in severe coronavirus disease 2019 to address microthrombosis. *Clin Kidney J*. 2020;13:477–479.
5. Fogarty H, Townsend L, Ni Cheallaigh C, et al. More on COVID-19 coagulopathy in Caucasian patients. *Br J Haematol*. 2020;189:1060–1061.



<https://doi.org/10.1016/j.rec.2020.09.020>
1885-5857/

© 2020 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2020.08.009>

<https://doi.org/10.1016/j.rec.2020.08.021>

Telematic cardiology consultation in the elderly. The 5 M framework can help



Consulta telemática de cardiología para ancianos. La regla de las 5 M puede ser una ayuda

To the Editor,

We read with great interest the excellent consensus document of the Spanish Society of Cardiology on teleconsulta-

tions for clinical cardiologists in the era of COVID-19 by Barrios et al.¹ Telematic cardiology consultations are now a reality in Spain² and a document to help organize them will always be welcome. However, as active members of the Geriatric Cardiology Section, we were disappointed to see that there was no specific reference to elderly patients, who make up a very high percentage of the patients we see in our everyday practice. Elderly patients, who are particularly vulnerable to coronavirus infection,³ need more help to understand that telematic medicine can be an effective way to communicate with their cardiologists and to be able to use it effectively. With this in mind, the 5 M framework^{4,5} (figure 1) can be a useful guide for teleconsultations:

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2020.06.032>

<https://doi.org/10.1016/j.rec.2020.09.022>